

# Small step from single neurons to networks

Johnatan (Yonatan) Aljadeff

*aljadeff@ucsd.edu*

September 22, 2020

# Outline

Information from inside a cell is valuable!

- E/I balance


- Spatial representations

Modeling can help when intracellular recordings are not available

A synapse is not a number

# Information from inside the cell helps us understand **balance of excitation and inhibition**


... during spontaneous activity



Okun, Lampl, 2008


# Information from inside the cell helps us understand **balance of excitation and inhibition**

... and also during evoked activity




Okun, Lampl, 2008

# Information from inside the cell helps us understand **formation of spatial representations**




Harvey et al. 2009

# Information from inside the cell helps us understand formation of spatial representations



Harvey et al. 2009

# Information from inside the cell helps us understand **formation of spatial representations**



Cohen et al. 2017

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$



# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**

- Assume consecutive spikes at times  $t = 0, T$ .

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**

- ▶ Assume consecutive spikes at times  $t = 0, T$ .
- ▶ We know that  $V(t = 0) = 0$ , and  $V(t = T) = \theta$ .

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**

- ▶ Assume consecutive spikes at times  $t = 0, T$ .
- ▶ We know that  $V(t = 0) = 0$ , and  $V(t = T) = \theta$ .
- ▶ Between the two spikes, the dynamics are  $\frac{dV}{dt} = -V + I$ .

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**

- ▶ Assume consecutive spikes at times  $t = 0, T$ .
- ▶ We know that  $V(t = 0) = 0$ , and  $V(t = T) = \theta$ .
- ▶ Between the two spikes, the dynamics are  $\frac{dV}{dt} = -V + I$ .
- ▶ We know the solution!

$$\log\left(\frac{I}{I - V(T)}\right) = \log\left(\frac{I}{I - \theta}\right) = T$$

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**

- ▶ Assume consecutive spikes at times  $t = 0, T$ .
- ▶ We know that  $V(t = 0) = 0$ , and  $V(t = T) = \theta$ .
- ▶ Between the two spikes, the dynamics are  $\frac{dV}{dt} = -V + I$ .
- ▶ We know the solution!

$$\log\left(\frac{I}{I - V(T)}\right) = \log\left(\frac{I}{I - \theta}\right) = T$$

- ▶ So the f-I curve is

$$f(I) = \frac{1}{T}$$

# When intracellular recording impossible, a model can help

To get some intuition, we start with the LIF neuron

$$\begin{aligned}\frac{dV}{dt} &= -V + I, & (\text{with constant input } I) \\ V &= 0 & \text{if } V > \theta\end{aligned}$$

**Let's use what we learned to compute the f-I curve.**


- ▶ Assume consecutive spikes at times  $t = 0, T$ .
- ▶ We know that  $V(t = 0) = 0$ , and  $V(t = T) = \theta$ .
- ▶ Between the two spikes, the dynamics are  $\frac{dV}{dt} = -V + I$ .
- ▶ We know the solution!

$$\log\left(\frac{I}{I - V(T)}\right) = \log\left(\frac{I}{I - \theta}\right) = T$$

- ▶ So the f-I curve is


$$f(I) = \frac{1}{T} = \begin{cases} 0 & I \leq \theta \\ \frac{1}{\log\left(\frac{I}{I - \theta}\right)} & I > \theta \end{cases}$$

# What does the f-I curve tell us?




If the spike frequency is measured extra-cellularly,  
does the f-I curve give us the input?

# What does the f-I curve tell us?




If the spike frequency is measured extra-cellularly, does the f-I curve give us the input?

Not necessarily. What if the input is noisy?






# What does the f-I curve tell us?




If the spike frequency is measured extra-cellularly, does the f-I curve give us the input?


Not necessarily. What if the input is noisy?




# Beyond the f-I curve




# Beyond the f-I curve



# Beyond the f-I curve



# Beyond the f-I curve



In today's exercise we'll explore the f-I curve and the ISI distribution for different types of inputs to a LIF neuron.

# Is a synapse a number?

What is the appropriate way to model the input  $I$  ?

# Is a synapse a number?

What is the appropriate way to model the input  $I$  ?


If a synapse **is** a number,  $I = w \times \text{presynaptic activity}$

# Is a synapse a number?

What is the appropriate way to model the input  $I$  ?

If a synapse **is** a number,  $I = w \times \text{presynaptic activity}$

Alas, **short-term plasticity** (facilitation and depression) exists.



Hubbard, 1963



# Synaptic model of short term plasticity, Tsodyks Markram

Dynamics of synaptic resources:

$$\text{recovered: } \frac{dx}{dt} = \frac{z}{\tau_{\text{rec}}} - U_{\text{SE}} x \delta(t - t_s)$$

$$\text{active: } \frac{dy}{dt} = -\frac{y}{\tau_{\text{in}}} + U_{\text{SE}} x \delta(t - t_s)$$

$$\text{inactive: } \frac{dz}{dt} = \frac{y}{\tau_{\text{in}}} - \frac{z}{\tau_{\text{rec}}}$$

# Synaptic model of short term plasticity, Tsodyks Markram

Dynamics of synaptic resources:

$$\text{recovered: } \frac{dx}{dt} = \frac{z}{\tau_{\text{rec}}} - U_{\text{SE}}^1 x \delta(t - t_s)$$

$$\text{active: } \frac{dy}{dt} = -\frac{y}{\tau_{\text{in}}} + U_{\text{SE}}^1 x \delta(t - t_s)$$

$$\text{inactive: } \frac{dz}{dt} = \frac{y}{\tau_{\text{in}}} - \frac{z}{\tau_{\text{rec}}}$$

# Synaptic model of short term plasticity, Tsodyks Markram

Dynamics of synaptic resources:

$$\text{recovered: } \frac{dx}{dt} = \frac{z}{\tau_{\text{rec}}} - U_{\text{SE}}^1 x \delta(t - t_s)$$


$$\text{active: } \frac{dy}{dt} = -\frac{y}{\tau_{\text{in}}} + U_{\text{SE}}^1 x \delta(t - t_s)$$

$$\text{inactive: } \frac{dz}{dt} = \frac{y}{\tau_{\text{in}}} - \frac{z}{\tau_{\text{rec}}}$$


To model facilitation,  $U_{\text{SE}}$  also changes with time

$$\text{facilitation: } \frac{dU_{\text{SE}}^1}{dt} = -\frac{U_{\text{SE}}^1}{\tau_{\text{facil}}} + U_{\text{SE}}(1 - U_{\text{SE}}^1)\delta(t - t_s)$$


# A toy example where short term plasticity is important




# A toy example where short term plasticity is important



# A toy example where short term plasticity is important



# A toy example where short term plasticity is important



In today's exercise we will explore quantitatively the effects of short term plasticity on the input a neuron receives and on the response of a LIF neuron model.