

GenRCA Rare Codon Analysis Report

Gene Name: YFP

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 582

DNA Sequence:

ATGTTCAAAGGAATCGTAGAAGGCATTGGGATCATTGAAAAGATCGACATTTACACTGAT
CTTGATAAATATGCTATTAGGTTTCCTGAAAATATGCTCAATGGTATCAAAAAAGAGTCT
TCCATCATGTTTAATGGTTGTTTTCTGACTGTGACTTCCGTTAACAGTAATATAGTGTGG
TTTGATATTTTTGAAAAAGAAGCTAGGAAATTGGATAACATTTAGAGAATACAAAGTTGGC
GACAGAGTAAACTTAGGAACATTCCCCAAATTTGGTGCTGCATCTGGTGGTCACATTCTA
AGTGCTAGAATTAGTTGTGTGGCCAGTATTATTGAGATCATCGAAAATGAAGATTATCAA
CAAATGTGGATTCAAATCCCTGAAAACTTTACGGAATTTCTGATTGATAAGGATTATATA
GCCGTCGATGGCATATCTCTAACCATTGACACAATCAAGAATAATCAATTTTTTATATCA
TTACCGTTGAAAATAGCACAAAACACAAACATGAAATGGAGGAAGAAGGGTGACAAAGTT
AACGTGGAACGTAGCAATAAAATTAACGCAAATCAATGCTGG

Protein Sequence:

MFKGIVEGIGIIEKIDIYTDLDKYAIRFPENMLNGIKKESSIMFNGCFLTVTSVNSNIVW
FDIFEKEARKLDTFREYKVGDRVNLGTFPKFGAASGGHILSARISCVASIIIEIENEDYQ
QMWIQIPENFTEFLIDKDYIAVDGISLTIDTIKNNQFFISLPLKIAQNTNMKWRKKGDKV
NVELSNKINANQCW



Negative CIS Elements	Negative repeat Elements
1	0

Indices based on non-uniform usage of synonymous codon

Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.28	[11]
ENC	Effective Number of Codons	20~61	53.87	[2, 3]
RCBS	Relative Codon Bias Strength	≥ 0	0.25	[41]
DCBS	Directional Codon Bias Score	≥ 1	1.67	[51]
CDC	Codon Deviation Coefficient	0~1	0.1	[61]
MILC	Measure Independent of Length and Composition	-1~1	-0.52	[71]
ICDI	Intrinsic Codon Deviation Index	0~1	0.26	[81]
SCUO	Synonymous Codon Usage Order	0~1	0.18	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.77	[111]
P	Codon Preference	≥ 1	1.12	[121]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.33	[131]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.75	[11]
CFD	Codon Frequency Distribution	0~1	0.02	[21]
FOP	Frequency of Optimal Codons	0~1	0.48	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.57	[51]
COUSIN18			0.35	
CBI	Codon Bias Index	-1~1	0.18	[61]
Dmean	Mean Dissimilarity-based Index	0~2	0.48	[71]
RCA	Relative Codon Adaptation	≥ 0	1.09	[81]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.34	[91]
B	Codon Usage Bias	0~2	0.35	[101]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.5	[11]
gtAI	Genetic tRNA Adaptation Index	0~1	0.6	[21]
P2	P2 Index	0~1	0.43	[31]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
	GC Content at the Third Position			

GC3	of Synonymous Codons	0~1	0.36	[1]
GC	GC Content		0.35	
GC1	GC Content at the First Position of Synonymous Codons		0.38	
GC2	GC Content at the Second Position of Synonymous Codons		0.3	
ENcp	Effective Number of Codon Pairs	20~61	20	[2]
CPS	Codon Pair Score	-1~1	0.03	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.79	[5]

Gene Name: Beta-galactosidase

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)

DNA Sequence Length: 2994


DNA Sequence:

```
ATGACCATGATAACGGATAGCTTAGCCGTTGTTCTCCAACGAAGAGACTGGGAAAATCCT
GGTGTGACCCAACCTTAATAGACTAGCTGCTCATCCTCCATTTGCCTCATGGAGAACTCT
GAAGAGGCGAGAAGCTGATAGGCCATCGCAACAACCTTCGGTCCCTCAACGGAGAGTGGAGA
TTTGCCTGGTTTCCAGCTCCTGAAGCGGTGCCAGAGTCTTGGTTGGAATGTGATCTTCCA
GAAGCTGACACAGTTGTCGTCCTCCATCCAACCTGGCAGATGCACGGATATGATGCTCCTATC
TACACCAATGTCACCTTATCCGATCACAGTCAATCCTCCTTTTGTTCCTACTGAAAACCCA
ACCGGCTGCTACTCCCTGACATTCAATGTTGACGAATCTTGGCTGCAAGAGGGACAGACG
AGAATAATTTTCGACGGTGTCAACTCTGCATTCCATTTATGGTGTAAATGGAAGATGGGTT
GGTTATGGTCAAGACAGTAGATTACCTTCTGAGTTCGATCTGAGCGCATTTCTTAGAGCT
GGTGAGAACCGTTTAGCAGTTATGGTGCTAAGATGGTCTGATGGTTCCTATTTAGAGGAT
CAAGACATGTGGAGGATGTCTGGGATTTTCCGTGATGTTTCGTTGCTCCACAAGCCAACT
ACTCAAATTTTCAGACTTCCATGTTGCTACCAGATTCAATGATGATTTTCAGCAGAGCAGTT
TTAGAGGCTGAAGTTCAAATGTGTGGAGAATTAAGAGATTACCTTCGTGTTACTGTTTCA
CTATGGCAAGGTGAAACTCAAGTGGCATCTGGTACCGCTCCTTTTGGCGGTGAAATTATT
GACGAGCGTGGGGGTTACGCTGATAGAGTTACTCTCAGATTAAATGTTGAAAACCCAAAG
TTATGGTCTGCTGAAATTCCAAACTTGTATCGTGCTGTTGTGGAGTTGCACACCGCAGAT
GGAACCTTTGATTGAGGCTGAAGCTTGTGATGTTGGGTTCCGTGAGGTCCGTATTGAAAAC
GGGTTGCTTTTGTCTAAACGGCAAACCTCTTTTGATTGCTGGCGTTAATCGTCACGAACAT
CACCTCTTCATGGACAAGTTATGGACGAGCAAACCATGGTCCAAGATATTCTATTGATG
AAACAAAACAACCTTCAATGCTGTCCGCTGCTCGCATTACCTAATCATCCCTTATGGTAC
ACTCTCTGTGACCGTTATGGGTTGTACGTTGTTGATGAGGCTAACATTGAGACTCATGGA
ATGGTACCTATGAACAGATTGACAGACGACCCAAGATGGCTTCCAGCTATGAGTGAAAGA
GTTACCAGAATGGTGCAGAGAGACAGAAATCACCCATCTGTCATTATCTGGTCTCTTGGA
AATGAATCTGGTCATGGTGCCAACCACGATGCTCTTTACAGATGGATCAAGTCGGTTGAT
CCATCTAGACCTGTTCAATATGAAGGTGGTGGCGCAGATACTACTGCTACTGACATTATC
TGCCCAATGTACGCCCGTGTGACGAGGACCAGCCGTTTCCAGCTGTTCCAAAATGGAGT
ATCAAAAAATGGCTGTCCCTTCCAGGTGAAAACCAGACCACTAATTTTATGCGAGTACGCC
CATGCTATGGGTAACTCCCTGGGTGGATTTGCCAAGTACTGGCAAGCATTCCGTCAATAC
CCAAGATTGCAGGGTGGTTTTGTTTGGGATTGGGTTGATCAATCTTTGATCAAGTACGAC
GAGAATGGAACCCCTTGGTCTGCCTATGGTGGAGACTTTGGTGATACTCCAAACGACAGA
CAGTTCTGTATGAACGGTCTAGTCTTTGCTGACAGGACTCCACACCCAGCCTTAACCTGAG
GCCAAGCATCAACAACAGTTTTTCCAATTTAGACTATCTGGTCAAACCATCGAAGTTACG
TCCGAATACCTGTTTCGACATTCTGACAATGAGCTGCTACATTGGATGGTTGCGCTAGAT
GGTAAGCCACTAGCTTCAGGCGAAGTGCCCTGGACGTTGCTCCTCAGGGGAAGCAACTA
ATTGAGCTTCCTGAATTGCCACAGCCTGAGTCTGCTGGTCAACTCTGGTTGACTGTACGT
GTTGTTTCAGCCCAACGCTACTGCTTGGTCCGAGGCGGGTCATATATCGGCCTGGCAGCAA
TGGAGGCTGGCTGAAAATTTGTGCGGTAACATTACCTGCTGCCTCCCATGCCATTCCACAT
TTGACTACAAGTGAGATGGACTTTTGTATCGAGTTGGGTAACAAGAGGTGGCAATTCAAC
AGACAATCTGGTTTCCTTAGTCAAATGTGGATTGGGGATAAAAAGCAACTTTTGACCCCT
CTAAGGGATCAGTTCACTAGAGCTCCACTTGATAACGACATTGGTGTTTCTGAAGCTACA
AGAATAGACCCAAATGCCTGGGTTGAAAGATGGAAGGCGGCTGGTCAATTATCAGGCTGAG
GCTGCTCTACTTCAGTGTACTGCAGATACGTTAGCTGATGCCGTTCTAATTACCACAGCC
```

CATGCTTGGCAGCATCAAGGAAAGACTCTGTTTATTTCCAGAAAAACCTATCGAATCGAT
GGATCGGGACAGATGGCTATCACCGTTGATGTTGAAGTTGCTAGCGATACCCACATCCA
GCCAGAATAGGTCTTAAGTGTCTAGCTCAAGTGGCTGAGAGAGTCAACTGGTTAGGA
TTAGGACCTCAGGAGAACTATCCCGACAGATTGACCGCAGCTTGCTTTGACAGATGGGAC
CTTCCATTATCCGATATGTACACTCCATATGTGTTTCCTTCTGAGAACGGACTGCGATGC
GGTACCAGAGAGTTGAACTACGGGCCTCACCAATGGAGAGGTGATTTCCAATTCAACATC
TCAAGATACAGTCAACAACAGTTGATGGAAACAAGTCACAGACATTTGTTGCATGCAGAA
GAGGGTACTTGGCTCAACATCGATGGGTTTCACATGGGAATTGGTGGAGACGAT

Protein Sequence:

MTMITDSLAVVLQRRDWENPGVTQLNRLAAHPPFASWRNSEEARTDRPSQQLRSLNGEWR
FAWFPAPEAVPESWLECDLPEADTVVPSNWQMHGYDAPIYTNVTYPITVNPPFVPTENP
TGCYSLTFNVDESWLQEGQTRII FDGVNSAFHLWCNCRWVGYGQDSRLPSEFDLSAFLRA
GENRLAVMVLRWSDGSYLEDQDMWRMSGIFRDVSLHKKPTTQISDFHVATRFNDDFSRAV
LEAEVQMCGELRDYLRVTVSLWQGETQVASGTAPFGGEI I DERGGYADRVTLRLNVENPK
LWSAEIPNLRYAVVELHTADGTLIEAEACDVGFREVRIENGLLLLNGKPLLIRGVNRHEH
HPLHGQVMDEQTMVQDILLMKQNNFNAVRCSHYPNHPLWYTLCDRYGLYVVDEANIETHG
MVPNMRLTDDPRWLPAMSERVTRMVQRDRNHPSVIIWSLGNESGHGANHDALYRWIKSVD
PSRPVQYEGGGADTTATDIICPMYARVDEDQFFPAVPKWSIKKWLSLPGETRPLILCEYA
HAMGNSLGGFAKYWQAFRQYPRLQGGFVWDWVDQSLIKYDENGNPWSAYGGDFGDTPNDR
QFCMNLVFADRTPHPALTEAKHQQQFFQFRLSGQTIEVTSEYLFRRHSDNELLHWMVALD
GKPLASGEVPLDVAPQGKQLIELPELPQPESAGQLWLTVRVVPNATAWSEAGHISAWQQ
WRLAENLSVTLPAASHAIPHLTTSEMDFCIELGNKRWFNRQSGFLSQMWIGDKKQLLTP
LRDQFTRAPLDNDIGVSEATRDPNAWVERWKAAGHYQAEAAALLQCTADTLADAVLITTA
HAWQHQGKTLFISRKTYRIDGSGQMAITVDVEVASDTPHPARIGLNCQLAQVAERNWL
LGPQENYPDLRLTAACFDRWDLPLSDMYTPYVFPSENGLRCTRELNYGPHQWRGDFQFNI
SRYSQQQLMETSHRHLHLHAEEGTWNLDGFHMGIGGDD



Negative CIS Elements	Negative repeat Elements
0	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.28	[11]
ENC	Effective Number of Codons	20~61	46.3	[2, 3]

RCBS	Relative Codon Bias Strength	≥ 0	0.25	[4]
DCBS	Directional Codon Bias Score	≥ 1	1.77	[5]
CDC	Codon Deviation Coefficient	0~1	0.13	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.38	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.11	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.11	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.82	[11]
P	Codon Preference	≥ 1	1.13	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.66	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.76	[1]
CFD	Codon Frequency Distribution	0~1	0.03	[2]
FOP	Frequency of Optimal Codons	0~1	0.52	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	1.55	[5]
COUSIN18			1.87	
CBI	Codon Bias Index	-1~1	0.29	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.09	[7]
RCA	Relative Codon Adaptation	≥ 0	1.13	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.12	[9]
B	Codon Usage Bias	0~2	0.2	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.52	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.59	[2]
P2	P2 Index	0~1	0.66	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.4	[1]
GC	GC Content		0.46	
GC1	GC Content at the First Position of Synonymous Codons		0.55	

GC2	GC Content at the Second Position of Synonymous Codons		0.43	
ENcp	Effective Number of Codon Pairs	20~61	35.62	[2]
CPS	Codon Pair Score	-1~1	-0.01	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.76	[5]

Gene Name: Cre

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 1029

DNA Sequence:

```
ATGTCCAATCTTTTGACTGTCCATCAAAATCTTCCTGCTCTACCTGTTGATGCAACCTCT
GACGAAGTCAGGAAGAATCTAATGGATATGTTTCAGAGACCGACAAGCATTTTCAGAGCAT
ACTTGGAAGATGCTGTTGAGCGTTTGTCGTTTCATGGGCAGCGTGGTGCAAACCTCAATAAC
CGTAAGTGGTTCCCCGCTGAACCTGAGGATGTGAGAGATTATCTATTGTATTTACAGGCT
CGTGGGCTGGCCGTTAAGACGATACAGCAACACTTAGGGCAGCTAAATATGTTGCATAGA
CGTTCCGGGCTACCTAGACCTTCAGATTCCAATGCAGTGTCTCTGGTCATGCGTCGGATA
CGTAAGGAGAATGTGGACGCCGGTGAACGAGCCAAGCAAGCTTTAGCGTTCGAAAGAACA
GACTTTGACCAAGTGAGAAGTCTCATGGAAAACCTCAGATCGATGTCAAGATATTAGGAAC
CTGGCATTCTTGGAATCGCTTACAACACTTTACTTCGTATCGCTGAAATTGCCAGGATC
AGGGTGAAAGATATTTCCAGGACTGATGGGGGCAGAATGTTGATTCATATTGGCCGAACA
AAAACGCTGGTTTCCACAGCAGGTGTAGAAAAGGCACTGTCGTTAGGAGTAACCAAACCTC
GTTGAAAGATGGATATCCGTTTCTGGTGTGCTGACGACCCTAACAACTATCTGTTTTGT
CGAGTCAGAAAAAATGGAGTGGCAGCTCCATCAGCTACTTCCCAGCTCAGTACCAGAGCT
CTGGAGGGTATCTTTGAGGCCACTCACAGGTTGATCTATGGTGCCAAGGATGATAGTGGT
CAAAGATATCTGGCCTGGTCTGGACACAGCGCCCGGGTGGTGCTGCCCCGTGACATGGCA
AGAGCGGGGGTTTCCATACCAGAAATTATGCAAGCTGGCGGGCTGGACGAATGTTAATATT
GTCATGAACATATACGGAATCTTGATTTCAGAGACTGGTGCCATGGTAAGACTACTAGAA
GATGGAGAC
```

Protein Sequence:

```
MSNLLTVHQNLPALPVDATSDVRKLNLMDFRDRQAFSEHTWKMLLSVCRSWAAWCKLNN
RKWFPAEPEDVRDYLLYLQARGLAVKTIQQHLGQLNMLHRRSGLPRPSDSNAVSLVMRRI
RKENVDAGERAKQALAFERTDFDQVRSLMENS DRCQDIRNLAFLGIAYNTLLRIAIEIARI
RVKDISRTDGGRM LIHIGRTKTLVSTAGVEKALSLGVTKLVERWISVSGVADDPNNYLC
RVRKNGVAAPSATSQLSTRALEGIFEATHRLIYGAKDDSGQRYLAWSGHSARVGAARDMA
RAGVSIPEIMQAGGWTNVNIVMNYIRNLDSETGAMVRLLEDGD
```



Negative CIS Elements	Negative repeat Elements
0	0

Indices based on non-uniform usage of synonymous codon

Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.13	[11]
ENC	Effective Number of Codons	20~61	55.81	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.17	[4]
DCBS	Directional Codon Bias Score	≥1	1.54	[5]
CDC	Codon Deviation Coefficient	0~1	0.09	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.52	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.1	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.07	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.87	[11]
P	Codon Preference	≥1	1.08	[12]
MCB	Maximum-likelihood Codon Bias	≥0	0.25	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.66	[1]
CFD	Codon Frequency Distribution	0~1	0.07	[2]
FOP	Frequency of Optimal Codons	0~1	0.4	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.19	[5]
COUSIN18			-0.07	
CBI	Codon Bias Index	-1~1	0.12	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.31	[7]
RCA	Relative Codon Adaptation	≥0	1.06	[8]
CUFS	Codon Usage Frequency Similarity	0~√2	0.18	[9]
B	Codon Usage Bias	0~2	0.25	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.47	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.55	[2]
P2	P2 Index	0~1	0.54	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.43	[1]
GC	GC Content		0.47	
GC1	GC Content at the First Position of Synonymous Codons		0.54	
GC2	GC Content at the Second Position of Synonymous Codons		0.44	
ENcp	Effective Number of Codon Pairs	20~61	61	[2]
CPS	Codon Pair Score	-1~1	0.05	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.74	[5]

Gene Name: Luciferase

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)


DNA Sequence Length: 1650

DNA Sequence:

```
ATGGAAGACGCCAAAAATATCAAGAAAGGTCCAGCCCCGTTTTATCCACTAGAGGATGGT
ACCGCCGGTGAACAACCTCCATAAGGCCATGAAGCGTTATGCCCTGGTTCCTGGAACATTT
GCTTTCACTGATGCACATATCGAGGTCAACATTACATATGCTGAATATTTTGAGATGTCC
GTCAGATTAGCTGAAGCAATGAAGAGGTACGGTTTGAACACCAATCATCGTATTGTTGTC
TGTTCCGAAAACAGTTTACAATTTTTTATGCCCGGTGTTGGGTGCATTATTTATTGGTGTT
GCTGTTGCTCCGGCTAACGATATTTACAATGAAAGAGAGTTGCTGAATAGTATGAACATC
TCACAGCCCACTGTCTGTTTTGTCTCAAAGAAAGGCTTACAAAAGATCTTAAATGTCCAG
AAGAACTCCCCATTATCCAAAAAATCATCATCATGGATTGCAAACTGATTACCAAGGG
TTCCAGTCAATGTATACGTTTGTCACTAGCCATCTGCCTCCGGGGTTAATGAGTATGAC
TTTGTTCCAGAATCTTTTGACAGAGATAAGACAATTGCCCTGATTATGAATTCTTCAGGT
TCTACAGGATTGCCCAAGGGTGTGCTTTGCCACATAGAACGGCTTGTGTCCGTTTCTCT
CACGCAAGAGATCCCATTTTTTGGGAACCAGATAATACCAGACACTGCCATCTTATCAGTG
GTTCCCTTCCACCACGGATTTGGGATGTTTACAACCTTGGGGTACTTGATCTGTGGATTC
CGAGTTGTTCTCATGTATAGGTTCTGAAGAGGAACTGTTTTTAAGATCGCTGCAAGACTAC
AAAATTCAAAGTGCCTACTTGTTCCTCACTTTATTTTTCTTTTTTTGCCAAAAGTACGTTG
ATTGACAAGTATGACTTGTCCAATCTGCATGAAATCGCAAGTGGAGGTGCTCCGTTAAGT
AAAGAAGTTGGCGAAGCTGTAGCTAAGCGTTTCCATTTGCCCTGGTATTCGTCAAGGTTAC
GGTTTAACGGAAACCACTAGTGCCATTTTAATAACACCAGAGGGCGACGACAAACCTGGA
GCCGTCGGTAAAGTTGTTCCCTTCTTTGAAGCCAAGGTAGTAGACTTGGACACTGGTAAA
ACATTGGGAGTGAATCAGAGAGGAGAGCTTTGTGTTGTTGTTGTTGTTGTTGTTGTTGTT
TACGTTAATAATCCTGAGGCTACTAATGCTTTAATTGACAAAGATGGTTGGTTACATTCT
GGTGACATTGCTTATTGGGATGAAGATGAGCATTCTTTATTGTCGATCGTCTTAAGTCC
TTGATTAAGTATAAGGGATATCAAGTGGCCCCAGCAGAATTGGAATCAATTTTGCTTCAA
CATCCTAATATATTTGACGCTGGTGTGCTGGATTGCCAGATGATGATGCTGGCGAACTC
CCAGCAGCAGTGGTAGTCTTAGAACATGGAAGAACTATGACGGAGAAAGAGATTGTTGAC
TACGTCGCTTCGCAAGTTACCACTGCAAAAGAACTGAGAGGTGGGGTTGTCTTTGTTCGAT
GAAGTTCCAAAAGGATTAACAGGTAAGCTTGATGCTAGAAAAATAAGAGAAATCTTAATT
AAAGCTAAAAAGGGTGGTAAATCCAACTG
```

Protein Sequence:

```
MEDAKNIKKGPAPFYPLEDGTAGEQLHKAMKRYALVPGTIAFTDAHIEVNIITYAEYFEMS
VRLAEAMKRYGLNTNHRIVVCSSENSLQFFMPVLGALFIGVAVAPANDIYNERELLNSMNI
SQPTVVFVSKKGLQKILNVQKKLP I IQK I IIMDSKTDYQGFQSMYTFVTSHLPPGFNEYD
FVPESFDRDKTIALIMNSSGSTGLPKGVALPHRTACVRFSHARDPIFGNQIIPDTAILS
VVFHHGFGMFTTLGYLICGFRVVMYRFEELFLRSLQDYKIQSALLVPTLFSFFAKSTL
IDKYDLSNLHEIASGGAPLSKEVGEAVAKRFHLPGRQGYGLTETTSAILITPEGDDKPG
AVGKVVPPFEAKVVDLDTGKTLGVNQRGELCVRGPMIMSGYVNNPEATNALIDKDWLHS
GDIAYWDEDEHFFIVDRLKSLIKYQVAPAELESILLQHPNIFDAGVAGLPDDDAGEL
PAAVVVLEHGKTMTEKEIVDYVASQVTTAKKLRGGVVVFVDEVKGLTGKLDARKIREILI
KAKKGGKSKL
```



Negative CIS Elements	Negative repeat Elements
2	0

Indices based on non-uniform usage of synonymous codon

Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.23	[11]
ENC	Effective Number of Codons	20~61	48.44	[2, 3]
RCBS	Relative Codon Bias Strength	≥ 0	0.18	[4]
DCBS	Directional Codon Bias Score	≥ 1	1.55	[5]
CDC	Codon Deviation Coefficient	0~1	0.1	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.44	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.16	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.11	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.86	[11]
P	Codon Preference	≥ 1	1.09	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.4	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.74	[11]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.48	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.67	[5]
COUSIN18			0.62	
CBI	Codon Bias Index	-1~1	0.22	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.2	[7]

RCA	Relative Codon Adaptation	≥ 0	1.1	[8]
CUFS	Codon Usage Frequency Similarity	$0 \sim \sqrt{2}$	0.19	[9]
B	Codon Usage Bias	$0 \sim 2$	0.22	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	$0 \sim 1$	0.51	[1]
gtAI	Genetic tRNA Adaptation Index	$0 \sim 1$	0.6	[2]
P2	P2 Index	$0 \sim 1$	0.5	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	$0 \sim 1$	0.38	[1]
GC	GC Content		0.42	
GC1	GC Content at the First Position of Synonymous Codons		0.51	
GC2	GC Content at the Second Position of Synonymous Codons		0.36	
ENcp	Effective Number of Codon Pairs	$20 \sim 61$	35.07	[2]
CPS	Codon Pair Score	$-1 \sim 1$	0.01	[3] , [4]
Codon Volatility	Codon Volatility	$0.5 \sim 1$	0.76	[5]

Gene Name: lacI transcription factor

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 1080

DNA Sequence:

ATGAAGCCAGTCACTCTCTATGACGTCGCCGAATATGCTGGAGTGTCTATCAGACAGTG
TCCAGAGTGGTGAACCAAGCCTCGCATGTTAGCGCGAAGACGCGTGAGAAGGTTGAAGCT
GCCATGGCAGAGCTGAACTACATTCCAAACAGAGTTGCTCAACAACCTAGCGGGTAAACAG
TCCCTTCTTATTGGTGTGGCAACATCTTCGCTGGCTTTGCATGCTCCTTCCCAGATTGTT
GCTGCCATTAAGAGCAGGGCTGACCAACTGGGTGCTTCGGTGGTAGTTTCCATGGTGGAG
AGATCCGGTGTTGAGGCCTGTAAGGCTGCGGTCCACAACCTGTTAGCTCAGAGAGTGTCA
GGGCTGATCATCAATTATCCTTTGGACGATCAGGATGCTATTGCAGTTGAGGCTGCCTGT
ACCAATGTTCCGGCTTTGTTCTTGGATGTCAGTGACCAGACTCCTATCAACTCAATCATC
TTTTCTCATGAAGACGGTACGAGGCTGGGAGTGGAACACCTGGTTGCACTAGGACACCAG
CAAATTGCTCTGCTAGCTGGCCCTCTGAGTTCGGTTAGTGCCAGACTGAGATTAGCTGGC
TGGCACAAGTACTTGACCAGAAATCAAATCCAGCCAATCGCCGAACGTGAGGGAGACTGG
TCCGCTATGTCCGGTTTCCAGCAAATATGCAAATGCTTAACGAAGGAATCGTCCCCACT
GCAATGTTGGTTGCAAACGACCAAATGGCCCTGGGTGCCATGAGAGCTATCACTGAATCT
GGTCTCAGGGTTGGTGCTGATATCTCCGTCGTTGGCTACGATGACACCGAAGATTCTTCG
TGTTACATTCTCCTCTCAACCACTATCAAACAAGACTTTCGTTTATTAGGACAGACCTCG
GTTGACAGATTGTTGCAATTGAGTCAAGGTCAAGCTGTTAAGGGTAACCAACTTCTACCA
GTTTCCCTTGTTAAGAGAAAACTACTCTTGCTCCCAATACTCAAACCTGCTTCCCAAGA
GCATTGCTGATTCCCTAATGCAATTAGCCAGACAGGTTTCCCGTTTGAATCCGGTCAA

Protein Sequence:

MKPVTLYDVAEYAGVSYQTVSRVVNQASHVSAKTREKVEAAMAELNYIPNRVAQQLAGKQ
SLIIGVATSSLALHAPSQIVAAIKSRADQLGASVVVSMVERSGVEACKAAVHNLLAQRVS
GLIINYPLDDQDAIAVEAACTNPALFLDVSDQTPINSIIFSHEDGTRLGVEHLVALGHQ
QIALLAGPLSSVSARLRLAGWHKYLTRNQIQPIAEREGDWSAMSGFQQTMQMLNEGIVPT
AMLVANDQMALGAMRAITESGLRVGADISVVGYYDDTEDSSCYIPPLTTIKQDFRLLGQTS
VDRLLQLSQGQAVKGNQLLPVSLVKRKTTLAPNTQTASPRALADSLMQLARQVSRLESGQ



Negative CIS Elements	Negative repeat Elements
0	0

Indices based on non-uniform usage of synonymous codon

Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.38	[11]
ENC	Effective Number of Codons	20~61	46.11	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.31	[4]
DCBS	Directional Codon Bias Score	≥1	1.89	[5]
CDC	Codon Deviation Coefficient	0~1	0.18	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.4	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.17	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.15	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.81	[11]
P	Codon Preference	≥1	1.2	[12]
MCB	Maximum-likelihood Codon Bias	≥0	0.59	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.72	[1]
CFD	Codon Frequency Distribution	0~1	0.03	[2]
FOP	Frequency of Optimal Codons	0~1	0.47	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	1.42	[5]
COUSIN18			1.8	
CBI	Codon Bias Index	-1~1	0.23	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.21	[7]
RCA	Relative Codon Adaptation	≥0	1.12	[8]
CUFS	Codon Usage Frequency Similarity	0~√2	0.23	[9]
B	Codon Usage Bias	0~2	0.28	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.52	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.58	[2]
P2	P2 Index	0~1	0.68	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.47	[1]
GC	GC Content		0.49	
GC1	GC Content at the First Position of Synonymous Codons		0.59	
GC2	GC Content at the Second Position of Synonymous Codons		0.43	
ENcp	Effective Number of Codon Pairs	20~61	32.37	[2]
CPS	Codon Pair Score	-1~1	-0.03	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.74	[5]

Gene Name: TetR transcription factor

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 654

DNA Sequence:

ATGGCTAGACTAAATAGAGAATCGGTGATAGATGCAGCACTTGAACCTCTAAACGAAACT
GGTATCGATGGGCTAACTACAAGAAAAGTGGCACAGAAATTAGGAATTGAGCAACCAACA
CTGTATTGGCACGTTAAAAATAAGAGAGCACTGTTAGATGCCCTTGCCGTGGAAATTCTT
GCCAGACATCATGACTATTTCCTTACCAGCTGCAGGAGAAAGTTGGCAATCTTTCTTGAGA
AACAAATGCGATGTCGTTCCGTAGAGCACTGTTAAGGTATAGAGATGGTGCAAAGGTCCAT
CTGGGAACAAGACCCGATGAGAAACAATACGACACAGTCGAAACACAATAAGGTTTCATG
ACTGAAAACGGGTTTTCTCTGAGAGATGGGCTCTATGCTATAAGTGCTGTCAGCCATTTT
ACTTTGGGAGCTGTACTGGAACAGCAGGAACATACTGCAGCATTAACAGATCGACCTGCT
GCTCCAGATGAAAACCTTGCCCTCCACTCTTGAGAGAGGCATTACAAATTATGGATAGTGAT
GATGGCGAACAGGCTTTTCTTCATGGATTGGAGAGTTTGATACGAGGCTTTGAAGTCCAA
TTGACAGCATTGCTTCAAATTGTAGGAGGCGATAAGCTGATAATACCATTCTGT

Protein Sequence:

MARLNRESVIDAALELLNETGIDGLTTRKLAQKLGIEQPTLYWHVKNKRALLDALAVEIL
ARHHDYSLPAAGESWQSFRLNNAMSFRRALLRYRDGAKVHLGTRPDEKQYDTVETQLRFM
TENGFSLRDGLYAISAVSHFTLGAVLEQQEHTAALTDSPAAPDENLPPLLREALQIMSDS
DGEQAFHLGLESIRGFVQLTALLQIVGGDKLIIPFC



Negative CIS Elements	Negative repeat Elements
0	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference

RSCU	Relative Synonymous Codon Usage	0~6	1.42	[11]
ENC	Effective Number of Codons	20~61	47.14	[2, 3]
RCBS	Relative Codon Bias Strength	≥ 0	0.3	[4]
DCBS	Directional Codon Bias Score	≥ 1	1.75	[5]
CDC	Codon Deviation Coefficient	0~1	0.16	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.45	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.24	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.19	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.78	[11]
P	Codon Preference	≥ 1	1.17	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.53	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.69	[11]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.45	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	1.06	[5]
COUSIN18			0.91	
CBI	Codon Bias Index	-1~1	0.2	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.46	[7]
RCA	Relative Codon Adaptation	≥ 0	1.11	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.31	[9]
B	Codon Usage Bias	0~2	0.42	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.45	[11]
gtAI	Genetic tRNA Adaptation Index	0~1	0.55	[2]
P2	P2 Index	0~1	0.51	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.33	[11]
GC	GC Content		0.43	

GC1	GC Content at the First Position of Synonymous Codons		0.57	
GC2	GC Content at the Second Position of Synonymous Codons		0.39	
ENcp	Effective Number of Codon Pairs	20~61	32.85	[2]
CPS	Codon Pair Score	-1~1	0.07	[3] , [4]
Codon Volatility	Codon Volatility	0.5~1	0.75	[5]

Gene Name: araC transcription factor

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 876

DNA Sequence:

ATGCCGAAGCTCAAAATGATCCACTGCTTCCAGGCTATTCTTTCAATGCTCATTGTTGGTT
GCCGGTTTAACCCCTATTGAAGCAAATGGATATCTGGATTTTTTTCATAGACAGGCCATTG
GGGATGAAAGGTTATATTTTGAACCTGACCATCAGAGGTCAAGGGGTTGTGAAAAACCAA
GGCAGAGAATTCGTATGCCGTCCTGGTGACATCCTGCTGTTTCCTCCTGGAGAAATCCAT
CATTATGGGCGTCACCCTGAGGCCAGGGAATGGTACCACCAATGGGTGTATTTTAGACCC
AGAGCTTACTGGCATGAATGGCTGAATTGGCCATCAATATTTGCAAATACTGGCTTTTTTC
CGACCAGACGAGGCTCACCAACCTCACTTCTCTGACCTATTTGGACAGATCATCAACGCT
GGACAAGGGGAGGGAAGATATTCTGAGTTATTGGCCATCAATTTGCTGGAGCAACTGCTG
CTACGGAGAATGGAAGCAATTAACGAATCATTGCATCCACCGATGGACAATAGAGTTCGG
GAAGCTTGCCAATATATATCCGATCATTTGGCTGACTCCAATTTTGATATAGCCTCAGTG
GCTCAACATGTTTGTCTTAGTCCCAGTAGACTATCTCATTTGTTTCAGACAGCAGCTGGGT
ATCTCCGTACTGAGTTGGAGAGAGGATCAACGTATAAGTCAAGCAAAGTTGCTTTTGTCA
ACCACCCGTATGCCTATAGCTACGGTAGGAAGAAACGTTGGATTTGATGACCAATTATAT
TTCAGTCGTGTCTTTAAGAAATGCACTGGAGCCTCTCCTAGCGAGTTTAGAGCTGGCTGT
GAAGAGAAAGTGAATGATGTTGCCGTCAGCTGTCC

Protein Sequence:

MAEAQNDPLLPGYSFNAHLVAGLTPIEANGYLDFFDIDRPLGMKGYILNLTIRGQGVVKNQ
GREFVCRPGDILLFPPGEIHYYGRHPEAREWYHQWVYFRPRAYWHEWLNWPSIFANTGFF
RPDEAHQPHFSDLFGQIINAGQGEGRYSELLAINLLEQLLLRRMEAINESLHPPMDNRVR
EACQYISDHLADSNDIASVAQHVCLSPSRLSHLFRQQLGISVLSWREDQRISQAKLLLS
TTRMPIATVGRNVGFDDQLYFSRVFKKCTGASPSEFRAGCEEKVNDVAVKLS



Negative CIS Elements	Negative repeat Elements
1	0

Indices based on non-uniform usage of synonymous codon

Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.29	[11]
ENC	Effective Number of Codons	20~61	51.22	[2, 3]
RCBS	Relative Codon Bias Strength	≥ 0	0.28	[4]
DCBS	Directional Codon Bias Score	≥ 1	1.71	[5]
CDC	Codon Deviation Coefficient	0~1	0.14	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.46	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.12	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.13	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.83	[11]
P	Codon Preference	≥ 1	1.18	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.45	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.72	[11]
CFD	Codon Frequency Distribution	0~1	0.03	[2]
FOP	Frequency of Optimal Codons	0~1	0.42	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.07	[5]
COUSIN18			-0.37	
CBI	Codon Bias Index	-1~1	0.13	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.33	[7]
RCA	Relative Codon Adaptation	≥ 0	1.12	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.23	[9]
B	Codon Usage Bias	0~2	0.31	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.49	[11]
gtAI	Genetic tRNA Adaptation Index	0~1	0.57	[2]
P2	P2 Index	0~1	0.53	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
-------	-------------	-------	-------	-----------

GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.41	[1]
GC	GC Content		0.45	
GC1	GC Content at the First Position of Synonymous Codons		0.54	
GC2	GC Content at the Second Position of Synonymous Codons		0.4	
ENcp	Effective Number of Codon Pairs	20~61	31.72	[2]
CPS	Codon Pair Score	-1~1	0.03	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.76	[5]

Gene Name: Regulatory protein GAL4

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)

DNA Sequence Length: 2643


DNA Sequence:

```
ATGAAATTACTATCCTCTATAGAACAGGCTTGTGATATTTGTCGATTGAAGAAGCTTAAG
TGTTCCAAGGAGAAACCAAAGTGTGCAAAATGTCTCAAAAACAACTGGGAATGTCGTTAC
TCGCCCCAAGACTAAAAGATCGCCCCCTAACTAGAGCGCATCTGACGGAGGTTGAGTCTAGG
CTAGAACGCTTGGAGCAGCTATTTTTTGTGATATTTCCAAGAGAAGATTTGGATATGATT
CTTAAATGGATTCACTTCAAGATATCAAGGCTCTTCTAACGGGACTTTTCGTTTCAGGAC
AATGTCAACAAAGATGCAGTAACGGATCGTTTGGCATCTGTTGAAACCGATATGCCTTTG
ACCCTAAGACAGCACCGTATTTCTGCAACATCAAGTTCGGAAGAGTCCTCTAACAAAGGA
CAAAGGCAGCTTACAGTATCCATTGATTCCGCAGCCCATCATGACAATTCAACAATCCCC
CTCGACTTTATGCCTAGAGATGCTCTCCATGGATTTGATTGGTCCGAAGAAGACGATATG
TCTGACGGTTTGGCATTTTTGAAAACCTGATCCTAATAACAACGGTTTTTTTTGGGGATGGC
AGTTTGTATGTATCCTACGATCCATTGGTTTCAAACCTGAAAATTATACAAATTCAAAT
GTGAATAGGCTTCCAACCATGATCACTGATAGGTATACACTCGCAAGTCGAAGTACCACT
TCCCGGCTTTTACAAAGCTATCTCAATAACTTCCATCCATACTGTCCAATAGTGCCTCC
CCAACCTTTAATGATGCTCTACAACAATCAGATTGAAATTGCCTCTAAAGACCAGTGGCAG
ATACTATTCAACTGCATCCTAGCCATTGGGGCATGGTGCATCGAAGGTGAATCTACAGAC
ATTGACGTTTTTCTATTATCAAAATGCAAAGTCACACCTAACATCCAAGGTGTTTGAATCG
GGATCTATCATACTTGTTACGGCATTGCATCTTCTTTCTAGATATACTCAATGGCGACAA
AAAATAACACATCTTACAACCTCCATTCACTTAGCATACGAATGGCCATTTCACTAGGT
TTGAATAGGGACCTTCCTTCGAGTTTCTCTGACTCGTCGATTTTGGAGCAAAGACGAAGA
ATTTGGTGGTCTGTGTATTCTCGGAAATCCAACATCATTACTATATGGAAGGAGCATT
CAGTTGAGCCAAAATACCATTTTCGTTCCATCTTCGGTGGATGATGTTTCAGCGGACAAC
ACAGGACCAACAATTTATCATGGAATTATTGAAACAGCAAGACTACTTCAAGTGTTCCAC
AAGATCTACGAGCTTGACAAAACCGTTACCGCTGAGAAGAGCCCTATATGTGCCAAGAAA
TGTTTAATGATCTGCAACGAAATCGAAGAAGTTTCCCGACAAGCTCCGAAGTTTTTACAG
ATGGATATAAGCAACCACTGCCCTGACCAACCTTTTGAAAAGAACACCCCTGGCTGAGTTT
ACGAGATTTGAGCTTAAATGGAAGCAGCTTTCACTGATAATCTACGTGCTCAGAGATTT
TTCACGAATTTACGCAAAAAGAGAGTCAGCTGGAACAAGATCAGAATGATCATCAATCA
TACGAAGTTAAGAGATGTTCAATAATGTTATCAGATGCAGCTCAACGAAGTGTGATGTCC
GTATCATCGTATATGGACAATCATAATGTTACACCATACTTTGCATGGAATTGTTTCATAT
TATTTATTCAATGCAGTGTTAGTCCCCATCAAGACATTATTAAGCAATTCTAAAAGCAAT
GCTGAAAATAACGAGACAGCTCAGCTTTTGCAACAGATAAATACTGTACTGATGCTTCTA
AAAAAGTTGGCTACATTCAAGATCCAACTTGTGAAAAGTATATTCAGGTGTTGGAAGAG
GTGTGTGCACCTTTTTTGCTTTCGCAATGTGCGATCCCATACCTCACATTAGCTACAAT
AACTCTAATGGAAGTGCCATCAAAAACATTGTGGGATCAGCCACGATTGCTCAGTATCCA
ACTTTACCAGAAGAAAATGTAAACAACATATCTGTTAAGTACGTTAGCCCTGGCTCGGTG
GGGCCTTCCCCAGTTCCATTGAAATCAGGTGCCTCATTTTCAGATTTGGTCAAACATATTA
TCCAACAGACCTCCAAGTAGAACTCCCCGGTTACTATTTCCAGATCGACCCCAAGCCAT
CGATCGGTGACTCCATTTCTGGGCCAACAGCAGCAGCTACAATCCCTTGTTCCATTGACA
CCTTCAGCACTTTTTGGTGGTGCAAACTTTAATCAATCTGGGAATATAGCCGATTCTCTCT
TTATCATTACGTTTACAACTCTAGCAATGGACCAAAATCTTATTACAACCCAGACAAAC
TCACAGGCTTTATCACAACCGATTGCTTCATCAAACGTTTCACGATAACTTCATGAATAAC
GAAATTACTGCTTCCAAGATAGACGATGGGAATAATTCTAAGCCGCTGAGCCCAGGCTGG
```

ACTGATCAAAC TGCCTATAATGCATTTGGTATTACGACGGGAATGTTCAATACGACTACT
ATGGACGATGTTTATAACTATTTATTTCGATGATGAAGATACCCCGCCCAATCCGAAGAAA
GAG

Protein Sequence:

MKLLSSIEQACDICRLKKLKCSKEKPKCAKCLKNNWECRYSPKTKRSPLTRAHLTEVESR
LERLEQLFLLIFPREDLDMILKMDSLQDIKALLTGLFVQDNVNKDAVTDRLASVETDMPL
TLRQHRISATSSSESSNKQQRQLTVSIDSAAHHDNSTIPLDFMPRDALHGFDWSEEDDM
SDGLPFLKTDPNNGFFGDSLLCILRSIGFKPENYTNSNVNRLPTMITDRYTLASRSTT
SRLQSYLNNFHPYCPIVHSPTLMMLYNNQIEIASKDQWQILFNCILAIGAWCIEGESTD
IDVFYYQNAKSHLTSKVFEESGSIILVTALHLLSRYTQWRQKNTSYNFHSFSIRMAISLG
LNRDLPSFSDSSILEQRRRIWWSVYSWEIQLSLLYGRSIQLSQNTISFPSSVDDVQRTT
TGPTIYHGIIETARLLQVFTKIYELDKTVTAEKSPICAKKCLMICNEIEEVSQRQAPKFLQ
MDISTTALTNLLKEHPWLSFTRFELKWKQLSLIIYVLRDFFTNFTQKKSQLEQDQNDHQS
YEVKRCSIMLSDAQRTVMSVSSYMDNHNVTPTYFAWNCSYYLEFNAVLVPIKTLLSNSKSN
AENNETAQLLQQINTVLMMLKKLATFKIQTCCKYIQVLEEVCAPFLLSQCAIPLPHISYN
NSNGSAIKNIVGSATIAQYPTLPEENVNINISVKYVSPGSGVPSVPLKSGASFSDLVKLL
SNRPPSRNSPVTIPRSTPSHRSVTPFLGQQQQQLQSLVPLTPSALFGGANFNQSGNIADSS
LSFTFTNSSNGPNLITTQNTSQALSQPIASSNVHDNFMNNEITASKIDDGNNSKPLSPGW
TDQTAYNAFGITTMGFNTTTMDDVYNYLFDDEDTPPNPKKE



Negative CIS Elements	Negative repeat Elements
2	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.1	[11]
ENC	Effective Number of Codons	20~61	53.75	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.13	[41]
DCBS	Directional Codon Bias Score	≥1	1.39	[51]
CDC	Codon Deviation Coefficient	0~1	0.07	[61]
MILC	Measure Independent of Length and Composition	-1~1	-0.47	[71]

ICDI	Intrinsic Codon Deviation Index	0~1	0.08	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.06	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.91	[11]
P	Codon Preference	≥ 1	1.04	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.24	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.7	[1]
CFD	Codon Frequency Distribution	0~1	0.05	[2]
FOP	Frequency of Optimal Codons	0~1	0.42	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.41	[5]
COUSIN18			0.57	
CBI	Codon Bias Index	-1~1	0.13	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.25	[7]
RCA	Relative Codon Adaptation	≥ 0	1.08	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.14	[9]
B	Codon Usage Bias	0~2	0.23	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.46	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.54	[2]
P2	P2 Index	0~1	0.47	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.4	[1]
GC	GC Content		0.41	
GC1	GC Content at the First Position of Synonymous Codons		0.43	
GC2	GC Content at the Second Position of Synonymous Codons		0.4	
ENcp	Effective Number of Codon Pairs	20~61	40.37	[2]
CPS	Codon Pair Score	-1~1	0.05	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.76	[5]

Gene Name: Ribonuclease Barnase

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)


DNA Sequence Length: 471

DNA Sequence:

```
ATGATGAAGATGGAAGGCATAGCGCTCAAAAAAAGGCTAAGTTGGATATCAGTCTGTCTC
CTGGTCCTTGTCAGTGCCGCTGGTATGCTTTTTTCCACAGCAGCAAAGACAGAAACAAGT
TCACACAAAGCGCATACAGAGGCACAGGTGATAAACACATTTGACGGTGTTGCCGACTAC
CTACAAACATACCATAAACTACCTGATAATTATATTACCAAATCAGAGGCTCAAGCCCTC
GGTTGGGTTGCTAGTAAAGGTAACCTGGCAGATGTTGCCCTGGCAAGTCTATCGGAGGC
GATATCTTCTCCAACAGAGAAGGAAAATTGCCAGGAAAATCAGGTCGTACTTGGAGGGAG
GCTGATATTAACTACACGTCTGGGTTTCAGAAATAGTGATAGAATCTTATACTCGTCTGAT
TGGTTGATCTACAAGACTACTGATCATTATCAAACCTTTCACCAAGATTAGA
```

Protein Sequence:

```
MMKMEGIALKKRLSWISVCLLVLSAAGMLFSTA AKTETSSHKAHTAQVINTFDGVADY
LQTYHKLPDNYITKSEAQALGWVASKGNLADVAPGKSIGGDIFSNREGKLPKSGRTWRE
ADINYTSGFRNSDRILYSSDWLIYKTTDHYQTFTKIR
```



Negative CIS Elements	Negative repeat Elements
2	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.23	[11]
ENC	Effective Number of Codons	20~61	56.83	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.25	[41]

DCBS	Directional Codon Bias Score	≥ 1	1.57	[5]
CDC	Codon Deviation Coefficient	0~1	0.12	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.6	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.19	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.13	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.82	[11]
P	Codon Preference	≥ 1	1.13	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.26	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.7	[1]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.42	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	1.06	[5]
COUSIN18			1.74	
CBI	Codon Bias Index	-1~1	0.13	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.42	[7]
RCA	Relative Codon Adaptation	≥ 0	1.08	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.27	[9]
B	Codon Usage Bias	0~2	0.36	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.47	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.57	[2]
P2	P2 Index	0~1	0.64	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.41	[1]
GC	GC Content		0.43	
GC1	GC Content at the First Position of Synonymous Codons		0.45	
GC2	GC Content at the Second Position of Synonymous Codons		0.44	

ENcp	Effective Number of Codon Pairs	20~61	23.26	[2]
CPS	Codon Pair Score	-1~1	0.05	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.76	[5]

Gene Name: Cas9

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)

DNA Sequence Length: 2994


DNA Sequence:

```
ATGGACAAAAAGTACTCCATCGGTCTAGACATTGGAACAAACTCAGTCGGGTGGGCCGTT
ATTACCGACGAGTACAAGGTTCCCGAGTAAGAAGTTCAAAGTATTGGGGAATACAGACCGT
CACTCTATCAAAAAAGAACTTAATTGGCGCACTATTGTTTGATAGTGGGGAAACTGCCGAA
GCAACAAGATTGAAGAGAAGTGC CGCTCGTAGGTATACTAGAAGGAAAAACCGTATTTGT
TATTTG CAGGAGATATTTTCTAATGAAATGGCTAAAGTTGACGATTCTTCTTCCACCGA
CTAGAAGAATCTTTTCTTGTGAGGAGGACAAAAACACGAACGACACCCTATTTTGGT
AACATCGTTGATGAAGTTGCCTACCATGAGAAATATCCGACGATCTACCATTTGCGAAAG
AAACTGGTTGATTCAACCGATAAAGCTGACTTGAGATTGATATACCTTGCTTTAGCTCAT
ATGATTAAATTTAGAGGCCATTTCTTGATCGAAGGAGATCTCAATCCTGACAATAGTGAT
GTGGATAAGTTATTTATCCAACCTGGTTCAAACCTACAATCAGCTCTTTGAGGAGAACCCC
ATAAATGCCTCTGGTGTGGATGCTAAGGCAATTTTGAGTGCCAGATTGAGCAAATCCAGA
AGACTTGAAAATCTGATTGCACAGCTTCCAGGCGAAAAAGAAATGGCCTATTTGGGAAT
CTGATAGCACTCTCATTTGGGTCTTACTCCAAACTTCAAATCCAACCTTTGATTTAGCGGAA
GACGCGAAAACCTTCAACTAAGTAAAGACACTTATGATGATGATTTGGATAATTTGCTGGCT
CAAATTGGTGATCAGTATGCCGATTTGTTTTTAGCTGCCAAGAACTTGAGTGATGCTATT
TTGCTTTCTGATATTCTGAGAGTTAACTGAAATTACGAAAGCTCCCTTGTCTGCTTCA
ATGATCAAAAGATATGATGAGCACCACCAAGACCTAACTTTGCTAAAAGCTTTGGTGCGT
CAACAACCTACCAGAAAAATACAAGGAGATTTTCTTTGACCAGTCGAAAAATGGTTACGCT
GGCTATATCGATGGAGGAGCCTCCCAAGAGGAATTTTACAAGTTTATCAAACCAATTCTA
GAAAAAATGGATGGTACCGAAGAGTTATTGGTTAAGCTTAACAGGGAAGACCTTTTGAGG
AAGCAAAGGACTTTTGACAATGGTTCAATTCCCCACCAGATTCATTTAGGTGAATTACAT
GCGATATTGAGGCGCCAGGAGGATTTTATCCGTTTTTGAAAGACAATCGTGAAAAGATC
GAAAAGATTTTGACTTTTTCGTATTTCCTTACTATGTTGGACCTTTAGCCAGAGGAAATAGC
AGATTGCGCTGGATGACCAGAAAATCTGAAGAGACGATCACTCCTTGGAACCTTTGAAGAA
GTCGTGGACAAAGGAGCATCCGCCCAATCATTTATCGAGAGGATGACCAATTTTGATAAG
AATCTACCCAATGAAAAGGTATTACCCAAACACTCTTTGTTGTATGAGTATTTCACTGTG
TATAACGAGTTGACTAAAGTGAAATATGTTACTGAAGGTATGCGCAAGCCTGCTTTTCTC
TCCGGGGAACAGAAGAAGGCCATCGTTGATCTTTTGTTCAAAACCTAATCGTAAAGTTACT
GTCAAGCAGCTGAAGGAAGATTACTTTAAGAAGATTGAATGTTTTGACAGCGTTGAAATC
AGCGGAGTTGAAGACCGTTTTAATGCGTCGTTAGGAACCTACCATGACCTTTTGAAAATA
ATAAAAGATAAAGACTTTCTGGATAATGAGGAAAACGAAGACATATTAGAAGATATTGTA
CTGACCCTAACATTGTTTGAGGATAGAGAAATGATTGAAGAGCGTCTGAAGACCTATGCT
CATCTCTTTGATGACAAAGTCATGAAACAGCTGAAGAGACGAAGATATACAGGCTGGGGA
AGATTAAGTCGAAAGCTAATTAATGGTATCCGGGATAAGCAGAGTGGTAAGACAATTCTG
GACTTCTTAAAATCTGATGGGTTTGCTAACAGAACTTCATGCAGTTGATTCATGATGAT
TCATTGACGTTCAAAGAGGATATACAGAAGGCTCAAGTGTCAAGGTCAAGGTGACTCTTG
CATGAGCATATTGCTAATCTAGCCGGGTCAACAGCCATAAAAAAGGGTATTCTTCAAACCT
GTCAAAGTTGTGGATGAGTTGGTCAAGGTGATGGGAAGGCACAAAACCCGAAAAATATAGTC
ATCGAAATGGCACGAGAAAAATCAAACCACTCAAAAAAGGTCAAAAAAACTCTAGAGAAAGA
ATGAAAAGAATTGAAGAAGGGATAAAAAGAGCTTGGATCGCAAAATCTGAAAGAACACCCA
GTGGAAGAACACCCAACCTGCAAAACGAGAAATGTACTTGTATTATTTACAAAATGGACGT
GATATGTATGTGGATCAGGAATTAGACATTAACAGGTTGAGTGACTATGATGTTGATCAC
```

ATTGTTCTCAAAGTTTTTTGAAGGATGATTCAATTGATAACAAAGTGCTAACAAGATCA
GACAAGAATAGAGGAAAGTCTGACAATGTACCTTCGGAAGAGGTCGTCAAGAAGATGAAG
AACTATTGGAGACAATTGCTGAATGCAAAGCTAATTACCCAGAGAAAAGTTTGATAATCTT
ACAAAAGCTGAAAGAGGAGGACTATCGGAGTTGGATAAGGCTGGGTTTATTAAAGACAG
TTAGTAGAACTAGACAGATCACTAAGCACGTTGCTCAGATTCTGGATTCCAGAATGAAC
ACCAAGTACGATGAAAATGATAAATTGATAAGAGAAAGTCAAGGTTATCACTTTGAAGAGT
AAATTGGTATCAGATTTTAGAAAAGATTTTCAGTTCTACAAGGTAAGAGAAATCAACAAC
TACCACCATGCCATGATGCCTACCTCAATGCTGTCTGTTGGTACTGCTTTGATA

Protein Sequence:

MDKKYSIGLDIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAE
ATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSFHRLEESFLVEEDKKHERHPIFG
NIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSD
VDKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGN
LIALSLGLTPNFKSNFDLAEDAKLQLSKDITYDDDLNLLAQIGDQYADLFLAAKNLSDAI
LLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYA
GYIDGGASQEEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELH
AILRRQEDFYFPFLKDNREKIEKILTFRIPIYVGPLARGNSRFWMTRKSEETITPWNFEE
VVDKGASQSFIERMTNFDKNLPNEKVLPHSLLYEYFTVYNELTKVKYVTEGMRKPAFL
SGEQKKAIVDLLFKTNRKVTQVQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLKI
IKDKDFLDNEENEDILEDIVLTTLTFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWG
RLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMLIHDDSLTFKEDIQKAQVSGQGDSL
HEHIANLAGSPAIIKKGILQTVKVVDLKVVMGRHKPENIVIEMARENQTTQKGQKNSRER
MKRIEEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYDVDH
IVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRKFDNL
TKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKS
KLVSDFRKDFQFYKVREINNYHHAHDAYLNAVVGTAII



Negative CIS Elements	Negative repeat Elements
2	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.18	[11]
ENC	Effective Number of Codons	20~61	50.33	[2, 3]

RCBS	Relative Codon Bias Strength	≥ 0	0.17	[4]
DCBS	Directional Codon Bias Score	≥ 1	1.52	[5]
CDC	Codon Deviation Coefficient	0~1	0.09	[6]
MILC	Measure Independent of Length and Composition	-1~1	-0.45	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.12	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.07	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.91	[11]
P	Codon Preference	≥ 1	1.06	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.36	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.75	[1]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.48	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.78	[5]
COUSIN18			0.88	
CBI	Codon Bias Index	-1~1	0.2	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.15	[7]
RCA	Relative Codon Adaptation	≥ 0	1.12	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.16	[9]
B	Codon Usage Bias	0~2	0.15	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.52	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.6	[2]
P2	P2 Index	0~1	0.47	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.41	[1]
GC	GC Content		0.39	
GC1	GC Content at the First Position of Synonymous Codons		0.47	

GC2	GC Content at the Second Position of Synonymous Codons		0.29	
ENcp	Effective Number of Codon Pairs	20~61	38.51	[2]
CPS	Codon Pair Score	-1~1	0.03	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.78	[5]

Gene Name: Toxin CcdB

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: Pichia pastoris (budding yeasts)


DNA Sequence Length: 303

DNA Sequence:

ATGCAATTCAAAGTTTATACGTACAAGAGAGAGAGCAGATATCGATTGTTTCGTCGATGTT
CAATCGGATATCATTGACACACCAGGACGTAGAATGGTGATCCCTCTGGCTAGTGCAAGA
TTGCTTAGTGACAAAGTTTCAAGAGAGTTGTATCCAGTCGTCCATATTGGAGACGAATCC
TGGAGAATGATGACTACAGACATGGCATCAGTCCCAGTTAGCGTCATTGGGGAAGAGGTA
GCAGACCTGTCTCATAGAGAAAACGATATCAAGAACGCAATTAACCTCATGTTTTGGGGT
ATT

Protein Sequence:

MQFKVYTYKRESRYRLFVDVQSDIIDTPGRRMVIPLASARLLSDKVSRELYPVVHIGDES
WRMMTTDMASVPVSVIGEEVADLSHRENDIKNAINLMFWGI



Negative CIS Elements	Negative repeat Elements
0	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.6	[11]
ENC	Effective Number of Codons	20~61	43.46	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.55	[41]
DCBS	Directional Codon Bias Score	≥1	1.89	[51]
CDC	Codon Deviation Coefficient	0~1	0.22	[61]

MILC	Measure Independent of Length and Composition	-1~1	-0.54	[7]
ICDI	Intrinsic Codon Deviation Index	0~1	0.33	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.28	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.64	[11]
P	Codon Preference	≥ 1	1.37	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.59	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.73	[1]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.46	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	1.24	[5]
COUSIN18			2.32	
CBI	Codon Bias Index	-1~1	0.21	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.5	[7]
RCA	Relative Codon Adaptation	≥ 0	1.11	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.4	[9]
B	Codon Usage Bias	0~2	0.5	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.53	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.61	[2]
P2	P2 Index	0~1	0.59	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.45	[1]
GC	GC Content		0.43	
GC1	GC Content at the First Position of Synonymous Codons		0.48	
GC2	GC Content at the Second Position of Synonymous Codons		0.37	
ENcp	Effective Number of Codon Pairs	20~61	20	[2]
CPS	Codon Pair Score	-1~1	-0.02	[3, 4]

Codon Volatility	Codon Volatility	0.5~1	0.76	[5]
---------------------	------------------	-------	------	---------------------

Gene Name: T7 RNA Polymerase

Reference Source: Codon Usage Database - Kazusa

Expression Host Organism: *Pichia pastoris* (budding yeasts)

DNA Sequence Length: 2649


DNA Sequence:

```
ATGAATACTATCAACATAGCCAAGAATGACTTTTCTGACATCGAACTTGCGGCCATTCCA
TTCAACACCCTTGCCGATCACTATGGAGAAAAGACTAGCAAGGGAGCAGTTGGCATTAGAA
CACGAGTCTACGAAATGGGGGAGGCCCGCTTCAGGAAAAATGTTTGAAAGGCAATTAAAA
GCAGGAGAAGTAGCCGATAATGCCGCTGCTAAGCCGTTGATTACAACGTTACTTCCAAAA
ATGATTGCAAGGATCAACGACTGGTTTGAAGAAGTGAAAGCTAAACGAGGTAAGAGACCC
ACTGCGTTTCAATTTCTCCAAGAAATTAAGCCAGAAGCAGTTGCATATATTACAATAAAG
ACAACGCTAGCGTGTCTCACCTCTGCCGATAATACCACGGTGCAGGCAGTTGCGTCTGCT
ATAGGGCGTGCCATTGAAGATGAGGCTCGTTTTGGCAGGATTCGAGACTTGAGGGCAAAG
CACTTCAAAAAAACGTTGAAGAACAATAAACAAACGCGTAGGTCATGTCTATAAAAAAG
GCATTTCATGCAAGTGGTTGAAGCTGACATGCTCAGCAAGGGACTTTTAGGAGGGGAGGCA
TGGTCTTCATGGCATAAGGAGGATAGCATACACGTTGGAGTACGATGTATTGAAATGCTA
ATTGAGTCTACAGGAATGGTGTCACTTCATAGACAAAAATGCTGGAGTTGTCCGTCAGAT
TCAGAGACCATTGAACTCGCACCTGAGTATGCCGAGGCCATTGCCACTAGAGCTGGAGCT
CTAGCCGGTATATCCCCAATGTTCCAACCCTGTGTCTGCTACCTCCTAAACCATGGACTGGT
ATTACTGGTGGTGGCTATTGGGCAAATGGAAGAAGACCACTAGCATTGGTAAGAACTCAT
TCTAAAAAGGCCTTAATGAGATACGAAGATGTTTACATGCCAGAAGTTTATAAAGCCATT
AATATTGCTCAAAACACCGCCTGGAAAATCAACAAGAAAGTATTAGCCGTTGCCAATGTT
ATCACTAAATGGAAGCATTGTCCAGTAGAAGATATTCCAGCAATTGAGAGGGAGGAATTA
CCAATGAAGCCCGAAGACATCGACATGAACCCCGAAGCCCTAACCGCGTGGAAGAGAGCT
GCTGCTGCTGTTTATCGCAAGGATAAGGCTCGTAAATCAAGAAGAATCTCCTTGAGTTT
ATGTTAGAGCAAGCCAATAAATTTGCCAATCACAAGGCTATATGGTTTCCATATAACATG
GATTGGAGAGGCCGTGTATATGCAGTTTCTATGTTCAATCCTCAAGGAAATGATATGACC
AAGGGATTGTAACTCTTGCAAAGGGGAAAACCCATTGGAAAAGAAGGTTACTACTGGTTG
AAAATTTCATGGAGCCAACGTGTCCGGAGTGGAACAAGGTTCCGTTTCCAGAACGAATCAAG
TTTATTGAAGAAAATCATGAGAACATTATGGCCTGTGCTAAATCTCCTTTAGAAAACACA
TGGTGGGCTGAACAGGATTACCATTTTGTTCCTTGCTTTCTGTTTTGAGTATGCAGGT
GTGCAGCATCATGGTTTGAGTTACAACGTAGTTTGGCCCTGGCGTTTGATGGATCCTGC
TCTGGTATTTCAGCACTTTTCTGCTATGCTTCGAGATGAGGTTGGAGGAAGAGCTGTTAAC
TTACTGCCTAGCGAGACAGTGCAGGATATCTACGGTATTGTTGCAAAGAAAGTCAACGAA
ATTTTGCAAGCAGATGCGATCAATGGGACAGACAATGAGGTTGTACAGTCACGGATGAA
AATACGGGGGAAATCAGTGAGAAAGTAAAGCTGGGTACCAAAGCATTAGCCGGACAATGG
CTAGCTTACGGTGTGACTAGATCGGTCACCAAAAGATCAGTGATGACGTTGGCCTATGGA
TCAAAGGAATTTGGGTTTAGACAACAAGTTCTTGAAGATACAATCCAGCCAGCAATTGAC
TCCGGTAAAGGGCTCATGTTCACTCAACCTAATCAAGCTGCAGGTTACATGGCCAAGCTA
ATTTGGGAGTCTGTTTCTGTTACAGTTGTTGCAGCTGTAGAGGCTATGAATTGGCTTAAG
AGCGCTGCAAACTATTGGCTGCCGAGGTAAGGACAAAGAAGACAGGTGAAATTCTAAGA
AAAAGATGTGCTGTTTCAATTGGGTGACTCCTGACGGTTTCCGGTTTGGCAGGAATACAAG
AAACCCATTCAAACAGATTGAACCTTATGTTTTTGGGTCAATTTAGACTTCAACCAACT
ATCAATACAAACAAAGACTCTGAAATTGATGCTCACAACAAAGAAATCAGGAATTGCTCCC
AACTTTGTTCACTCCCAGGATGGTAGTCATCTGAGAAAAGACAGTCGTTTGGGCTCACGAA
AAATATGGTATCGAGAGTTTTGCTCTGATCCATGACTCCTTTGGAACATTCTCTGCTGAT
GCTGCCAATTTGTTCAAAGCTGTACGAGAGACCATGGTTGATACCTACGAATCCTGTGAT
```

GTCTTGGCTGATTCTACGACCAATTTGCTGATCAATTACATGAATCTCAGTTGGATAAA
ATGCCTGCACTACCTGCAAAGGGTAATCTGAATCTTAGAGACATTCTGAATCAGATTTT
GCATTCGCT

Protein Sequence:

MNTINIAKNDFSDIELAAIPFNTLADHYGERLAREQLALEHESYEMGEARFRKMFERQLK
AGEVADNAAAKPLITTLPLPKMIARINDWFEEVKAKRGKRPTAFQFLQEIKPEAVAYITIK
TTLACLTADNTTVQAVASAI GRAIEDEARFGRIRDLEAKHFKNVEEQLNKRVGHVYKK
AFMQVVEADMLSKGLLGGEAWSSWHKEDSIHVGVRCIEMLIESTGMVSLHRQNAGVVGQD
SETIELAPEYAEAIATRAGALAGISPMFQPCVVPPKPWTGITGGGYWANGRRPLALVRTH
SKKALMRYEDVYMPEVYKAINIAQNTAWKINKKVLAVANVITKWKHCPVEDIPAIEREEL
PMKPEDIDMNPEALTAWKRAAAAVYRKDKARKSRRISLEFMLEQANKFANHKAIWFPYNM
DWRGRVYAVSMFNPQGNDMTKGLLTAKGKPIGKEGYWLKIHGANCAGVDKVPFPERIK
FIEENHENIMACAKSPLENTWAEQDSPFCFLAFCFEYAGVQHHGLSYNCSLPLAFDGSC
SGIQHFSAMLRDEVGGRAVNLLPSETVQDIYGIVAKKVNEILQADAINGTDNEVVTVTDE
NTGEISEKVKLGTKALAGQWLAYGVTRSVTKRSVMTLAYGSKEFGFRQQVLEDTIQPAID
SGKGLMFTQPNQAAGYMAKLIWESVSVTVVAAVEAMNWLKSAAKLLAAEVKDKKTGEILR
KRCVHWVTPDGFVPVQYKKPIQTRLNLMFLGQFRLQPTINTNKDSEIDAHKQESGIAP
NFVHSQDGSHLRKTVVWAHEKYGIESFALIHDSFGTIPADAANLFKAVRETMVDITYESCD
VLADFYDQFADQLHESQLDKMPALPAKGNLNLRDILESDFAF



Negative CIS Elements	Negative repeat Elements
1	0

Indices based on non-uniform usage of synonymous codon				
Index	Description	Range	Value	Reference
RSCU	Relative Synonymous Codon Usage	0~6	1.15	[11]
ENC	Effective Number of Codons	20~61	50.3	[2, 3]
RCBS	Relative Codon Bias Strength	≥0	0.16	[41]
DCBS	Directional Codon Bias Score	≥1	1.53	[51]
CDC	Codon Deviation Coefficient	0~1	0.1	[61]
MILC	Measure Independent of Length and Composition	-1~1	-0.44	[71]

ICDI	Intrinsic Codon Deviation Index	0~1	0.11	[8]
SCUO	Synonymous Codon Usage Order	0~1	0.08	[9, 10]
Ew	Weighted Sum of Relative Entropy	0~1	0.87	[11]
P	Codon Preference	≥ 1	1.06	[12]
MCB	Maximum-likelihood Codon Bias	≥ 0	0.36	[13]

Indices based on codon frequency in a reference set of genes

Index	Description	Range	Value	Reference
CAI	Codon Adaptation Index	0~1	0.74	[1]
CFD	Codon Frequency Distribution	0~1	0.04	[2]
FOP	Frequency of Optimal Codons	0~1	0.48	[3, 4]
COUSIN59	Codon Usage Similarity Index	∞	0.86	[5]
COUSIN18			1.04	
CBI	Codon Bias Index	-1~1	0.2	[6]
Dmean	Mean Dissimilarity-based Index	0~2	0.13	[7]
RCA	Relative Codon Adaptation	≥ 0	1.1	[8]
CUFS	Codon Usage Frequency Similarity	0~ $\sqrt{2}$	0.13	[9]
B	Codon Usage Bias	0~2	0.16	[10]

Indices based on adaptation to the tRNA levels and their supply

Index	Description	Range	Value	Reference
tAI	tRNA Adaptation Index	0~1	0.51	[1]
gtAI	Genetic tRNA Adaptation Index	0~1	0.59	[2]
P2	P2 Index	0~1	0.5	[3]

Indices based on complex patterns of codon usage

Index	Description	Range	Value	Reference
GC3	GC Content at the Third Position of Synonymous Codons	0~1	0.38	[1]
GC	GC Content		0.43	
GC1	GC Content at the First Position of Synonymous Codons		0.53	
GC2	GC Content at the Second Position of Synonymous Codons		0.39	
ENcp	Effective Number of Codon Pairs	20~61	35.63	[2]
CPS	Codon Pair Score	-1~1	0.03	[3, 4]
Codon Volatility	Codon Volatility	0.5~1	0.76	[5]

