

Supplementary Figure 1. Co-occurring MDR patterns of *bla*_{CMY-2}, *bla*_{CTX-M-55} and *bla*_{CTX-M-65} plasmid types

The nodes represent genes, and the connecting branches (edges) represent the frequency of co-occurrence at a given threshold. The threshold of each co-occurring network is represented by the highest standard deviation (SD) that demonstrates MDR. The number of connections represent the number of isolates demonstrating that co-occurring pattern. A) Most common co-occurring AMR determinants demonstrating MDR in association with *bla*_{CMY-2} plasmid types. B) Most common co-occurring AMR determinants demonstrating MDR in association with *bla*_{CTX-M-55} plasmid types. C) Most common co-occurring AMR determinants demonstrating MDR in association with *bla*_{CTX-M-65} plasmid types.

Supplementary Tables

Supplementary Table S1: Phenotypic and genotypic susceptibility profiles of all isolates in the study.

Supplementary Table S2: Summary table of final comparison between phenotypic and genotypic data with corresponding CLSI breakpoints for antimicrobials used in study.

Supplementary Table S3: Discordant isolates and their corresponding genotypic-phenotypic susceptibility results.

Supplementary Table S4: The prevalence rate of each resistance gene detected in concordant isolates.

Supplementary Table S5: Frequency of resistant and intermediate AMR determinant combinations according to antimicrobial class with its corresponding phenotypic susceptibilities.

Supplementary Table S6: Genomic profiles of multidrug resistant isolates harbouring antimicrobial resistant (AMR) determinants *bla*_{CMY-2}, *bla*_{CTX-M-55} and *bla*_{CTX-M-65}.