1978: Viola rupestris ssp. relicta

Relevant synonyms	—
Norwegian name	Kalkfiol
Feature group	Vascular plants
Expert(s)	Hanne Hegre
Date	2025-12-17

Alpine (ALP)

Occurrence status	Present regularly (PRE)	
Overall conclusion	🔴 **Unfavourable - bad (U2)**	↘️ **Deteriorating**
Range	🟢 Favourable (FV)	➡️ Stable
Population	🟡 Unfavourable - inadequate (U1)	↘️ Deteriorating
Habitat for species	🔴 Unfavourable - bad (U2)	↘️ Deteriorating
Future prospects	🔴 Unfavourable - bad (U2)	

Atlantic (ATL)

Occurrence status	Present regularly (PRE)	
Overall conclusion	🔴 **Unfavourable - bad (U2)**	↘️ **Deteriorating**
Range	🟢 Favourable (FV)	➡️ Stable
Population	🟡 Unfavourable - inadequate (U1)	↘️ Deteriorating
Habitat for species	🔴 Unfavourable - bad (U2)	↘️ Deteriorating
Future prospects	🔴 Unfavourable - bad (U2)	

Pressures

- PA05: Abandonment of management/use of grasslands and other agricultural and agroforestry systems (e.g. cessation of grazing, mowing or traditional farming) [ALP, ATL]
- PI03: Problematic native species [ALP, ATL]
- PJ01: Temperature changes and extremes due to climate change [ALP, ATL]
- PJ10: Change of habitat location, size, and / or quality due to climate change [ALP]

Data sources, references

- Solstad, H., Elven, R., Arnesen, G., Eidesen, P.B., Gaarder, G., Hegre, H., Høitomt, T., Mjelde, M. and Pedersen, O. 2021. Karplanter: Vurdering av kalkfiol Viola rupestris subsp. relicta for Norge. Rødlista for arter 2021. Artsdatabanken. https://lister.artsdatabanken.no/rodlisteforarter/2021/30355. Downloaded 12.12.2025
- Nordal, I. and Marcussen, T. 2005. Kalkfiol Viola rupestris ssp. relicta er bisentrisk i Skandinavia – ny forekomst på Møre. Blyttia 63:67–71. [Viola rupestris ssp. relicta is bicentric in Scandinavia – new finding in Møre, W. Norway.]
- Nordal, I. Jonsell, B. and Marcussen, T. 2005. Viola rupestris: molecular analyses to elucidate postglacial migration in Western Europe. Journal of Biogeography 32: 1453–1459.
- Marcussen, T., Karlsson, T., Wind, P. and Jonsell, B. 2010. Violaceae. In: Jonsell, B. and Karlsson, T (ed.), Flora Nordica. 6. Thymelaeaceae to Apiaceae: 12–52. The Swedish Museum of Natural History, Stockholm.



The map is based on officially registered occurrences (https://artskart.artsdatabanken.no/) and expert inputs from the ongoing red list assessment. Some previous reports of this subspecies probably belong to subsp. rupestris.