Outputs from the **ACEfit** function of APACE software

Filename	Description	Comments/Caveats
Basic summaries		
Stdev.nii	Phenotypic standard deviation	
Heritability		
h2_dist.pdf	Cumulative distribution of all in-mask h ²	
h2_hist.pdf	Histogram of positive in-mask h ²	Percentage of positive h ² is provided
ACE_A_h2.nii	Narrow sense heritability, h ²	
ACE_A_LRT.nii	Likelihood ratio test (LRT) statistic for testing H ₀ : h ² =0	Basis for all thresholding procedures (computed and saved if image-wise inference is made)
ACE_A_LRT_vox_Pasym.nii	-log ₁₀ P-value for testing H ₀ : h ² =0 using large sample, parametric null distribution	Not trustworthy except for very large samples (computed and saved if image-wise inference is made)
Common Environment		
ACE_C_c2.nii	Estimate of c ² , shared environmental component	
ACE_C_LRT.nii	LRT statistic for testing H ₀ : c ² =0	(computed and saved if image- wise inference is made)
ACE_C_LRT_vox_Pasym.nii	-log ₁₀ P-value for testing H ₀ : c ² =0 using large sample, parametric null distribution	Not trustworthy except for very large samples; note that no permutation test is possible for c ² . (computed and saved if image-wise inference is made)
Error		
ACE_E_e2.nii	Estimate of e ² , unique error component	

Additional outputs from permutation inference of APACE software

Filename	Description	Comments/Caveats
Heritability		
H0dist_mean.pdf	Null distribution of mean of all in-mask h ²	Permutation-based p-value is
		displayed in plot title
H0dist_wh2.pdf	Null distribution of variance-weighted average	Permutation-based p-value is
	of all in-mask h ²	displayed in plot title
H0dist_median.pdf	Null distribution of median (Q2) of all in-mask h ²	Permutation-based p-value is
		displayed in plot title
H0dist_q3.pdf	Null distribution of the third quartile (Q3) of all	Permutation-based p-value is
	in-mask h ²	displayed in plot title
H0dist_mGTmedian.pdf	Null distribution of in-mask h ² >Q2(h ²)	Permutation-based p-value is
		displayed in plot title
H0dist_mGTq3.pdf	Null distribution of in-mask h ² >Q3(h ²)	Permutation-based p-value is
		displayed in plot title
H0dist_test_statistic.pdf	Null distribution of maximum LRT statistic	Permutation-based p-value is
		displayed in plot title. Only produced if
		image-wise inference is made.
h2_FDRplot.pdf	Scatter plot of voxel/element-wise FDR-	Only produced if image-wise inference
	adjusted p-values vs. expected p-values for	is made.
	heritability	
ACE_A_LRT_vox_P.nii	$-\log_{10}$ P-value for testing H ₀ : h^2 =0 using	Only produced if image-wise inference
	nonparametric permutation-based null	is made.
	distribution	
ACE_A_LRT_vox_FWEP.nii	Voxel/element-wise FWE-corrected -log ₁₀ P-	Only produced if image-wise inference
	value for testing H_0 : $h^2=0$ using nonparametric	is made.
	permutation-based null distribution	
ACE_A_LRT_vox_FDRP.nii	Voxel/element-wise FDR-adjusted -log ₁₀ P-	Only produced if image-wise inference
	value for testing H_0 : $h^2=0$ using nonparametric	is made.
	permutation-based null distribution	
H0dist_cluster_size.pdf	Null distribution of maximum suprathreshold	Permutation-based p-value is
	cluster size	displayed in plot title. Only produced if

		image-wise cluster inference is made.
ACE_A_LRT_clus.nii	Suprathreshold cluster sizes on LRT statistic for	Cluster forming threshold of u=2.706,
	testing H ₀ : h ² =0	the large permutation-based sample
		α=0.05 threshold. Only produced if
		image-wise cluster inference is made.
ACE_A_LRT_clus_FWEP.nii	Cluster FWE-corrected -log ₁₀ P-value (size	Only produced if image-wise cluster
	statistic)	inference is made
H0dist_cluster_mass.pdf	Null distribution of maximum suprathreshold	Permutation-based p-value is
	cluster mass	displayed in plot title. Only produced if
		image-wise cluster inference is made.
ACE_A_LRT_mass.nii	Suprathreshold cluster mass on LRT statistic	Cluster forming threshold of u=2.706,
	for testing H ₀ : h ² =0	the large permutation-based sample
		α=0.05 threshold. Only produced if
		image-wise cluster inference is made.
ACE_A_LRT_mass_FWEP.nii	Cluster FWE-corrected -log ₁₀ P-value (mass	Only produced if image-wise cluster
	statistic)	inference is made.
Pairwise Correlation		
H0dist_Tstat_rMZ_rDZ.pdf	Null distribution of 2-sample t-statistic for	Permutation-based p-value is
	testing H ₀ : r _{MZ} =r _{DZ}	displayed in plot title.
HOdist_Diff_rMZ_rDZ.pdf	Null distribution of mean difference of r _{MZ} and	Permutation-based p-value is
	r_{DZ}	displayed in plot title.
HOdist_Tstat_rDZ_rSib.pdf	Null distribution of 2-sample t-statistic for	Permutation-based p-value is
	testing H ₀ : r _{DZ} =r _{Sib}	displayed in plot title.
HOdist_Diff_rDZ_rSib.pdf	Null distribution of mean difference of r _{DZ} and	Permutation-based p-value is
	r _{Sib}	displayed in plot title.