

Chapter 4

General Vector Spaces

4.1. Real Vector Spaces

4.2. Subspaces

4.3. Spanning Sets

4.4. Linear Independence

4.5. Coordinates and Basis

4.6. Dimension

4.8. Row Space, Column Space, and Null Space

4.9. Rank, Nullity, and the Fundamental Matrix Spaces

Chapter 4.3

Spanning Sets

Linear Combination

DEFINITION 1

If \mathbf{w} is a vector in a vector space V , then \mathbf{w} is said to be a *linear combination* of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r$ in V if \mathbf{w} can be expressed in the form

$$\mathbf{w} = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \cdots + k_r\mathbf{v}_r$$

where k_1, k_2, \dots, k_r are scalars.

These scalars are called the *coefficients* of the linear combination.

Linear Combination

THEOREM 4.3.1

If $S = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r\}$ is a nonempty set of vectors in a vector space V , then:

The set W of all possible linear combinations of the vectors in S is a subspace of V .



Proof

exe

Let $V = \mathbb{R}^2$

$$S_1 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$$

$$S_2 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$



THEOREM 4.3.1

Proof Let W be the set of all possible linear combinations of the vectors in S . We must show that W is closed under addition and scalar multiplication. To prove closure under addition, let

$$\mathbf{u} = c_1\mathbf{w}_1 + c_2\mathbf{w}_2 + \cdots + c_r\mathbf{w}_r \quad \text{and} \quad \mathbf{v} = k_1\mathbf{w}_1 + k_2\mathbf{w}_2 + \cdots + k_r\mathbf{w}_r$$

be two vectors in W . It follows that their sum can be written as

$$\mathbf{u} + \mathbf{v} = (c_1 + k_1)\mathbf{w}_1 + (c_2 + k_2)\mathbf{w}_2 + \cdots + (c_r + k_r)\mathbf{w}_r$$

which is a linear combination of the vectors in S . Thus, W is closed under addition. We leave it for you to prove that W is also closed under scalar multiplication and hence is a subspace of V .

Span



The subspace W in Theorem 4.3.1 is called the subspace of V *spanned* by S . The vectors $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r$ in S are said to *span* W , and we write

$$W = \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r\} \quad \text{or} \quad W = \text{span}(S)$$

Let $V = \mathbb{R}^2$

$$S_1 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$$

$$S_2 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$



$$W_1 = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$$

$$W_2 = \text{span}(S_2)$$

The Standard Unit Vectors Span R^n

EXAMPLE 1

Recall that the standard unit vectors in R^n are

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0), \quad \mathbf{e}_2 = (0, 1, 0, \dots, 0), \dots, \quad \mathbf{e}_n = (0, 0, 0, \dots, 1)$$


These vectors span R^n since every vector $\mathbf{v} = (v_1, v_2, \dots, v_n)$ in R^n can be expressed as

$$\mathbf{v} = v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + \dots + v_n \mathbf{e}_n$$


which is a linear combination of $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

A Geometric View of Spanning in R^2 and R^3

EXAMPLE 2



(a) $\text{span}\{\mathbf{v}\}$ is the line through the origin determined by \mathbf{v}



(b) $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$ is the plane through the origin determined by \mathbf{v}_1 and \mathbf{v}_2

A Spanning Set for P_n

EXAMPLE 3

The polynomials $1, x, x^2, \dots, x^n$ span the vector space P_n since each polynomial \mathbf{p} in P_n can be written as

$$\mathbf{p} = a_0 + a_1x + \cdots + a_nx^n$$

which is a linear combination of $1, x, x^2, \dots, x^n$.

We can denote this by writing $P_n = \text{span}\{1, x, x^2, \dots, x^n\}$


Two Important Types of Problems

- Given a nonempty set S of vectors in R^n and a vector \mathbf{v} in R^n , determine whether \mathbf{v} is a linear combination of the vectors in S .
- Given a nonempty set S of vectors in R^n , determine whether the vectors span R^n .

Let $V = \mathbb{R}^2$

$$S_1 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$$


$$S_2 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$



If $\mathbf{v} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \in \mathbb{R}^2$

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} \stackrel{?}{=} a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 2 \\ 2 \end{bmatrix} + c \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

for some real numbers a, b, c



If $\text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \stackrel{?}{=} \mathbb{R}^2$

Linear Combination Problem

EXAMPLE 4

Consider the vectors $\mathbf{u} = (1, 2, -1)$ and $\mathbf{v} = (6, 4, 2)$ in R^3 .

Show that $\mathbf{w} = (9, 2, 7)$ is a linear combination of \mathbf{u} and \mathbf{v} and $\mathbf{w}' = (4, -1, 8)$ is *not* a linear combination of \mathbf{u} and \mathbf{v} .

Solution

$$\mathbf{w} = k_1 \mathbf{u} + k_2 \mathbf{v}$$

$$(9, 2, 7) = k_1(1, 2, -1) + k_2(6, 4, 2) = (k_1 + 6k_2, 2k_1 + 4k_2, -k_1 + 2k_2)$$

EXAMPLE 4 Cont.

Equating corresponding components gives

$$\begin{aligned}k_1 + 6k_2 &= 9 \\2k_1 + 4k_2 &= 2 \\-k_1 + 2k_2 &= 7\end{aligned}$$

Solving this system

$$k_1 = -3, k_2 = 2, \text{ so } \mathbf{w} = -3\mathbf{u} + 2\mathbf{v}$$

Similarly,

$$\mathbf{w}' = k_1\mathbf{u} + k_2\mathbf{v}$$

$$(4, -1, 8) = k_1(1, 2, -1) + k_2(6, 4, 2) = (k_1 + 6k_2, 2k_1 + 4k_2, -k_1 + 2k_2)$$

$$\begin{aligned}k_1 + 6k_2 &= 4 \\2k_1 + 4k_2 &= -1 \\-k_1 + 2k_2 &= 8\end{aligned}$$

This system of equations is inconsistent
 \mathbf{w}' is not a linear combination of \mathbf{u} and \mathbf{v} .

Testing for Spanning

EXAMPLE 5

Determine whether the vectors $\mathbf{v}_1 = (1, 1, 2)$, $\mathbf{v}_2 = (1, 0, 1)$, and $\mathbf{v}_3 = (2, 1, 3)$ span the vector space R^3 .

Solution

determine whether an arbitrary vector $\mathbf{b} = (b_1, b_2, b_3)$ in R^3 can be expressed as a linear combination

$$\mathbf{b} = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + k_3\mathbf{v}_3$$

$$(b_1, b_2, b_3) = k_1(1, 1, 2) + k_2(1, 0, 1) + k_3(2, 1, 3)$$

$$(b_1, b_2, b_3) = (k_1 + k_2 + 2k_3, k_1 + k_3, 2k_1 + k_2 + 3k_3)$$

EXAMPLE 5 Cont.

$$k_1 + k_2 + 2k_3 = b_1$$

$$k_1 + k_3 = b_2$$

$$2k_1 + k_2 + 3k_3 = b_3$$

Theorem 2.3.8, the system is consistent if and only if its coefficient matrix

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 0 & 1 \\ 2 & 1 & 3 \end{bmatrix} \text{ has a nonzero determinant.}$$

But this is *not* the case here; $\det(A) = 0$ so \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 do not span R^3 .

The Procedure for Identifying Spanning Sets

Step 1. Let $S = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r\}$ be a given set of vectors in V , and let \mathbf{x} be an arbitrary vector in V .

Step 2. Set up the augmented matrix for the linear system that results by equating corresponding components on the two sides of the vector equation $k_1\mathbf{w}_1 + k_2\mathbf{w}_2 + \dots + k_r\mathbf{w}_r = \mathbf{x}$

Step 3. Use the techniques developed in Chapters 1 and 2 to investigate the consistency or inconsistency of that system.

If it is consistent for *all* choices of \mathbf{x} , the vectors in S span V , and if it is inconsistent for *some* vector \mathbf{x} , they do not.

Testing for Spanning

EXAMPLE 6

Determine whether the set S spans P_2 .

$$(a) S = \{1 + x + x^2, -1 - x, 2 + 2x + x^2\}$$

$$(b) S = \{x + x^2, x - x^2, 1 + x, 1 - x\}$$

Solution (a) An arbitrary vector in P_2 is of the form $\mathbf{p} = a + bx + cx^2$,

$$k_1(1 + x + x^2) + k_2(-1 - x) + k_3(2 + 2x + x^2) = a + bx + cx^2$$

$$(k_1 - k_2 + 2k_3) + (k_1 - k_2 + 2k_3)x + (k_1 + k_3)x^2 = a + bx + cx^2$$

EXAMPLE 6 Cont.

$$\begin{bmatrix} 1 & -1 & 2 & a \\ 1 & -1 & 2 & b \\ 1 & 0 & 1 & c \end{bmatrix}$$

and whose coefficient matrix is

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & -1 & 2 \\ 1 & 0 & 1 \end{bmatrix} \quad \begin{matrix} \leftarrow \\ \leftarrow \end{matrix}$$

S does **not** span P_2 .

Solution (b) Using the same procedure the augmented matrix

$$\begin{bmatrix} 0 & 0 & 1 & -1 & a \\ 1 & 1 & 1 & -1 & b \\ 1 & -1 & 0 & 0 & c \end{bmatrix}$$

EXAMPLE 6 Cont.

The reduced row echelon form

$$\begin{bmatrix} 1 & 0 & 0 & 0 & \frac{-a+b+c}{2} \\ 0 & 1 & 0 & 0 & \frac{-a+b-c}{2} \\ 0 & 0 & 1 & -1 & a \end{bmatrix}$$

is consistent for every choice a , b , and c . Thus,

$$\text{span}(S) = P_2$$

Testing for Spanning

EXAMPLE 7

In each part, determine whether the set S spans M_{22} .

$$(a) S = \left\{ \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \right\}$$

$$(b) S = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix} \right\}$$

exe

EXAMPLE 7

Solution (a) An arbitrary vector in M_{22} is of the form $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, so Equation (2) becomes

$$k_1 \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + k_2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + k_3 \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} + k_4 \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

which we can rewrite as

$$\begin{bmatrix} k_1 + k_2 + k_3 + k_4 & 2k_1 + 2k_3 + k_4 \\ k_3 + k_4 & k_1 + k_2 + k_4 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Equating corresponding entries produces a linear system whose augmented matrix is

$$\left[\begin{array}{cccc|c} 1 & 1 & 1 & 1 & a \\ 2 & 0 & 2 & 1 & b \\ 0 & 0 & 1 & 1 & c \\ 1 & 1 & 0 & 1 & d \end{array} \right] \text{ and whose coefficient matrix is } A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 0 & 2 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$$

As in part (a) of Example 6, the coefficient matrix is square, so we can apply parts (e) and (g) of Theorem 2.3.8. We leave it for you to verify that $\det(A) = -2 \neq 0$, so the system is consistent for *every* choice of a, b, c , and d , which implies that $\text{span}(S) = M_{22}$.

EXAMPLE 7

Solution (b) Using the same procedure as in part (a), the augmented matrix for the linear system corresponding to Equation (2) is

$$\begin{bmatrix} 1 & -1 & 0 & 0 & a \\ 0 & 0 & 0 & 1 & b \\ 0 & 1 & 1 & -1 & c \\ 0 & 0 & 0 & 1 & d \end{bmatrix} \text{ and the coefficient matrix is } A = \begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ which}$$

is square, so once again we can apply parts (e) and (g) of Theorem 2.3.8. Since the second and fourth rows of this matrix are identical, it follows that $\det(A) = 0$. Thus, the system is inconsistent for *some* choice of a , b , c , and d , which implies that S does not span M_{22} .

Spanning Sets are not Unique

THEOREM 4.3.2

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ and $S' = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ are nonempty sets of vectors in a vector space V , then

$$\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\} = \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$$

if and only if each vector in S is a linear combination of those in S' , and each vector in S' is a linear combination of those in S .

This Theorem states that two different sets of vectors may span the same space.

Chapter 4-3 Objectives

- Given a set S of vectors in R^n and a vector \mathbf{v} in R^n , determine whether \mathbf{v} is a linear combination of the vectors in S .
- Given a set S of vectors in R^n , determine whether the vectors in S span R^n .
- Determine whether two nonempty sets of vectors in a vector space V span the same subspace of V .