

Chapter 4

General Vector Spaces

4.1. Real Vector Spaces

4.2. Subspaces

4.3. Spanning Sets

4.4. Linear Independence

4.5. Coordinates and Basis

4.6. Dimension

4.8. Row Space, Column Space, and Null Space

4.9. Rank, Nullity, and the Fundamental Matrix Spaces


Chapter 4.2

Subspaces

Subspaces

DEFINITION 1

A subset W of a vector space V is called a *subspace* of V if W is itself a vector space under the addition and scalar multiplication defined on V .



E.g. The vectors \mathbf{u} and \mathbf{v} are in W , but the vectors $\mathbf{u} + \mathbf{v}$ and $k\mathbf{u}$ are not.

Show that a Nonempty Set is a Subspace

Those axioms that are **not** inherited by W are

Axiom 1 — Closure of W under addition

Axiom 4 — Existence of a zero vector in W

Axiom 5 — Existence of a negative in W for every vector in W

Axiom 6 — Closure of W under scalar multiplication

so these must be verified to prove that it is a subspace of V .

However, the following theorem shows that if Axiom 1 and Axiom 6 hold in W , then Axioms 4 and 5 hold in W as a consequence and hence **need not** be verified.

Subspace Test

THEOREM 4.2.1

If W is a nonempty set of vectors in a vector space V , then W is a subspace of V if and only if the following conditions are satisfied.

- (a) If \mathbf{u} and \mathbf{v} are vectors in W , then $\mathbf{u} + \mathbf{v}$ is in W .
- (b) If k is a scalar and \mathbf{u} is a vector in W , then $k\mathbf{u}$ is in W .

Proof

exe

THEOREM 4.2.1

Proof

If W is a subspace of V , then all the vector space axioms hold in W , including Axioms 1 and 6, which are precisely conditions (a) and (b).

Conversely, assume that conditions (a) and (b) hold. Since these are Axioms 1 and 6, and since Axioms 2, 3, 7, 8, 9, and 10 are inherited from V , we only need to show that Axioms 4 and 5 hold in W .

For this purpose, let \mathbf{u} be any vector in W .

It follows from condition (b) that the product $k\mathbf{u}$ is also a vector in W for every scalar k . In particular, $0\mathbf{u} = \mathbf{0}$ and $(-1)\mathbf{u} = -\mathbf{u}$ are in W , which shows that Axioms 4 and 5 hold in W .

The Zero Subspace

EXAMPLE 1

If V is any vector space, and if $W = \{\mathbf{0}\}$ is the subset of V that consists of the zero vector only, then W is closed under addition and scalar multiplication since


$$\mathbf{0} + \mathbf{0} = \mathbf{0} \quad \text{and} \quad k\mathbf{0} = \mathbf{0} \quad \text{for any scalar } k.$$

We call W the *zero subspace* of V .


Subspaces of R^2 and of R^3

EXAMPLE 2

Lines through the origin are subspaces of R^2 and of R^3



(a) W is closed under addition.




(b) W is closed under scalar multiplication.

Subspaces of R^2 and of R^3

EXAMPLE 3

Planes through the origin are subspaces of R^3



The vectors $\mathbf{u} + \mathbf{v}$ and $k\mathbf{u}$ both lie in the same plane as \mathbf{u} and \mathbf{v} .

Subspaces of R^2 and of R^3

Subspaces of R^2

- $\{\mathbf{0}\}$
- Lines through the origin
- R^2

Subspaces of R^3

- $\{\mathbf{0}\}$
 - Lines through the origin
 - Planes through the origin
 - R^3
-


A Subset of \mathbb{R}^2 That Is Not a Subspace

EXAMPLE 4

Let W be the set of all points (x, y) in \mathbb{R}^2 for which $x \geq 0$ and $y \geq 0$.

This set is not a subspace of \mathbb{R}^2 because it is not closed under scalar multiplication.

$\mathbf{v} = (1, 1)$ is a vector in W ,
but $(-1)\mathbf{v} = (-1, -1)$ is not.



Subspaces of M_{nn}

EXAMPLE 5

the set of symmetric $n \times n$ matrices is closed under addition and scalar multiplication and hence is a subspace of M_{nn} .

Similarly, the sets of upper triangular matrices, lower triangular matrices, and diagonal matrices are subspaces of M_{nn} .

A Subset of M_{nn} That Is Not a Subspace

EXAMPLE 6

The set W of invertible $n \times n$ matrices is not a subspace of M_{nn} :

$$U = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix} \quad \text{and} \quad V = \begin{bmatrix} -1 & 2 \\ -2 & 5 \end{bmatrix}$$

The matrix $0U$ is the 2×2 zero matrix and hence is not invertible, and the matrix $U + V$ has a column of zeros so it also is not invertible.

The Subspace of All Polynomials

EXAMPLE 7

Recall that a *polynomial* is a function that can be expressed in the form

$$p(x) = a_0 + a_1x + \cdots + a_nx^n$$

where a_0, a_1, \dots, a_n are constants. Denote this space by P_n .

It is evident that the sum of two polynomials is a polynomial and that a constant times a polynomial is a polynomial.

Thus, the set W of all polynomials is a subspace of $F(-\infty, \infty)$.

(i.e. denote this space by P_∞ .) (i.e. a function space)

Subspaces Test

EXAMPLE 8

Determine whether the indicated set of matrices is a subspace of M_{22} .

(a) The set U consisting of all matrices of the form

$$\begin{bmatrix} x & 0 \\ 2x & y \end{bmatrix}$$

(b) The set W consisting of all 2×2 matrices A such that

$$A \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

EXAMPLE 8 cont.

(a) If A and B are matrices in U , then they can be expressed in the form

$$A = \begin{bmatrix} a & 0 \\ 2a & b \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} c & 0 \\ 2c & d \end{bmatrix}$$

for some real numbers a, b, c , and d .

$$A + B = \begin{bmatrix} a + c & 0 \\ 2(a + c) & b + d \end{bmatrix} \quad \text{is a matrix in } U$$

Thus, U is closed under addition.

$$kA = \begin{bmatrix} ka & 0 \\ 2ka & kb \end{bmatrix} \quad \text{is also a matrix in } U$$

Moreover, U is closed under scalar multiplication.

Hence, U is a subspace of M_{22} .

EXAMPLE 8 cont.

(b) Let $A = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$

This is a vector in W since

$$A \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

However, $2A$ does not satisfy Equation (3) since

$$(2A) \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ -2 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

This alone establishes that W is not a subspace of M_{22} .

Subspaces Test

EXAMPLE 9

Determine whether the indicated set of polynomials is a subspace of P_2 .

(a) The set U consisting of all polynomials of the form

$$\mathbf{p} = 1 + ax - ax^2, \text{ where } a \text{ is a real number.}$$

(b) The set W consisting of all polynomials \mathbf{p} in P_2

such that $\mathbf{p}(2) = 0$.

EXAMPLE 9 cont.

(a)

The set U is not a subspace of P_2 because it is not closed under addition.

polynomials $\mathbf{p} = 1 + x - x^2$ and $\mathbf{q} = 1 + 2x - 2x^2$ are in U , but

$\mathbf{p} + \mathbf{q} = 2 + 3x - 3x^2$ is not.

(b)

If \mathbf{p} and \mathbf{q} are polynomials in W , and k is any real number, then

$$(\mathbf{p} + \mathbf{q})(2) = \mathbf{p}(2) + \mathbf{q}(2) = 0 + 0 = 0$$

$$(k\mathbf{p})(2) = k \cdot \mathbf{p}(2) = k \cdot 0 = 0.$$

Since $\mathbf{p} + \mathbf{q}$ and $k\mathbf{p}$ are in W , it follows that W is a subspace of P_2 .

Building Subspaces

THEOREM 4.2.2

If W_1, W_2, \dots, W_r are subspaces of a vector space V , then the intersection of these subspaces is also a subspace of V .

Proof

exe

THEOREM 4.2.2

Proof Let W be the intersection of the subspaces W_1, W_2, \dots, W_r . This set is not empty because each of these subspaces contains the zero vector of V , and hence so does their intersection. Thus, it remains to show that W is closed under addition and scalar multiplication.

To prove closure under addition, let \mathbf{u} and \mathbf{v} be vectors in W . Since W is the intersection of W_1, W_2, \dots, W_r , it follows that \mathbf{u} and \mathbf{v} also lie in each of these subspaces. Moreover, since these subspaces are closed under addition and scalar multiplication, they also all contain the vectors $\mathbf{u} + \mathbf{v}$ and $k\mathbf{u}$ for every scalar k , and hence so does their intersection W . This proves that W is closed under addition and scalar multiplication. ■

Solution Spaces of Homogeneous Systems

THEOREM 4.2.3

The solution set of a homogeneous system $A\mathbf{x} = \mathbf{0}$ of m equations in n unknowns is a subspace of R^n .

Proof

[Note]

Because the solution set of a homogeneous system in n unknowns is actually a subspace of R^n , we will generally refer to it as the *solution space* of the system.

Solution Spaces of Homogeneous Systems

EXAMPLE 10 Give a geometric description of the solution set.

$$(a) \begin{bmatrix} 1 & -2 & 3 \\ 2 & -4 & 6 \\ 3 & -6 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$(b) \begin{bmatrix} 1 & -2 & 3 \\ -3 & 7 & -8 \\ -2 & 4 & -6 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$(c) \begin{bmatrix} 1 & -2 & 3 \\ -3 & 7 & -8 \\ 4 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$(d) \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

EXAMPLE 10 cont.

(a) The solutions are $x = 2s - 3t$, $y = s$, $z = t$

it follows that $x = 2y - 3z$ or $x - 2y + 3z = 0$

This is a plane through the origin that has $\mathbf{n} = (1, -2, 3)$ as a normal.

(b) The solutions are $x = -5t$, $y = -t$, $z = t$

A parametric equations for the line through the origin that is parallel to the vector $\mathbf{v} = (-5, -1, 1)$.

(c) The only solution is $x = 0, y = 0, z = 0$,
so the solution space consists of the single point $\{\mathbf{0}\}$.

(d) This linear system is satisfied by all real values of x, y , and z ,
so the solution space is all of R^3 .

The Linear Transformation Viewpoint

The solution space of $A\mathbf{x} = \mathbf{0}$ is the set of vectors in R^n , that T_A maps into the zero vector in R^m . This set is sometimes called the *kernel* of the transformation.

THEOREM 4.2.4.

If A is an $m \times n$ matrix, then the kernel of the matrix transformation $T_A : R^n \rightarrow R^m$ is a subspace of R^n .

Chapter 4-2 Objectives

- Determine whether a subset of a vector space is a subspace.
- Show that a subset of a vector space is a subspace.
- Show that a nonempty subset of a vector space is not a subspace by demonstrating that the set is either not closed under addition or not closed under scalar multiplication.