Surgery Name Chronic Kidney Disease Review

Review Status: Review Required (CKD Stage 3-5 with >6 months since last eGFR, ACR >30, or high-risk)

Current EMIS Status: No EMIS CKD entry

Results Overview

Patient Information

NHS Number: 169663Age: 94 | Gender: Female

CKD Overview

• Stage: Stage 3b | ACR criteria: A1

• Albumin-Creatinine Ratio (ACR): 0.01 mg/mmol

• Creatinine:

Current: 131 μmol/L | Date: 2023-07-10
 3 Months Prior: 0 μmol/L | Date: nan

eGFR:

Current: 32 ml/min/1.73m² | Date: 2023-07-10
 3 Months Prior: 0 ml/min/1.73m² | Date: nan

• Anaemia Overview:

• Haemoglobin: 136 g/L | Date: nan

• Current Status: Normal

• Anaemia Management: No Action Needed

Electrolyte and Mineral Bone Disorder (MBD) Management

• Potassium: 4.7 mmol/L | Status: Normal | Date: nan

• Phosphate: 1.57 mmol/L | Status: Hyperphosphatemia | Date: nan

• Calcium: 2.35 mmol/L | Status: Normal | Date: nan

MBD Status: Check CKD-MBD
 Vitamin D Level: 37 ng/mL | Date: nan

• Vitamin D Status: Normal

Blood Pressure

Classification: Normal | Date: nan
 Systolic / Diastolic: 128 / 84 mmHg

• Target BP: <130/80 | BP Status: Above Target

Kidney Failure Risk

2-Year Risk: 0.03%5-Year Risk: 0.11%

The patient's 2- and 5-year kidney failure risk scores estimate the likelihood that their kidney disease will progress to kidney failure within the next 2 or 5 years. These scores are calculated based on the patient's current kidney function and other risk factors such as age, blood pressure, and existing health conditions. Understanding these risk scores helps in predicting disease progression and planning appropriate treatment strategies.

Care & Referrals

Multidisciplinary Care: Not Indicated
 Modality Education: Not Indicated
 Nephrology Referral: Not Indicated

• Persistent Proteinuria: No Referral Needed

Medication Review

• Current Medication: Ramipril 5mg

• Review Medications: No adjustments needed

• Contraindicated Medications: No contraindications

• **Suggested Medications:** Statin (e.g., Atorvastatin, Rosuvastatin, Simvastatin), ACE inhibitors (if not contraindicated) for proteinuria, Phosphate binders (e.g., Sevelamer, Lanthanum carbonate, Calcium acetate)

• Statin Recommendation: Consider Statin

Diabetes and HbA1c Management

HbA1c Level: 47 mmol/mol | Date: nan
 HbA1c Management: On Target

Lifestyle and Preventative Advice

Lifestyle Recommendations: Encourage a balanced, low-sodium diet. Advise moderate, regular physical activity
while monitoring for fatigue. Reinforce the importance of smoking cessation, weight management, and avoiding overthe-counter NSAIDs.

NICE Guideline Recommendations

CKD Stage G3b Recommendations:

- Monitoring and Risk Management: Continue primary care management with renal function tests every 6 months, or
 more frequently if ACR is greater than 3 mg/mmol. Use the KFRE to assess progression risk; refer to nephrology if the 5year risk exceeds 5% or if there's a rapid decline in eGFR.
- Referral Considerations: Consider nephrology referral for further evaluation and management, especially if complications like anaemia, electrolyte imbalances, or bone mineral disorders arise.
- **Lifestyle and Preventive Measures:** Aggressively manage BP and cardiovascular risk factors. Optimize dosing of ACE inhibitors or ARBs. Continue statin therapy as indicated.
- Patient Education: Reinforce the importance of lifestyle modifications and adherence to treatment plans to slow CKD progression.

Surgery Name, Add 1, Add 2, Belfast, BT123456

Telephone: 028 90*********