

INDICATION

For use in patients with GI bleeding who have already undergone an endoscopy. The pre-endoscopy Rockall score may be used prior to endoscopy.

Determines risk of re-bleeding and mortality

ADDITIONAL INFORMATION

COMORBIDITIES LIST	
Pneumonia	Cardiac failure
Dementia	Ischaemic heart disease
Recent major operation	Asthma
Malignancy	COPD
CVA/TIA	Diabetes mellitus
Haematological malignancy	Rheumatoid arthritis
Hypertension	Liver failure
Trauma/burns	Renal failure
Other cardiac disease	Disseminated malignancy
Major sepsis	Other liver disease

INTERPRETATION

SCORE	REBLEED	MORTALITY
0	4.9%	0%
1	3.4%	0%
2	5.3%	0.2%
3	11.2%	2.9%
4	14.1%	5.3%
5	24.1%	10.8%
6	32.9%	17.3%
7	43.8%	27%
≥8	41.8%	41.1%

CALCULATION

Total score = sum of scores for each question

QUESTION		POINTS
Age	<60 years	+0
	60-79 years	+1
	≥80 years	+2
Shock	None	+0
	HR ≥100	+1
	SBP <100	+2
Comorbidities	No major comorbidity	+0
	Any comorbidity EXCEPT Renal failure / Liver failure / Disseminated malignancy	+2
	Renal failure / Liver failure / Disseminated malignancy	+3
Diagnosis	Mallory-Weiss tear	+0
	No lesion identified / no hemorrhage	+0
	All other diagnoses	+1
	Malignancy of UGI tract	+2
Major stigmata of recent haemorrhage	None	+0
	Dark spot only	+0
	Blood in upper GI tract	+2
	Adherent clot	+2
	Visible or spurting vessek	+2