

INDICATION

Determines severity in Crohn's disease

INTERPRETATION

SCORE	RISK CATEGORY
<150	Possible remission
150-220	Mild
220-450	Moderate to Severe
>450	Severe

CALCULATION

Total score = sum of scores for each question

QUESTION	POINTS	MULTIPLIER
% below ideal weight (Weight / Ideal body weight / Sex)	$(\text{Ideal} - \text{observed}) / \text{ideal} \times 100$	$\times 1$
Haematocrit:	Expected – observed hematocrit (Men 47% Women 42%)	$\times 6$
Number of soft / liquid stools	Total number past 7 days	$\times 2$
Abdominal pain: average daily rating 7 days	0 = None 1 = Mild 2 = Moderate 3 = Severe	$\times 35$
General well-being: average daily rating 7 days	0 = Generally well 1 = Slightly under par 2 = Poor 3 = Very poor 4 = Terrible	$\times 49$
Anti-diarrhoea medication use in last 7 days	1 = Yes 0 = No	$\times 30$
Abdominal mass	0 = None 2 = Questionable 5 = Definite	$\times 10$
Extra-intestinal / physical complaints: Arthritis / Arthralgia / Iritis / Uveitis / Erythema nodosum / Pyoderma gangrenosum / Aphthous stomatitis / Anal fissure / Fistula / Abscess / Other fistula / Temp >37.8°C or 100°F	1 = Yes 0 = No	$\times 20$