MAL DIAGNOSIS:

PART 1: BREAST, RICHT, SEGMENTAL MASTECTOMY -

- INFILTRATING DUCTAL CARCINOMA, NOS TYPE, MODERATELY DIFFERENTIATED.
- THE TUMOR MEASURES 2.3 CM (GROSS MEASUREMENT), NOTTINGHAM SCORE 6/9 (TUBULES 2. NUCLEAR GRADE 2, MITOSES 2), OVERALL GRADE 2/3.
- C. DUCTAL CARCINOMA IN-SITU (DCIS), CRIBRIFORM TYPE WITH ASSOCIATED COMEDO NECROSIS, NUCLEAR GRADE 2, COMPRISING 5% OF TOTAL TUMOR VOLUME AND IS PRESENT IN ASSOCIATION WITH INVASIVE CARCINOMA.
- D. NO DEFINITIVE LYMPHOVASCULAR INVASION IS SEEN.
- E. MARGINS: THE INVASIVE CARCINOMA EXTENDS TO THE SUPERIOR MARGIN, REMAINDER OF MARGINS ARE FREE.
- F. ATYPICAL DUCTAL HYPERPLASIA, FLAT EPITHELIAL ATYPIA, FIBROCYSTIC CHANGES WITH ASSOCIATED CALCIFICATIONS, DUCTAL EPITHELIAL HYPERPLASIA, FIBROADENOMATOID NODULE AND PREVIOUS BIOPSY SITE CHANGES.
- G. ER-POSITIVE, PR-POSITIVE, HER ZINEU-NEGATIVE (INC SCORE 0), CROSS REFER

PART 2: #1 SENTINEL LYMPH NODE, RIGHT AXILLA, BIOPSY -

- A. ONE LYMPH NODE POSITIVE FOR MICRO METASTATIC CARCINOMA (1/1).
- THE METASTATIC FOCUS MEASURES 1MM (ON GLASS SLIDE), NO EXTRA CAPSULAR EXTENSION IS IDENTIFIED.

PART 3: #2 SENTINEL LYMPH NODE, RIGHT AXILLA, BIOPSY -ONE BENIGN LYMPH NODE, NO TUMOR SEEN (0/1).

Carcinoma, injultrating ductal, Nos 8500/3 Site: breat, NOS C50.9 3/13/11 lu

UUID:D644B589-A6ED-4DCC-97D4-751E22AE3A7D

N LANCO E PROPINCIO EN PROPINCIO DE PROPINCI DE PROPINCIO DE PROPINCIO DE PROPINCIO DE PROPINCIO DE PROPINCIO

Redacted

TCGA-BH-A0DQ-01A-PR

PART 4: #1 NON-SENTINEL LYMPH NODE, RIGHT AXILLA, BIOPSY ONE BENIGN LYMPH NODE, NO TUMOR SEEN (0/1).

SYNOPTIC - PRIMARY INVASIVE CARCINOMA OF BREAST

LATERALITY: PROCEDURE:

Rlaht

LOCATION:

Segmental Not specified

SIZE OF TUMOR:

Maximum dimension invasive component: 2.3 cm

MULTICENTRICITY/MULTIFOCALITY OF INVASIVE FOCI:

TUMOR TYPE (Invasive component):

NOTTINGHAM SCORE:

Ductal adenocarcinoma, NOS

Nuclear grade: 2

Tubule formation: 2 Mitotic activity score: 2 Total Nottingham score: 6

Nottingham grade (1, 2, 3): 2

ANGIOLYMPHATIC INVASION:

No

DERMAL LYMPHATIC INVASION:

CALCIFICATION:

Not applicable

TUMOR TYPE, IN SITU:

No Cribriform

Comedo

DCIS admixed with invasive carcinoma

Percent of tumor occupied by in situ component: 5 %

SURGICAL MARGINS INVOLVED BY INVASIVE COMPONENT:

Yes, focal

SURG MARGINS INVOLVED BY IN SITU COMPONENT:

LYMPH NODES POSITIVE:

No

LYMPH NODES EXAMINED:

1

METHOD(S) OF LYMPH NODE EXAMINATION:

SENTINEL NODE METASTASIS:

H/F stain

SIZE OF NODAL METASTASES:

LYMPH NODE METASTASIS(-ES) WITH EXTRACAPSULAR EXTENSION:

Diameter of largest lymph node metastasis: 1 inm METASTASES TO IPSILATERAL INTERNAL MAMMARY LYMPH NODE (IF APPLICABLE):

SKIN INVOLVED (ULCERATION): NON-NEOPLASTIC BREAST TISSUE: ADH T STAGE, PATHOLOGIC:

N STAGE, PATHOLOGIC: M STAGE, PATHOLOGIC:

pT2 pN1mi

ESTROGEN RECEPTORS: PROGESTERONE RECEPTORS:

pMX positive

HER2/NEU: positive

Zero or 1 +

