ventative mastectomy.		
P DATE:		
OCEDURE: right modified radical mast	ectomy, left total mastectomy.	
ECIFIC CLINICAL QUESTION: Not pro	vided. Col	Nection Date.
TSIDE TISSUE DIAGNOSIS: No. IOR MALIGNANCY: No.	-	
EMORADIATION THERAPY: No.		
HER DISEASES: No.		
INAL DIAGNOSIS:		•
RT 1: BREAST RIGHT, MASTECTO		
A. INVASIVE LOBULAR CAR B. NOTTINGHAM GRADE 2.0	cinoma, classical type Tubule Formation: 3, nuclear pleo	MORDINGM. 2 MITOTIC ACTIVITY. 4
TOTAL SCORE: 6/9).		·
C. THE INVASIVE TUMOR MI	EASURES 4.0 CM IN LARGEST DIMENSION	N.
E. LYMPHOVASCULAR SPA	SITU (LCIS) IS ALSO IDENTIFIED.	
F. RESECTION MARGINS AF	RE NEGATIVE FOR CARCINOMA.	
G. INVASIVE CARCINOMA IS	1.0 CM FROM THE NEAREST (POSTERIC NA IS LOCATED AT THE JUNCTION OF U	OR) MARGIN.
OUTER QUADRANT,		PPER OUTER QUADRANT AND LOWER
I. NIPPLE IS NEGATIVE FOR		
J. SKIN IS NEGATIVE FOR T		ROADENOMA, FIBROCYSTIC CHANGES,
COLUMNAR CELL CHANC	ies and pseudoangiomatqus stroi	MAL HYPERPLASIA
L. PREVIOUS BIOPSY SITE (	CHANGES.	
PROGESTERONE RECEP	ELLS ARE POSITIVE FOR ESTROGEN REC FOR (H-ecore 270) AND NEGATIVE FOR H	CEPTUR (H-ecore 270), POSITIVE FOR
PATHOLOGY REPORT		
RT 2: BREAST, LEFT, TOTAL MASI	FCTOMY.	
A. FIBROADENOMA.		
B. COLUMNAR CELL CHANG	es.	
C. FIBROCYSTIC CHANGES. D. PSEUDOANGIOMATOUS!	TRAMA: LIVINGMO: AGIA	•
E. PRIOR BIOPSY SITE CHA		
F. UNREMARKABLE NIPPLE	AND SKIN.	
A. METASTATIC CARCINOM B. THE LARGEST METASTA C. NO EXTRACAPSULAR EX	A, DISSECTION — A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.	NODES (2/18).
A. METASTATIC CARCINOM. B. THE LARGEST METASTA	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.	NODES (2/18).
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX  ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY:	A INVOLVES TWO OF EIGHTEEN LYMPH FICE FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant	NODES (2/18).
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE:	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant	
A. METASTATIC CARCINOM B. THE LARGEST METASTA: C. NO EXTRACAPSULAR EX  ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION:	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invesive componen	nt: 40 mm
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICTTY/MULTIFOCALE	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invesive componen	nt: 40 mm
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (invasive componen	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invesive componen	nt: 40 mm
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension investive componently OF INVASIVE FOCI:  No ti:	nt: 40 mm
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (invasive componen	A INVOLVES TWO OF EIGHTEEN LYMPH ITC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invesive component OF INVASIVE FOCI:  No ti: infiltrating lobular carcinoma Classical Nuclear grade: 2	at 40 mm /CD-0-3 Carcinoma infiltrating Jobular, NOS 8520
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Lower outer quadrant Maximum dimension invasive component OF INVASIVE FOCI:  No ti: Infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubule formation: 3 Mitotic activity score: 1	at 40 mm /CD-0-3 Carcinoma infiltrating Jobular, NOS 8520
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component	A INVOLVES TWO OF EIGHTEEN LYMPH IC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right   Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension investive component OF INVASIVE FOCI: No it: Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitolic activity score: 1 Total Nottingham score: 6	t: 40 mm /CO-0-3 Carcinoma infiltrating Jobular, NOS 8520, Site breast NOS CSO. 9
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Lower outer quadrant Maximum dimension invasive component OF INVASIVE FOCI:  No ti: Infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubule formation: 3 Mitotic activity score: 1	t: 40 mm /CO-0-3 Carcinoma infiltrating Jobular, NOS 8520, Site breast NOS CSO. 9
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component PY OF INVASIVE FOCI: No 1): Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No	t: 40 mm /CO-0-3 Carcinoma infiltrating Jobular, NOS 8520, Site breast NOS CSO. 9
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: CALCIFICATION:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component OF INVASIVE FOCI:  No tit: infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No	t: 40 mm /CO-0-3 Carcinoma infiltrating Jobular, NOS 8520, Site breast NOS CSO. 9
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU:	A INVOLVES TWO OF EIGHTEEN LYMPH IC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension investive component OF INVASIVE FOCI:  No  ti:  infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 6 Nottingham grade (1, 2, 3): 2 Yea No No LCIS	at 40 mm /CD-0-3 Carcinoma infiltrating Jobular, NOS 8520
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: CALCIFICATION:	A INVOLVES TWO OF EIGHTEEN LYMPH IC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension investive component OF INVASIVE FOCI:  No  ti:  infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 6 Nottingham grade (1, 2, 3): 2 Yea No No LCIS	t: 40 mm /CO-0-3 Carcinoma infiltrating Jobular, NOS 8520, Site breast NOS CSO. 9
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component OF INVASIVE FOCI:  No ti: infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS  IY INVASIVE COMPONENT: No Distance of invasive tumor to closest ma:	carcinoma infiltrating lobular, NOS 8520, Site breast NOS CSO. 9  May 100 May
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU:	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component FY OF INVASIVE FOCI: No  ti: Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS FY INVASIVE COMPONENT: No Distance of invasive tumor to closest man No	carcinoma infiltrating lobular, NOS 8520, Site breast NOS CSO. 9  May 100 May
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive POCI:  No  11: infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 6 Nottingham grade (1, 2, 3): 2 Yes No No LCIS YINVASIVE COMPONENT: No Distance of invasive tumor to closest mai No 2 18	carcinoma infiltrating lobular, NOS 8520, Site breast NOS CSO. 9  May 100 May
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive FOCI:  No  1: infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IY INVASIVE COMPONENT: No Distance of invasive tumor to closest main Na 2 18 MINATION:	carcinoma infiltrating lobular, NOS 8520, Site breast NOS CSO. 9  May 100 May
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXAMINED:	A INVOLVES TWO OF EIGHTEEN LYMPH FIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive componen FY OF INVASIVE FOCI: No  th: infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS FY INVASIVE COMPONENT: No Distance of invasive tumor to closest mai Na 2 18 MINATION: H/E stain	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive FOCI:  No  1: infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IY INVASIVE COMPONENT: No Distance of invasive tumor to closest main Na 2 18 MINATION:	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive FOCI:  No  1: infiltrating lobular carcinoma Classical Nuclear grads: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IY INVASIVE COMPONENT: No Distance of invasive tumor to closest main No 2 18 MINATION: H/E stain Diameter of largest lymph node metastas WITH EXTRACAPSULAR EXTENSION:	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES: LYMPH NODE METASTASES: LYMPH NODE VULCERATION:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Maximum dimension invasive component V OF INVASIVE FOCI:  No  It:  Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 6 Nottingham grade (1, 2, 3): 2 Yes No No LCIS Y INVASIVE COMPONENT: No Distance of invasive tumor to closest mai No 2 18 MINATION: H/E stain Diameter of largest lymph node metastas WITH EXTRACAPSULAR EXTENSION: No	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E LYMPH NODES POSITIVE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES: LYMPH NODE METASTASIS(-ES) V SKIN INVOLVED (ULCERATION): NON-NEOPLASTIC BREAST TISSU T STAGE, PATHOLOGIC:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Maximum dimension invasive component V OF INVASIVE FOCI:  No  It:  Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubula formation: 3 Mitotic activity score: 1 Total Nottingham score: 6 Nottingham grade (1, 2, 3): 2 Yes No No LCIS Y INVASIVE COMPONENT: No Distance of invasive tumor to closest mai No 2 18 MINATION: H/E stain Diameter of largest lymph node metastas WITH EXTRACAPSULAR EXTENSION: No	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE C LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED BE PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES: LYMPH	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive POCI:  No  1: Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IY INVASIVE COMPONENT: No Distance of invasive tumor to closest main Na 2 18 MINATION: H/E stain Diameter of largest lymph node metastas WITH EXTRACAPSULAR EXTENSION: No E: Fibroadenoma, FCD pT2 pN1a	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES POSITIVE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASIS(-ES) W SKIN INVOLVED (ULCERATION): NON-NEOPLASTIC BREAST TISSU T STAGE, PATHOLOGIC: M STAGE;	A INVOLVES TWO OF EIGHTEEN LYMPH ITC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component IV OF INVASIVE FOCI: No III: Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IV INVASIVE COMPONENT: No Distance of invasive tumor to closest man Na 2 18 MINATION: H/E stain Diameter of lergest lymph node metastas VITH EXTRACAPSULAR EXTENSION: No No E: Fibroadenoma, FCD pT2 pN1s Not applicable	carcinoma infiltrating lobular, NOS 8520, Sita breast Nes CSO. 9  pur 5/2
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (Invasive component HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E  PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES: LYMPH	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Lower outer quadrant Maximum dimension invasive component of Invasive POCI:  No  1: Infiltrating lobular carcinoma Classical Nuclear grade: 2 Tubule formation: 3 Mitotic activity score: 1 Total Nottingham score: 5 Nottingham grade (1, 2, 3): 2 Yes No No LCIS IY INVASIVE COMPONENT: No Distance of invasive tumor to closest main Na 2 18 MINATION: H/E stain Diameter of largest lymph node metastas WITH EXTRACAPSULAR EXTENSION: No E: Fibroadenoma, FCD pT2 pN1a	citeria   CO-0-3
A. METASTATIC CARCINOM. B. THE LARGEST METASTA' C. NO EXTRACAPSULAR EX ASE SYNOPSIS: SYNOPTIC - PRIMARY INVASIVE O LATERALITY: PROCEDURE: LOCATION: SIZE OF TUMOR: MULTICENTRICITY/MULTIFOCALF TUMOR TYPE (INVASIVE COMPONENT HISTOLOGIC TYPE: NOTTINGHAM SCORE:  ANGIOLYMPHATIC INVASION: DERMAL LYMPHATIC INVASION: CALCIFICATION: TUMOR TYPE, IN SITU: SURGICAL MARGINS INVOLVED E PAGET'S DISEASE OF NIPPLE: LYMPH NODES POSITIVE: LYMPH NODES POSITIVE: LYMPH NODES EXAMINED: METHOD(S) OF LYMPH NODE EXA SIZE OF NODAL METASTASES: LYMPH NODE METASTASES: LYMPH NODE METASTASIS(-ES) V SKIN INVOLVED (ULCERATION): NON-NEOPLASTIC BREAST TISSU T STAGE, PATHOLOGIC: N STAGE, PATHOLOGIC: N STAGE; ESTROGEN RECEPTORS:	A INVOLVES TWO OF EIGHTEEN LYMPH TIC FOCUS MEASURES 1.6 CM. TENSION IS IDENTIFIED.  CARCINOMA OF BREAST Right Modified radical mastectomy Upper outer quadrant Maximum dimension invasive component to the folial control of the contro	carcinoma, infiltrating lobular, NOS 8520, Site: breast Nes CSO. 9  Marin: 10 mm

UUID: 9D040C65-C225-4D41-BA4C-6BB21EC1CD32
TCGA-BH-A28Q-01A-PR
Redacted

Reviewer initials

Task Reviewer

Task Reviewer

Total Discrepancy

Total Discrepancy

Reviewer initials

Task Reviewer

Task Re