

# hw06

November 2, 2021


## 1 Homework 6

Nathan LeRoy

```
[2]: # better image quality
import matplotlib as mpl
%matplotlib inline
mpl.rcParams['figure.dpi'] = 300
```


### 1.1 Problem 1

We can find an analytical solution to the convolution using a graphical approach. Sketching out the two functions we get:



Let  $f(x) = \Pi(\frac{x-1}{3})$  (orange) and  $g(x) = \Pi(\frac{x-3}{2})$  (blue). We can find the solution to  $f(x) * g(x)$  graphically.

Following the procedure for graphically interpreting the convolution, we can fix  $f(x)$  and flip  $g(x)$  about the y axis. Then slide  $g(x)$  over until the functions overlap, and we can infer the bounds of our convolution:



From the graphs we can see that there will be 3 integrals to evaluate:

### 1.1.1 Section 1

Integrate from  $u = -\frac{1}{2}$  to  $u = x - 2$ :

$$\int_{-\frac{1}{2}}^{x-2} f(u)g(x-u)du = \int_{-\frac{1}{2}}^{x-2} (1)(1)du = u \Big|_{-\frac{1}{2}}^{x-2} = x - 2 - -\frac{1}{2} = x - 1\frac{1}{2}$$

The bounds of this equation in  $x$  space can be calculated based on our right edge:

**Left bound:**  $u = -\frac{1}{2} = x - 1 \Rightarrow x = 1\frac{1}{2}$

**Right bound:**  $u = 1\frac{1}{2} = x - 2 \Rightarrow x = 3\frac{1}{2}$

### 1.1.2 Section 2

Integrate from  $u = x - 4$  to  $u = x - 2$ :

$$\int_{x-4}^{x-2} f(u)g(x-u)du = \int_{x-4}^{x-2} (1)(1)du = u \Big|_{x-4}^{x-2} = x - 2 - (x - 4) = 2$$

The bounds of this equation in  $x$  space can be calculated using the edges of the plot:

**Left bound:**  $u = -\frac{1}{2} = x - 4 \Rightarrow 3\frac{1}{2}$

**Right bound:**  $u = 2\frac{1}{2} = x - 2 \Rightarrow 4\frac{1}{2}$

### 1.1.3 Section 3

Integrate from  $u = x - 4$  to  $u = 2\frac{1}{2}$ :

$$\int_{x-4}^{2\frac{1}{2}} f(u)g(x-u)du = \int_{x-4}^{2\frac{1}{2}} (1)(1)du = u \Big|_{x-4}^{2\frac{1}{2}} = 2\frac{1}{2} - (x - 4) = 6\frac{1}{2} - x$$

The bounds of this equation in  $x$  space can be calculated using the edges of the plot:


**Left bound:**  $u = \frac{1}{2} = x - 4 \Rightarrow 4\frac{1}{2}$

**Right bound:**  $u = 2\frac{1}{2} = x - 4 \Rightarrow 6\frac{1}{2}$

Thus, our final piecewise function looks like this:


$$h(x) = \begin{cases} x - 1\frac{1}{2} & 1\frac{1}{2} \leq x \leq 3\frac{1}{2} \\ 2 & 3\frac{1}{2} \leq x \leq 4\frac{1}{2} \\ -x + 6\frac{1}{2} & 4\frac{1}{2} \leq x \leq 6\frac{1}{2} \\ 0 & \text{elsewhere} \end{cases}$$

Here is a graph of the analytical convolution:



## 1.2 Problem 2

Here is a plot of our system:



By intuition, we can see that there will be 6 distinct regions and subsequent integrals to calculate:

1. Initially as we shift  $f_2(x)$  to the right.
2. When  $\Pi_2(x + 2)$  begins to overlap with  $\Pi_1(x - 2)$ .
3. When  $\Pi_2(x + 2)$  begins to **stop** overlapping with  $\Pi_1(x - 2)$ .
4. Initially as we shift  $f_2(x)$  to the left.
5. When  $\Pi_2(x - 2)$  begins to overlap with  $\Pi_1(x + 2)$ .
6. When  $\Pi_2(x - 2)$  begins to **stop** overlapping with  $\Pi_1(x + 2)$ .

### 1.2.1 Section 1 (two integrals):

$$\int_{x-2\frac{1}{2}}^{-1\frac{1}{2}} f(u)g(x-u)du + \int_{x+1\frac{1}{2}}^{2\frac{1}{2}} f(u)g(x-u)du = \int_{x-2\frac{1}{2}}^{-1\frac{1}{2}} (1)(1)du + \int_{x+1\frac{1}{2}}^{2\frac{1}{2}} (1)(1)du$$

$$\left. u \right|_{x-2\frac{1}{2}}^{-1\frac{1}{2}} + \left. u \right|_{x+1\frac{1}{2}}^{2\frac{1}{2}} = -1\frac{1}{2} - (x - 2\frac{1}{2}) + 2\frac{1}{2} - (x + 1\frac{1}{2})$$

$$2 - 2x$$

### 1.2.2 Section 2:

$$\int_{1\frac{1}{2}}^{x-1\frac{1}{2}} f(u)g(x-u)du = \int_{1\frac{1}{2}}^{x-1\frac{1}{2}} (1)(1)du$$

$$\left. u \right|_{1\frac{1}{2}}^{x-1\frac{1}{2}} = 1\frac{1}{2} - (x - 1\frac{1}{2})$$

$$x - 3$$

**1.2.3 Section 3:**

$$\int_{x-2\frac{1}{2}}^{2\frac{1}{2}} f(u)g(x-u)du = \int_{x-2\frac{1}{2}}^{2\frac{1}{2}} f(1)(1)du$$

$$u \Big|_{x-2\frac{1}{2}}^{2\frac{1}{2}} = 2\frac{1}{2} - (x - 2\frac{1}{2})$$

$$5 - x$$

**1.2.4 Section 4 (two integrals):**

$$\int_{-2\frac{1}{2}}^{x-1\frac{1}{2}} f(u)g(x-u)du + \int_{1\frac{1}{2}}^{x+2\frac{1}{2}} f(u)g(x-u)du = \int_{-2\frac{1}{2}}^{x-1\frac{1}{2}} (1)(1)du + \int_{1\frac{1}{2}}^{x+2\frac{1}{2}} (1)(1)du$$

$$u \Big|_{-2\frac{1}{2}}^{x-1\frac{1}{2}} + u \Big|_{1\frac{1}{2}}^{x+2\frac{1}{2}} = (x - 1\frac{1}{2}) - (-2\frac{1}{2}) + (x + 2\frac{1}{2}) - 1\frac{1}{2}$$

$$2x + 2$$

**1.2.5 Section 5:**

$$\int_{x+1\frac{1}{2}}^{-1\frac{1}{2}} f(u)g(x-u)du = \int_{x+1\frac{1}{2}}^{-1\frac{1}{2}} (1)(1)du$$

$$u \Big|_{x+1\frac{1}{2}}^{-1\frac{1}{2}} = -1\frac{1}{2} - (x + 1\frac{1}{2})$$

$$-3 - x$$

**1.2.6 Section 6:**

$$\int_{-2\frac{1}{2}}^{x+2\frac{1}{2}} f(u)g(x-u)du = \int_{-2\frac{1}{2}}^{x+2\frac{1}{2}} (1)(1)du$$


$$u \Big|_{-2\frac{1}{2}}^{x+2\frac{1}{2}} = (x + 2\frac{1}{2}) - (-2\frac{1}{2})$$

$$x + 5$$

Thus, our final piecewise function will look like this:

$$h(x) = \begin{cases} 2 - 2x & 0 \leq x \leq 1 \\ x - 3 & 3 \leq x \leq 4 \\ 5 - x & 4 \leq x \leq 5 \\ 2x + 2 & -1 \leq x \leq 0 \\ -3 - x & -4 \leq x \leq -3 \\ x + 5 & -5 \leq x \leq -4 \\ 0 & \text{elsewhere} \end{cases}$$

This plot looks like this:



### 1.3 Problem 3

We can plot our system in Python:


```
[3]: import matplotlib.pyplot as plt
import numpy as np

locs = np.linspace(-6,6, num=13)
x=[0,0,0,1,1,1,1,1,1,1,1,1,1]
h=[0,0,0,0,0,0,0,0,0,0,0,1,1,1,1]

plt.rcParams["figure.figsize"] = (6,3)
plt.stem(
    locs,
    x,
    linefmt='b-',
    markerfmt="bo",
    basefmt="b-",
    label="$x[n+3]$"
)
plt.stem(
    locs,
    h,
    linefmt='r-',
    markerfmt="ro",
    basefmt="r-",
    label="$h[n-3]$"
)
```

```
)
plt.legend()
```


[3]: <matplotlib.legend.Legend at 0x110246b50>



We can flip  $h[n = 3]$  and begin shifting to the right in increments of  $k$  to intuitively build an analytical solution for our convolution:

```
[4]: plt.rcParams["figure.figsize"] = (6,3)
plt.stem(
    locs,
    x,
    linefmt='b-',
    markerfmt="bo",
    basefmt="b-",
    label="$x[n+3]$"
)
plt.stem(
    locs,
    np.flip(h),
    linefmt='r-',
    markerfmt="ro",
    basefmt="r-",
    label="$h[n-3]$"
)
plt.legend()
```

[4]: <matplotlib.legend.Legend at 0x1103b92b0>



Right off the bat we overlap by one discrete time unit. The total overlapping “area” is  $1 \cdot 1 = 1$ . Shifting over by  $k = 1$  we get a total “area” of  $1 \cdot 1 + 1 \cdot 1 = 2$ . This idea can be extended to build a pattern. Let the convolution  $x[n+3] * h[n-3] = y[n]$ :

$$k = 0 \rightarrow y[n] = 1$$

$$k = 1 \rightarrow y[n] = 2$$

$$k = 2 \rightarrow y[n] = 3$$

$\vdots$

This yields the general solution:

$$y[n] = n + 1$$

## 1.4 Problem 4

### 1.4.1 a.) $m[n] = x[n] * z[n]$

We can plot the functions in python:

```
[5]: locs = np.linspace(-9,3, num=13)
x = [0,0,0,0,1,1,1,1,1,1,1,0,0]
z = np.flip([0,2,2,2,2,2,2,1,1,1,1,0,0,0])

plt.rcParams["figure.figsize"] = (6,3)
plt.stem(
    locs,
    x,
    linefmt='b-',
    markerfmt="bo",
    basefmt="b-",
```




```

        label="$x[n]$"
    )
    plt.stem(
        locs,
        z,
        linefmt='r-',
        markerfmt="ro",
        basefmt="r-",
        label="$z[n]$"
    )
    plt.legend()

```

[5]: <matplotlib.legend.Legend at 0x1104e6190>



We can flip  $z[n]$  about the y axis and then start shifting:

```

[6]: plt.rcParams["figure.figsize"] = (6,3)
    plt.stem(
        locs,
        x,
        linefmt='b-',
        markerfmt="bo",
        basefmt="b-",
        label="$x[n]$"
    )
    plt.stem(
        locs,
        np.flip(z),
        linefmt='r-',
        markerfmt="ro",


```

```

    basefmt="r-",
    label="$z[n]$"
)
plt.legend()

```

[6]: <matplotlib.legend.Legend at 0x11057d970>



As we shift  $z[n]$  for positive and negative values of  $k$ , we finally achieve the following vector:

$$y[n] = [1, 2, 3, 4, 6, 8, 10, 11, 12, 11, 10, 8, 6, 4, 2]$$


Plotted, we achieve the following:

```

[7]: y = [1,2,3,4,6,8,10,11,12,11,10,8,6,4,2]
plt.stem(
    y,
    linefmt='r-',
    markerfmt="ro",
    basefmt="r-",
    label="$y[n]$"
)
plt.legend()

```

[7]: <matplotlib.legend.Legend at 0x1105f3550>



**1.4.2 b.)**  $m[n] = x[n]*y[n]$


We can solve this using python:

```
[9]: x = np.array([0,1,1,1,1,1,1,1,0,0,0,0])
y = np.array([0,0,1,1,1,1,-1,-1,-1,-1,0,0])

z = np.convolve(x,y)

plt.stem(
    z,
    linefmt='r-',
    markerfmt="ro",
    basefmt="r-",
    label="$m[n]$"
)
```

[9]: <StemContainer object of 3 artists>



### 1.5 Problem 5

To find a matrix  $H$  that satisfies the equation  $y = Hx$ , we can start with the definition of a discrete-time convolution:

$$\sum_{k=0}^{\infty} x[k]h[n-k]$$

We can set our lower bound to  $k = 0$  as both equations are 0 for all values  $n < 0$ . From this we can start investigating the values of  $y[n]$ . Note that for all values of  $k > n$ , those summation components will be directly equal to 0. Values of  $k > n$  will evaluate to  $h[-1], h[-2] \dots$  etc, which is, by our definition of  $h[n]$  equal to zero and will evaluate to  $x[n] \cdot 0 = 0$ .

$$n = 0 \rightarrow y[0] = x[0]h[0]$$

$$n = 1 \rightarrow y[0] = x[0]h[1] + x[1]h[0]$$

$$n = 2 \rightarrow y[0] = x[0]h[2] + x[1]h[1] + x[2]h[0] = x[2]h[0] + x[1]h[1] + x[0]h[2]$$

$$n = 3 \rightarrow y[0] = x[0]h[3] + x[1]h[2] + x[2]h[1] + x[3]h[0] = x[3]h[0] + x[2]h[1] + x[1]h[2] + x[0]h[3]$$

$\vdots$

etc

We can see a pattern emerging and build up our vectors:

$$y = \begin{bmatrix} y[0] \\ y[1] \\ y[2] \\ \vdots \\ y[L + M - 1] \end{bmatrix} = \begin{bmatrix} h[0] & 0 & 0 & 0 & \dots & 0 \\ h[1] & h[0] & 0 & 0 & \dots & 0 \\ h[2] & h[1] & h[0] & 0 & \dots & 0 \\ \vdots & & & & & \\ h[M-1] & h[M-2] & h[M-3] & \dots & \dots & 0 \\ 0 & \dots & \dots & 0 & h[M-1] & h[0] \end{bmatrix} \begin{bmatrix} x[0] \\ x[1] \\ x[2] \\ \vdots \\ x[L-1] \end{bmatrix}$$

We can see now that we have a general solution to the originally proposed equation:

$$y = Hx$$