

wwPDB NMR Structure Validation Summary Report (i)

Jul 2, 2020 – 12:17 AM CDT

PDB ID : 2KKO

Title: Solution NMR structure of the homodimeric winged helix-turn-helix DNA-

binding domain (fragment 1-100) Mb0332 from Mycobacterium bovis, a possible ArsR-family transcriptional regulator. Northeast Structural Genomics

Consortium Target MbR242E.

Authors: Ramelot, T.A.; Cort, J.R.; Wang, D.; Ciccosanti, C.; Jiang, M.; Nair, R.;

Rost, B.; Swapna, G.; Acton, T.B.; Xiao, R.; Everett, J.K.; Montelione, G.T.;

Kennedy, M.A.; Northeast Structural Genomics Consortium (NESG)

Deposited on : 2009-06-26

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)

NmrClust : Kelley et al. (1996)

MolProbity: 4.02b-467

Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.6.dev1

BMRB Restraints Analalysis : v1.2

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.6.dev1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 63%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m NMR}$ archive $(\#{ m Entries})$
Clashscore	136327	12091
Ramachandran outliers	132723	10835
Sidechain outliers	132532	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	108	84%	•	14%
1	В	108	83%	•	14%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mode				
1	A:7-A:99, B:8-B:100 (186)	0.55	1	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 5, 7, 8, 10, 12, 13, 14, 15, 17, 18, 20
2	6, 9, 16
3	4, 11
Single-model clusters	2; 19



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3328 atoms, of which 1678 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called POSSIBLE TRANSCRIPTIONAL REGULATORY PROTEIN (POSSIBLY ARSR-FAMILY).

Mol	Chain	Residues		Atoms				Trace	
1	Λ	108	Total	С	Н	N	О	S	0
1	A	108	1664	505	839	165	153	2	U
1	D	108	Total	С	Н	N	О	S	0
1	В	100	1664	505	839	165	153	2	

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	LEU	-	EXPRESSION TAG	UNP Q7U294
A	102	GLU	-	EXPRESSION TAG	UNP Q7U294
A	103	HIS	-	EXPRESSION TAG	UNP Q7U294
A	104	HIS	-	EXPRESSION TAG	UNP Q7U294
A	105	HIS	-	EXPRESSION TAG	UNP Q7U294
A	106	HIS	-	EXPRESSION TAG	UNP Q7U294
A	107	HIS	-	EXPRESSION TAG	UNP Q7U294
A	108	HIS	-	EXPRESSION TAG	UNP Q7U294
В	101	LEU	-	EXPRESSION TAG	UNP Q7U294
В	102	GLU	-	EXPRESSION TAG	UNP Q7U294
В	103	HIS	-	EXPRESSION TAG	UNP Q7U294
В	104	HIS	-	EXPRESSION TAG	UNP Q7U294
В	105	HIS	-	EXPRESSION TAG	UNP Q7U294
В	106	HIS	-	EXPRESSION TAG	UNP Q7U294
В	107	HIS	-	EXPRESSION TAG	UNP Q7U294
В	108	HIS	-	EXPRESSION TAG	UNP Q7U294



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: POSSIBLE TRANSCRIPTIONAL REGULATORY PROTEIN (POSSIBLY ARSR-FAMILY)



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: POSSIBLE TRANSCRIPTIONAL REGULATORY PROTEIN (POSSIBLY ARSR-FAMILY)



• Molecule 1: POSSIBLE TRANSCRIPTIONAL REGULATORY PROTEIN (POSSIBLY ARSR-FAMILY)









Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 150 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.20
CNS	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2kko_nmr.cif
Number of chemical shift lists	1
Total number of shifts	2501
Number of shifts mapped to atoms	1563
Number of unparsed shifts	386
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	552
Assignment completeness (well-defined parts)	63%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	697	730	730	6±3
1	В	696	725	724	6±3
All	All	27860	29100	29080	198

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 150 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	$egin{aligned} \mathbf{Models} \end{aligned}$	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:B:38:GLU:HG2	1:B:80:ARG:HB3	0.86	1.48	12	2
1:A:24:ASN:HB2	1:B:16:ALA:HA	0.71	1.60	4	1
1:A:62:LEU:HA	1:A:66:GLY:HA3	0.68	1.62	15	2
1:A:69:GLU:HB2	1:A:82:ALA:HA	0.66	1.65	13	3
1:B:22:LEU:HG	1:B:28:LEU:HD21	0.66	1.68	2	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR



entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed Outliers		Percentiles
1	A	93/108 (86%)	91±1 (97±1%)	2±1 (2±1%)	0±0 (0±0%)	47 81
1	В	93/108 (86%)	91±1 (98±1%)	2±1 (2±1%)	0±0 (0±1%)	40 79
All	All	3720/4320 (86%)	3630 (98%)	77 (2%)	13 (0%)	47 81

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	24	ASN	5
1	В	24	ASN	5
1	В	25	GLY	1
1	В	23	ALA	1
1	A	74	GLY	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	67/80 (84%)	65±2 (97±3%)	2±2 (3±3%)	45	88
1	В	67/80 (84%)	65±1 (97±2%)	2±1 (3±2%)	47	89
All	All	2680/3200 (84%)	2594 (97%)	86 (3%)	46	89

5 of 37 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	В	38	GLU	6
1	A	24	ASN	6
1	A	38	GLU	5
1	A	69	GLU	5
1	В	69	GLU	4



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 63% for the well-defined parts and 59% for the entire structure.

7.1 Chemical shift list 1

File name: 2kko nmr.cif

Chemical shift list name: nef_chemical_shift_list_2kko.mr

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2501
Number of shifts mapped to atoms	1563
Number of unparsed shifts	386
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	552
Number of shift outliers (ShiftChecker)	4

The following errors were found when reading this chemical shift list.

• Chemical shift has been reported more than once. First 5 (of 386) occurrences are reported below.

Shift ID	Chain	Res	Type	Atom		Shift Dat	a
	Chain	nes	Type	Atom	Value	Uncertainty	Ambiguity
7	A	2	ALA	HB%	1.410	0.020	1
8	A	2	ALA	HB%	1.410	0.020	1
15	A	3	GLY	HA%	4.030	0.020	1
26	A	4	GLN	HG%	2.390	0.020	1
45	A	6	ASP	HB%	2.710	0.020	1

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atoms found in structure. First 5 (of 552) occurrences are reported below.

Chain Ros Type		Atom	n Shift Data Value Uncertainty Amb				
	laili	rtes	туре	Atom	Value	Uncertainty	Ambiguity
	A	100	HIS	HBy	3.42	0.02	2
	В	58	ASN	HD2y	6.68	0.02	2

Continued on next page...



Continued from previous page...

Chain	Pag	Tuno	Atom		Shift Dat	a
Chain	nes	туре	Atom	Value	Uncertainty	Ambiguity
A	62	LEU	HBx	1.19	0.02	2
A	45	ALA	HB%	1.4	0.02	1
A	84	GLU	HG%	2.35	0.02	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	210	-0.46 ± 0.04	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	192	0.10 ± 0.08	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	205	-0.28 ± 0.06	None needed ($< 0.5 \text{ ppm}$)
^{15}N	206	0.05 ± 0.23	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 63%, i.e. 1406 atoms were assigned a chemical shift out of a possible 2231. 42 out of 42 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	914/930 (98%)	356/372 (96%)	372/372 (100%)	186/186 (100%)
Sidechain	466/1243 (37%)	68/713 (10%)	374/449 (83%)	24/81 (30%)
Aromatic	$26/58 \; (45\%)$	13/30 (43%)	13/26 (50%)	0/2 (0%)
Overall	1406/2231 (63%)	437/1115 (39%)	759/847 (90%)	210/269 (78%)

7.1.4 Statistically unusual chemical shifts $\stackrel{ ext{(i)}}{ ext{(i)}}$

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	В	1	MET	СВ	76.00	44.20 - 21.80	19.2
1	A	1	MET	СВ	76.00	44.20 - 21.80	19.2
1	В	1	MET	CA	73.50	67.38 - 44.88	7.7
1	A	1	MET	CA	73.50	67.38 - 44.88	7.7




7.1.5 Random Coil Index (RCI) plots (i)

The images below report random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:





8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2612
Intra-residue ($ i-j =0$)	570
Sequential (i-j =1)	590
Medium range ($ i-j >1$ and $ i-j <5$)	746
Long range (i-j ≥5)	580
Inter-chain	126
Total dihedral-angle restraints	336
Total hydrogen bond restraints	456
Total disulfide bond restraints	0
Number of unmapped restraints	0
Number of restraints per residue	13.1
Number of long range restraints per residue	2.9

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	2.8	0.2
0.2-0.5 (Medium)	0.3	0.3
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation.



Bins $(^{\circ})$	Average number of violations per model	$\mathrm{Max}\;(^\circ)$
1.0-10.0 (Small)	2.1	3.2
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis

9.1 Summary of distance violations


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Postpoints type	Count	$\%^{1}$	Vio	lated	.3	Consis	tentl	${f y}$ Violated 4
Restraints type	Count	/0	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	570	21.8	11	1.9	0.4	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	482	18.5	9	1.9	0.3	0	0.0	0.0
Sidechain-Sidechain	88	3.4	2	2.3	0.1	0	0.0	0.0
Sequential (i-j =1)	590	22.6	9	1.5	0.3	0	0.0	0.0
Backbone-Backbone	156	6.0	2	1.3	0.1	0	0.0	0.0
Backbone-Sidechain	390	14.9	5	1.3	0.2	0	0.0	0.0
Sidechain-Sidechain	44	1.7	2	4.5	0.1	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	746	28.6	9	1.2	0.3	0	0.0	0.0
Backbone-Backbone	348	13.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	276	10.6	5	1.8	0.2	0	0.0	0.0
Sidechain-Sidechain	122	4.7	4	3.3	0.2	0	0.0	0.0
Long range ($ i-j \ge 5$)	580	22.2	14	2.4	0.5	0	0.0	0.0
Backbone-Backbone	78	3.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	188	7.2	6	3.2	0.2	0	0.0	0.0
Sidechain-Sidechain	314	12.0	8	2.5	0.3	0	0.0	0.0
Inter-chain	126	4.8	2	1.6	0.1	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	62	2.4	2	3.2	0.1	0	0.0	0.0
Sidechain-Sidechain	64	2.5	0	0.0	0.0	0	0.0	0.0
Total	2612	100.0	45	1.7	1.7	0	0.0	0.0
Backbone-Backbone	582	22.3	2	0.3	0.1	0	0.0	0.0
Backbone-Sidechain	1398	53.5	27	1.9	1.0	0	0.0	0.0
Sidechain-Sidechain	632	24.2	16	2.5	0.6	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

9.2 Distance violation statistics for each model

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	3	M (8)	M (SD^6 (Å)
Model ID	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)
1	0	1	0	0	0	1	0.18	0.18	0.0
2	1	1	1	1	0	4	0.15	0.16	0.02
3	1	2	0	1	0	4	0.2	0.28	0.08
4	0	2	1	1	0	4	0.13	0.15	0.01
5	2	0	0	0	0	2	0.15	0.17	0.03
6	2	0	0	0	0	2	0.11	0.11	0.0
7	0	0	1	2	0	3	0.11	0.11	0.0
8	0	0	1	1	0	2	0.14	0.15	0.01
9	0	1	0	0	0	1	0.13	0.13	0.0
10	0	0	0	1	0	1	0.13	0.13	0.0
11	1	1	2	2	2	8	0.14	0.2	0.03
12	4	0	0	0	0	4	0.14	0.18	0.03

Continued on next page...




Continued from previous page...

Madal ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (A)	$ SD^*(A) $
13	1	0	0	3	0	4	0.14	0.18	0.02
14	0	1	0	1	0	2	0.23	0.25	0.02
15	2	0	1	2	0	5	0.18	0.3	0.08
16	0	1	0	0	0	1	0.18	0.18	0.0
17	2	0	2	0	0	4	0.15	0.21	0.04
18	0	0	2	1	0	3	0.12	0.14	0.01
19	3	0	0	1	0	4	0.12	0.13	0.01
20	2	0	0	2	0	4	0.15	0.17	0.02

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model



The mean and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2567(IR:559, SQ:581, MR:737, LR:566, IC:124) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
7	8	7	10	2	34	1	5.0
0	1	2	3	0	6	2	10.0
2	0	0	1	0	3	3	15.0
2	0	0	0	0	2	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble




9.4 Most violated distance restraints in the ensemble

9.4.1 Histogram: Distribution of mean distance violations

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints

The following table provides the mean and the standard deviation of the top ten absolute value of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)
(1,968)	1:A:71:ARG:HD2	1:A:71:ARG:HH11	4	0.12	0.01
(1,968)	1:A:71:ARG:HD2	1:A:71:ARG:HH12	4	0.12	0.01
(1,968)	1:A:71:ARG:HD2	1:A:71:ARG:HH21	4	0.12	0.01
(1,968)	1:A:71:ARG:HD2	1:A:71:ARG:HH22	4	0.12	0.01
(1,968)	1:A:71:ARG:HD3	1:A:71:ARG:HH11	4	0.12	0.01
(1,968)	1:A:71:ARG:HD3	1:A:71:ARG:HH12	4	0.12	0.01
(1,968)	1:A:71:ARG:HD3	1:A:71:ARG:HH21	4	0.12	0.01
(1,968)	1:A:71:ARG:HD3	1:A:71:ARG:HH22	4	0.12	0.01
(1,2097)	1:B:71:ARG:HD2	1:B:71:ARG:HH11	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD2	1:B:71:ARG:HH12	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD2	1:B:71:ARG:HH21	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD2	1:B:71:ARG:HH22	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD3	1:B:71:ARG:HH11	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD3	1:B:71:ARG:HH12	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD3	1:B:71:ARG:HH21	4	0.13	0.02
(1,2097)	1:B:71:ARG:HD3	1:B:71:ARG:HH22	4	0.13	0.02

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)
(1,1448)	1:B:26:ARG:H	1:B:26:ARG:HG2	3	0.19	0.08
(1,1448)	1:B:26:ARG:H	1:B:26:ARG:HG3	3	0.19	0.08
(1,1408)	1:B:21:ALA:HB1	1:B:65:GLY:H	3	0.13	0.01
(1,1408)	1:B:21:ALA:HB2	1:B:65:GLY:H	3	0.13	0.01
(1,1408)	1:B:21:ALA:HB3	1:B:65:GLY:H	3	0.13	0.01
(1,1275)	1:B:8:LYS:HA	1:B:8:LYS:HD2	3	0.19	0.02
(1,1275)	1:B:8:LYS:HA	1:B:8:LYS:HD3	3	0.19	0.02
(1,897)	1:A:67:LEU:HD11	1:A:86:VAL:HA	2	0.11	0.0
(1,897)	1:A:67:LEU:HD12	1:A:86:VAL:HA	2	0.11	0.0
(1,897)	1:A:67:LEU:HD13	1:A:86:VAL:HA	2	0.11	0.0
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD21	2	0.26	0.01
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD22	2	0.26	0.01
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD23	2	0.26	0.01
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD21	2	0.26	0.01
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD22	2	0.26	0.01
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD23	2	0.26	0.01
(1,310)	1:A:24:ASN:H	1:A:28:LEU:HD11	2	0.14	0.01
(1,310)	1:A:24:ASN:H	1:A:28:LEU:HD12	2	0.14	0.01
(1,310)	1:A:24:ASN:H	1:A:28:LEU:HD13	2	0.14	0.01
(1,299)	1:A:22:LEU:HD21	1:A:28:LEU:HD21	2	0.11	0.0
(1,299)	1:A:22:LEU:HD21	1:A:28:LEU:HD22	2	0.11	0.0
(1,299)	1:A:22:LEU:HD21	1:A:28:LEU:HD23	2	0.11	0.0
(1,299)	1:A:22:LEU:HD22	1:A:28:LEU:HD21	2	0.11	0.0
(1,299)	1:A:22:LEU:HD22	1:A:28:LEU:HD22	2	0.11	0.0
(1,299)	1:A:22:LEU:HD22	1:A:28:LEU:HD23	2	0.11	0.0
(1,299)	1:A:22:LEU:HD23	1:A:28:LEU:HD21	2	0.11	0.0
(1,299)	1:A:22:LEU:HD23	1:A:28:LEU:HD22	2	0.11	0.0
(1,299)	1:A:22:LEU:HD23	1:A:28:LEU:HD23	2	0.11	0.0
(1,2146)	1:B:76:ARG:HD2	1:B:77:GLN:H	2	0.16	0.0


¹Number of violated models, ²Standard deviation

9.5 All distance violations

9.5.1 Histogram: Distribution of distance violations

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations

The following table lists the top ten absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1448)	1:B:26:ARG:H	1:B:26:ARG:HG2	15	0.3
(1,1448)	1:B:26:ARG:H	1:B:26:ARG:HG3	15	0.3
(1,1966)	1:B:62:LEU:HB2	1:B:67:LEU:HD21	3	0.28
(1,1966)	1:B:62:LEU:HB2	1:B:67:LEU:HD22	3	0.28
(1,1966)	1:B:62:LEU:HB2	1:B:67:LEU:HD23	3	0.28
(1,1966)	1:B:62:LEU:HB3	1:B:67:LEU:HD21	3	0.28
(1,1966)	1:B:62:LEU:HB3	1:B:67:LEU:HD22	3	0.28
(1,1966)	1:B:62:LEU:HB3	1:B:67:LEU:HD23	3	0.28
(1,1437)	1:B:23:ALA:H	1:B:24:ASN:H	3	0.28
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD21	15	0.26
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD22	15	0.26
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD23	15	0.26
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD21	15	0.26
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD22	15	0.26
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD23	15	0.26
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD21	14	0.25

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD22	14	0.25
(1,837)	1:A:62:LEU:HB2	1:A:67:LEU:HD23	14	0.25
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD21	14	0.25
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD22	14	0.25
(1,837)	1:A:62:LEU:HB3	1:A:67:LEU:HD23	14	0.25
(1,2378)	1:B:101:LEU:HD21	1:B:102:GLU:H	14	0.21
(1,2378)	1:B:101:LEU:HD22	1:B:102:GLU:H	14	0.21
(1,2378)	1:B:101:LEU:HD23	1:B:102:GLU:H	14	0.21
(1,1275)	1:B:8:LYS:HA	1:B:8:LYS:HD2	17	0.21
(1,1275)	1:B:8:LYS:HA	1:B:8:LYS:HD3	17	0.21
(1,42)	1:A:16:ALA:HB1	1:B:25:GLY:HA2	11	0.2
(1,42)	1:A:16:ALA:HB1	1:B:25:GLY:HA3	11	0.2
(1,42)	1:A:16:ALA:HB2	1:B:25:GLY:HA2	11	0.2
(1,42)	1:A:16:ALA:HB2	1:B:25:GLY:HA3	11	0.2
(1,42)	1:A:16:ALA:HB3	1:B:25:GLY:HA2	11	0.2
(1,42)	1:A:16:ALA:HB3	1:B:25:GLY:HA3	11	0.2
(1,2382)	1:B:101:LEU:H	1:B:102:GLU:H	16	0.18
(1,2151)	1:B:76:ARG:H	1:B:76:ARG:HG3	12	0.18



10 Dihedral-anlge violation analysis


10.1 Summary of dihedral-angle violations

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

A1 - 4	C	$\%^1$	Vic	\mathbf{lated}^{i}	3	Consis	tentl	${ m ly~Violated^4}$
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\frac{1}{\%}$
PHI	168	50.0	7	4.2	2.1	0	0.0	0.0
PSI	168	50.0	20	11.9	6.0	0	0.0	0.0
Total	336	100.0	27	8.0	8.0	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-anlge restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-anlge type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-anlge violation statistics in each model

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Num	ber o	of violations	Mean (°)	Max (°)	Std. deviation (°)
Wiodei 1D	PHI	PSI	Total	Mean ()	Max ()	sta. deviation ()
1	1	1	2	1.2	1.3	0.1
2	1	3	4	1.27	1.8	0.3
3	0	4	4	2.38	3.2	0.74
4	0	2	2	1.25	1.4	0.15
5	0	2	2	1.4	1.7	0.3
6	1	0	1	1.1	1.1	0.0
7	0	0	0	0.0	0.0	0.0
8	1	0	1	1.2	1.2	0.0
9	0	4	4	1.55	1.8	0.18
10	1	1	2	1.3	1.3	0.0
11	0	2	2	1.2	1.3	0.1
12	1	1	2	1.45	1.8	0.35
13	0	1	1	2.0	2.0	0.0
14	0	4	4	1.55	2.2	0.38
15	0	2	2	1.55	1.7	0.15
16	0	3	3	1.33	1.5	0.17
17	1	1	2	1.4	1.5	0.1
18	3	0	3	1.27	1.6	0.24
19	0	1	1	1.1	1.1	0.0
20	0	0	0	0.0	0.0	0.0



10.2.1 Bar graph: Dihedral violation statistics for each model



The mean and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Violation statistics in the ensemble

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Num	ber o	of violated restraints	Fractio	n of the ensemble
PHI	PSI	Total	Count ¹	%
5	10	15	1	5.0
1	8	9	2	10.0
1	2	3	3	15.0
0	0	0	4	20.0
0	0	0	5	25.0
0	0	0	6	30.0
0	0	0	7	35.0
0	0	0	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0
0	0	0	12	60.0

Continued on next page...




Continued from previous page...

Num	ıber o	of violated restraints	1			
PHI	PSI	Total	Count ¹	%		
0	0	0	13	65.0		
0	0	0	14	70.0		
0	0	0	15	75.0		
0	0	0	16	80.0		
0	0	0	17	85.0		
0	0	0	18	90.0		
0	0	0	19	95.0		
0	0	0	20	100.0		

¹ Number of models with violations

10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble




10.4 Most violated dihedral-anlge restraints

10.4.1 Histogram: Distribution of mean dihedral-anlge violations

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints

The following table provides the mean and the standard deviation of the top ten absolute value of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean (°)	\mathbf{SD}^2 (°)
(1,168)	1:A:100:HIS:N	1:A:100:HIS:CA	1:A:100:HIS:C	1:A:101:LEU:N	3	1.93	0.68
(1,104)	1:A:62:LEU:N	1:A:62:LEU:CA	1:A:62:LEU:C	1:A:63:LYS:N	2	1.95	0.25
(1,116)	1:A:71:ARG:N	1:A:71:ARG:CA	1:A:71:ARG:C	1:A:72:ARG:N	3	1.5	0.22
(1,204)	1:B:27:ARG:N	1:B:27:ARG:CA	1:B:27:ARG:C	1:B:28:LEU:N	2	1.4	0.3
(1,36)	1:A:27:ARG:N	1:A:27:ARG:CA	1:A:27:ARG:C	1:A:28:LEU:N	2	1.35	0.25
(1,106)	1:A:63:LYS:N	1:A:63:LYS:CA	1:A:63:LYS:C	1:A:64:SER:N	2	1.4	0.0
(1,246)	1:B:49:GLY:N	1:B:49:GLY:CA	1:B:49:GLY:C	1:B:50:MET:N	2	1.25	0.15
(1,336)	1:B:100:HIS:N	1:B:100:HIS:CA	1:B:100:HIS:C	1:B:101:LEU:N	2	1.2	0.1
(1,200)	1:B:22:LEU:N	1:B:22:LEU:CA	1:B:22:LEU:C	1:B:23:ALA:N	2	1.15	0.05
(1,32)	1:A:22:LEU:N	1:A:22:LEU:CA	1:A:22:LEU:C	1:A:23:ALA:N	2	1.1	0.0


¹ Number of violated models, ²Standard deviation



10.5 All violated dihedral-angle restraints

10.5.1 Histogram : Distribution of violations

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints

The following table lists the absolute value of the top ten violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,272)	1:B:62:LEU:N	1:B:62:LEU:CA	1:B:62:LEU:C	1:B:63:LYS:N	3	3.2
(1,168)	1:A:100:HIS:N	1:A:100:HIS:CA	1:A:100:HIS:C	1:A:101:LEU:N	3	2.9
(1,104)	1:A:62:LEU:N	1:A:62:LEU:CA	1:A:62:LEU:C	1:A:63:LYS:N	14	2.2
(1,274)	1:B:63:LYS:N	1:B:63:LYS:CA	1:B:63:LYS:C	1:B:64:SER:N	3	2.1
(1,118)	1:A:72:ARG:N	1:A:72:ARG:CA	1:A:72:ARG:C	1:A:73:GLU:N	13	2.0
(1,222)	1:B:36:GLN:N	1:B:36:GLN:CA	1:B:36:GLN:C	1:B:37:GLY:N	9	1.8
(1,131)	1:A:78:TYR:C	1:A:79:TYR:N	1:A:79:TYR:CA	1:A:79:TYR:C	12	1.8
(1,116)	1:A:71:ARG:N	1:A:71:ARG:CA	1:A:71:ARG:C	1:A:72:ARG:N	2	1.8
(1,204)	1:B:27:ARG:N	1:B:27:ARG:CA	1:B:27:ARG:C	1:B:28:LEU:N	5	1.7
(1,104)	1:A:62:LEU:N	1:A:62:LEU:CA	1:A:62:LEU:C	1:A:63:LYS:N	15	1.7

