



# Supporting Information:

## NMRlipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PE and PG lipids

Pavel Buslaev,<sup>†</sup> Fernando Favela-Rosales,<sup>‡</sup> Patrick Fuchs,<sup>¶</sup> Matti Javanainen,<sup>§</sup>  
Jesper J. Madsen,<sup>||,⊥</sup> Josef Melcr,<sup>§, #</sup> Markus S. Miettinen,<sup>@</sup> O. H. Samuli  
Ollila,<sup>\*,△</sup> Chris G. Papadopoulos,<sup>∇</sup> Antonio Peón,<sup>††</sup> Thomas J. Piggot,<sup>‡‡</sup> and  
Pierre Poulain<sup>¶</sup>

<sup>†</sup>*University of Jyväskylä*

<sup>‡</sup>*Departamento de Investigación, Tecnológico Nacional de México, Campus Zacatecas  
Occidente, México*

<sup>¶</sup>*Paris, France*

<sup>§</sup>*Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences,  
Flemingovo nám. 542/2, CZ-16610 Prague 6, Czech Republic*

<sup>||</sup>*Department of Chemistry, The University of Chicago, Chicago, Illinois, United States of  
America*

<sup>⊥</sup>*Department of Global Health, College of Public Health, University of South Florida, Tampa,  
Florida, United States of America*

<sup>#</sup>*Groningen Biomolecular Sciences and Biotechnology Institute and The Zernike Institute  
for Advanced Materials, University of Groningen, 9747 AG Groningen, The Netherlands*

<sup>@</sup>*Department of Theory and Bio-Systems, Max Planck Institute of Colloids and Interfaces,  
14424 Potsdam, Germany*

<sup>△</sup>*Institute of Biotechnology, University of Helsinki*

<sup>∇</sup>*I2BC - University Paris Sud  
S2*

<sup>††</sup>*Spain*

<sup>‡‡</sup>*Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, United*

# S1 R-PDLF and SDROSS experiments



Figure S1: (A) Chemical structure of POPE with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPE sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar slices from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.

1.A, B etc. labels to be put in the figure.



Figure S2: (A) Chemical structure of POPG with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPG sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDPLF spectra (D) Dipolar slices from the 2D R-PDPLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.




Figure S3: Simpson simlaton of S-DROSS curve of  $\beta$ -carbon of POPG.

## **S2 Comparison of headgroup order parameters from different force fields to experiments**

The quality of PE and PG headgroup conformational ensembles in different simulations against NMR experiments is evaluated in figures S4 and S5 using C-H bond order parameters as in our previous studies for PC and PS lipids.<sup>1,2</sup> Conclusions are the same for all lipids: None of the force fields correctly captures the lipid headgroup conformational ensembles, but CHARMM36 gives results closest to experiments.




Figure S4: The headgroup and glycerol backbone order parameters of PE lipids from experiments (POPE and signs this work, DPPE from Ref. 3 and E.coliPE from Ref. 4) and simulations with different force fields.

2.This should be clarified as in NMRlipidsI and error bars should be added. Probably larger error bars for united atom models based on the report by Fuchs et al.




Figure S5: The headgroup and glycerol backbone order parameters of PG lipids from experiments (POPG and signs from this work and from Ref. 5, DPPG with 100mM NaCl from Ref. 6, and E. Coli PG results from Ref. 4). and simulations with different force fields.



## S3 Lipid headgroups in mixtures with PC and PE or PG lipids

Headgroup order parameters of PC lipids are unchanged upon addition of zwitterionic lipids or cholesterol in experiments, but increase upon addition of negatively charged PG or PS lipids because headgroup dipole tilts more parallel to the membrane plane after incorporation of negative charges into the membrane.<sup>7,10,11</sup> The response of PC headgroup order parameters to the addition of PE or PG lipids from different simulations is compared with experiments in figure S6. by the tested CHARMM36 and Berger-OPLS force fields, although CHARMM36 slightly overestimates the changes. The good performance of Berger-OPLS simulations is notable because the response of headgroup order parameters to cholesterol was significantly overestimated by the Berger/HÅultje force field in our previous work.<sup>1</sup>

None of the force fields fully reproduces the PC headgroup order parameter response to the increasing amount of PG, which may be related to the counterion binding affinity (see also the next section).<sup>2</sup> In all force fields except Slipids, the order parameters of different hydrogens attached to the  $\alpha$ -carbon are responding differently when mixed with PE or PG lipids. 4.Maybe we should figure out what is the reason for this?

Maybe we should analyze the P-N vector angle from different simulations?

<https://github.com/NMRLipids/NMRLipidsIVPEandPG/issues/10>.

For  $\beta$ -carbon order parameter in PG headgroup, experiments report mild increase<sup>8</sup> or no change<sup>5</sup> upon addition of PC lipids (Fig. S7). Simulations with all the tested force fields give only very small changes also for the  $\alpha$ -carbon order parameter (Figs. S11 and S7). Therefore, the simulations are generally in line with experiments, suggesting that the interactions with PC do not essentially effect the PG headgroup structure. This suggests that the interactions between PG and PC headgroups are captured better in simulations than for PS headgroup, where all the force fields significantly overestimated the structural response of PS headgroup to the interactions with PC lipids.<sup>11</sup>




Figure S6: Modulation of POPC headgroup order parameters with increasing amount of POPE (left) and POPG (right) in bilayer from experiments<sup>7,8</sup> and simulations with different force fields. Signs are determined as discussed in.<sup>1,9</sup>

**3.P-N vector angles from Berger-OPLS and GROMOS-CKP simulations are yet to be analyzed.**




Figure S7: Modulation of PG lipid headgroup order parameters with the increasing amount of PC in lipid bilayer from experiments<sup>5,8</sup> and simulations with different force fields.

**5.P-N angles from Slipids and GROMOS-CKP yet to be calculated.**

### S3.1 Sodium binding to PE and PG lipid bilayers

Sodium binding affinity to PE lipids has not been measured experimentally, but large differences to PC would be surprising. In simulations, the sodium binding affinity to POPE depends on the used force field (Fig. S8), but lesser extend than reported previously for PC.<sup>12</sup>

6. This will be finished once we have all the simulation details and Lipid17 simulations with correct dihedrals from issue <https://github.com/NMRLipids/NMRLipidsIVPEandPG/issues/12>, Because some simulation and ion parameters are not identical with the previous work,<sup>12</sup> we compare POPE results to the POPC simulations ran with identical parameters (Fig. S12). In Lipid17 with the strongest sodium binding affinity to POPE, the binding affinity is approximately similar to POPC. Slipids and CHARMM36 exhibit slightly, and GROMOS-CKP substantially weaker binding to POPE than to POPC. Assuming that the binding to POPE would be similar than to POPC, the sodium binding affinity to POPE is potentially realistic in CHARMM36, Slipids, and GROMOS-CKP simulations here, but substantially overestimated in Lipid17 simulation.

Simulations with PG lipids give similar dependence on force field as observed in POPE simulations: Lipid 17 simulations with Dang ion parameters exhibits stronger counter-ion binding affinity to pure POPG bilayer than CHARMM36, Slipids, and GROMOS-CKP simulations, which are roughly similar (Fig. S9). Lipid17 also exhibits less increase in POPC headgroup order parameters upon addition of POPG than other simulations (Fig. S6), and lower area per molecule ( $59.5 \text{ \AA}^2$ ) than in experiments ( $66.1 \text{ \AA}^2$ ). In our previous study about PS lipids,<sup>11</sup> such behaviour was related to the overestimated counterions binding and shielding the electrostatic repulsion between PG headgroups in bilayers. Even though the area per lipid in CHARMM36, Slipids, and GROMOS-CKP simulations is in good agreement with experiments (Fig. S9), the experimental increase in POPC headgroup order parameters upon addition of POPG are not fully reproduced (Fig. S6). Therefore we conclude that the counter-ion binding affinity is overestimated in Lipid17 simulations, while the other simulations are more realistic, but slight overbinding cannot be excluded.




Figure S8: Sodium (solid line) and chloride ion density profiles along membrane normal from different simulations with PE lipids.




Figure S9: Counterion densities and area per lipids from simulations with PG lipids. Experimental area for POPG at 303 K is 66.1  $\text{\AA}^2$  and 67  $\text{\AA}^2$  for DPPC at 323 K.<sup>13</sup>

## S4 Calcium density distributions




Figure S10: Calcium ion density profiles along membrane normal from simulations of POPC:POPG (1:1) mixtures with different force fields.

## S5 Changes of PG headroup order parameters upon addition of PC




Figure S11: Modulation of PG lipid headgroup order parameters with the increasing amount of PC in lipid bilayer from experiments<sup>5,8</sup> and simulations with different force fields.



## S6 Sodium binding to POPC simulations





Figure S12: Sodium (solid line) and chloride ion density profiles along membrane normal from different simulations with PC lipids.

**7. Discussion about differences to the NMRlipids II to be discussed once we have the details on ions models.**

## S7 Calcium binding to POPC:POPG (4:1) mixtures



8.Lipid17ecc data to be analyzed and added.




Figure S14: Calcium ion density profiles along membrane normal from simulations of POPC:POPG (4:1) mixtures with different force fields.




Figure S15: Changes in POPC CHARMM36 dihedrals with increasing amount of CaCl<sub>2</sub>.




Figure S16: Changes in POPC lipid17ecc dihedrals with increasing amount of  $\text{CaCl}_2$ .




Figure S17: Changes in POPG lipids dihedrals with increasing amount of  $\text{CaCl}_2$ .




Figure S18: Changes in POPG lipid17 dihedrals with increasing amount of  $\text{CaCl}_2$ .

## S8 Changes in headgroup conformations upon addition of $\text{CaCl}_2$

## S9 Simulated systems

### S9.1 CHARMM36

*POPE* [33.Simulation details by M. Javanainen.](#)

*POPE with additional NaCl* [34.Simulation details by A. Peon.](#)

*POPG* [35.Simulation details by Ollila.](#)

*POPG with additional NaCl* [36.Simulation details by A. Peon.](#)

*POPC:POPE mixtures* Data is available at.<sup>64,65</sup> 300 K with v-rescale ( $\tau=0.1$  ps), 1 bar with PR semiisotropic ( $\tau=4$  ps, compressibility= $4.5\text{e-}5$  bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [37.Full simulation details by Fuchs et al.](#)

*POPC:POPG mixture with additional calcium* [38.Simulation details by J. Madsen.](#)

*POPC:POPG mixture with additional NaCl* [39.Simulation details by A. Peon.](#)

### S9.2 CHARMM36ua

*POPE* Data is available at.<sup>17</sup> [40.Simulation details by T. Piggot.](#)

### S9.3 Slipids

*POPE* Data is available at.<sup>20</sup> [41.Simulation details by T. Piggot.](#)

*POPE with additional NaCl* [42.Simulation details by A. Peon.](#) I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.

*DPPE* Data is available at.<sup>19</sup> [43.Simulation details by F. Favela.](#)

*POPG* Data is available at.<sup>46</sup> [44.Simulation details by F. Favela.](#) I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.



Table S1: List of MD simulations with PE lipids.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	<sup>a</sup> N <sub>l</sub>	<sup>b</sup> N <sub>w</sub>	<sup>c</sup> N <sub>c</sub>	<sup>d</sup> T (K)	<sup>e</sup> t <sub>sim</sub> (ns)	<sup>f</sup> t <sub>anal</sub> (ns)	<sup>g</sup> files
POPE	CHARMM36 <sup>?</sup>	0	144	5760	0	310	500	400	14
POPE	CHARMM36 <sup>?</sup>	0	500	25000	0	310	500	100	15
POPE	CHARMM36 <sup>?</sup>	0.11	500	25000	50	310	500	100	16
POPE	CHARMM36ua <sup>?</sup>	0	336	15254	0	310	2×200	2×100	17
DPPE	Slipids <sup>18</sup>	0	288	9386	0	336	200	100	19
POPE	Slipids <sup>18</sup>	0	336	?	0	310	2×200	2×100	20
POPE	Slipids <sup>18</sup>	0	500	25000	0	310	500	100	21
POPE	Slipids / Åqvist <sup>18,22</sup>	0.11	500	25000	50	310	500	100	23
DPPE	GROMOS-CKP <sup>?</sup>	0	128	3655	0	342	2×500	2×400	24
POPE	GROMOS-CKP <sup>?</sup>	0	128	3552	0	313	2×500	2×400	25
POPE	GROMOS-CKP <sup>?</sup>	0	500	25000	0	310	500	100	26
POPE	GROMOS-CKP <sup>?</sup>	0.11	500	25000	50	310	500	100	27
DOPE	GROMOS-CKP <sup>?</sup>	0	128	4789	0	271	2×500	2×400	28
POPE	GROMOS 43A1-S3 <sup>?</sup>	0	128	3552	0	313	2×200	2×100	29
POPE	OPLS-UA vdW on H <sup>?</sup>	0	128	3328	0	303	2×200	2×100	30
POPE	OPLS-UA <sup>?</sup>	0	128	3328	0	303	2×200	2×100	31
POPE	OPLS-MacRog <sup>32</sup>	0	144	5760	0	310	500	350	33
POPE	OPLS-MacRog <sup>32</sup>	0	128	5120	0	300	500	300	34
POPE	Berger-Vries <sup>?</sup>	0	128	3552	0	303	2×200	2×100	35
POPE	Berger-largeH <sup>?</sup>	0	128	3552	0	303	2×200	2×100	36
DOPE	Berger-Vries <sup>?</sup>	0	128	4789	0	271	2×200	2×100	37
DOPE	Berger-largeH <sup>?</sup>	0	128	4789	0	271	2×300	2×100	38
POPE	LIPID17 <sup>39</sup>	0	500	25000	50	310	500	100	40
POPE	LIPID17 <sup>39</sup>	0.11	500	25000	50	310	500	100	41

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

9.Citation for CHARMM36 PE?

10.Which ion model is used in <sup>16</sup>?

11.Citation for GROMOS-CKP?

12.Citation for GROMOS 43A1-S3?

13.Citation for OPLS-UA models?

14.Citations for Berger-\* simulations?

15.LIPID17 simulations with correct dihedrals still coming

Table S2: List of MD simulations with PG lipids.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	<sup>a</sup> N <sub>l</sub>	<sup>b</sup> N <sub>w</sub>	<sup>c</sup> N <sub>c</sub>	<sup>d</sup> T (K)	<sup>e</sup> t <sub>sim</sub> (ns)	<sup>f</sup> t <sub>anal</sub> (ns)	<sup>g</sup> files
POPG/K <sup>+</sup>	CHARMM36? <sup>16</sup>	0	118	4110	0	298	100	100	42
POPG	CHARMM36?	0.11	500	25000	49	310	500	100	43
POPG	CHARMM36?	0	500	25000	0	310	500	100	44
POPG/Na <sup>+</sup>	Slipids / Åqvist <sup>22,45</sup>	0	288	10664	0	298	250	100	46
DPPG/Na <sup>+</sup>	Slipids / Åqvist <sup>22,45</sup>	0	288	11232	0	314	200	100	47
DPPG/Na <sup>+</sup>	Slipids / Åqvist <sup>22,45</sup>	0	288	11232	0	298	400	100	48
POPG	Slipids / Åqvist <sup>22,45</sup>	0	500	25000	0	310	500	100	49
POPG	Slipids / Åqvist <sup>22,45</sup>	0.11	500	25000	49	310	500	100	50
POPG	LIPID17 / Dang <sup>39,51</sup>	0	500	25000	0	310	500	100	52
POPG	LIPID17?	0.11	500	25000	49	310	500	100	53
POPG	GROMOS-CKP?	0	500	25000	0	310	500	100	54
POPG	GROMOS-CKP?	0.11	500	25000	49	310	500	100	55

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

17. Citations and ion model for CHARMM36?

18. Lipid17 simulation with correct dihedral potentials still coming.

19. Citation and ion model for GROMOS-CKP?

Table S3: List of MD simulations with PE and PG lipids mixed with PC.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	CaCl <sub>2</sub> (M)	<sup>a</sup> N <sub>l</sub>	<sup>b</sup> N <sub>w</sub>	<sup>c</sup> N <sub>c</sub>	<sup>d</sup> T (K)	<sup>e</sup> t <sub>sim</sub> (ns)	<sup>f</sup> t <sub>anal</sub> (ns)	<sup>g</sup> files
POPC	CHARMM36 <sup>?</sup>	0	0	500	25000	0	310	500	100	56
POPC:POPG (7:3)	CHARMM36 <sup>?</sup>	0	0	350	25000	0	310	500	100	57
POPC:POPG (1:1)	CHARMM36 <sup>?</sup>	0	0	150:150	31500	0	298	500	400	58
POPC:POPG (1:1)	CHARMM36 <sup>?</sup>	0	0.1	150:150	31329	57	298	400	300	59
POPC:POPG (1:1)	CHARMM36 <sup>?</sup>	0	1.08	150:150	29766	578	298	500	400	60
POPC:POPG (4:1)	CHARMM36 <sup>?</sup>	0	0	350:88	26280	0	298	500	400	61
POPC:POPG (4:1)	CHARMM36 <sup>?</sup>	0	0.1	350:88	26280	47	298	500	400	62
POPC:POPG (4:1)	CHARMM36 <sup>?</sup>	0	1.0	350:88	24927	451	298	500	400	63
POPC	CHARMM36 <sup>?</sup>	0	0	256	8704	0	300	300	250	64
POPC:POPE (1:1)	CHARMM36 <sup>?</sup>	0	0	128	8704	0	300	300	250	65
POPC	OPLS-MacRog <sup>32</sup>	0	0	128	5120	0	300	500	300	66
POPC:POPE (1:1)	OPLS-MacRog <sup>32</sup>	0	0	128	5120	0	300	500	300	67
POPC	Slipid <sup>18</sup>	0	0	512	23943	0	298	170	100	68
POPC:POPE (1:1)	Slipid <sup>18</sup>	0	0	128	5120	0	298	500	300	69
POPC	GROMOS-CKP / ?? ? ?	0	0	500	25000	0	310	500	100	70
POPC:POPG (7:3)	GROMOS-CKP / ?? ? ?	0	0	350:150	25000	0	310	500	100	71
POPC	Slipid <sup>18</sup>	0	0	500	25000	0	310	500	100	72
POPC:POPG (7:3)	Slipid / Åqvist <sup>18,22</sup>	0	0	350:150	25000	0	310	500	100	73
POPC:POPG (1:1)	Slipid / Dang <sup>18,51,74,75</sup>	0	0	128:128	12800	0	298	500	400	76
POPC:POPG (1:1)	Slipid / Dang <sup>18,51,74,75</sup>	0	0.1	128:128	12800	23	298	500	400	76
POPC:POPG (1:1)	Slipid / Dang <sup>18,51,74,75</sup>	0	0.2	128:128	12800	46	298	1500	500	76
POPC:POPG (1:1)	Slipid / Dang <sup>18,51,74,75</sup>	0	0.5	128:128	12800	115	298	1500	500	76
POPC:POPG (1:1)	Slipid / Dang <sup>18,51,74,75</sup>	0	1.0	128:128	12800	230	298	1500	500	76

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

20. Citation and ion model for GROMOS-CKP?

21. Citation and description for "Berger" model?

22. Lipid17 POPC and POPC:POPG mixtures (<https://doi.org/10.5281/zenodo.3241242> and <https://doi.org/10.5281/zenodo.3237656>) should be added after simulated with corrected dihedrals.

23. Upcoming Lipid17ecc with POPC:POPS (4:1) mixture simulations to be added.

Table S4: List of MD simulations with PE and PG lipids mixed with PC.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	CaCl <sub>2</sub> (M)	<sup>a</sup> N <sub>l</sub>	<sup>b</sup> N <sub>w</sub>	<sup>c</sup> N <sub>c</sub>	<sup>d</sup> T (K)	<sup>e</sup> t <sub>sim</sub> (ns)	<sup>f</sup> t <sub>anal</sub> (ns)	<sup>g</sup> files
POPC:POPG (4:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	0	350:88	26265	0	298	400	350	77
POPC:POPG (4:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	0.1	350:88	26124	47	298	400	250	78
POPC:POPG (4:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	1.0	350:88	24840	475	298	1200	200	79
POPC:POPG (1:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	0	150:150	31572	0	298	320	200	80
POPC:POPG (1:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	0.1	150:150	31401	57	298	718	198	81
POPC:POPG (1:1)	Lipid17 / Dang <sup>39,51,75</sup>	0	1.0	150:150	29865	569	298	720	200	82
POPC:POPG (1:1)	Lipid17ecc / ECC-ions <sup>83-85</sup>	0	0	150:150	31572	0	298	347.8	333	86
POPC:POPG (1:1)	Lipid17ecc / ECC-ions <sup>83-85</sup>	0	0.1	150:150	29865	54	298	400	300	87
POPC:POPG (1:1)	Lipid17ecc / ECC-ions <sup>83-85</sup>	0	1.0	150:150	29865	569	298	600	400	88
POPC	Berger <sup>?</sup> <b>24.</b>	0	0	256	10240	0	300	300	200	89
POPC:POPE (1:1)	Berger <sup>?</sup> <b>25.</b>	0	0	128	11008	0	300	300	200	90
POPC:DOPE (1:1)	Berger <sup>?</sup> <b>26.</b>	0	0	128	10240	0	300	300	200	91
DOPC	Berger <sup>?</sup> <b>27.</b>	0	0	256	11008	0	300	300	200	92
DOPC:DOPE (1:1)	Berger <sup>?</sup> <b>28.</b>	0	0	128	11008	0	300	300	200	93

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

**29. Citation and ion model for GROMOS-CKP?**

**30. Citation and description for "Berger" model?**

**31. Lipid17 POPC and POPC:POPG mixtures** (<https://doi.org/10.5281/zenodo.3241242> and <https://doi.org/10.5281/zenodo.3237656>) should be added after simulated with corrected dihedrals.

**32. Upcoming Lipid17ecc with POPC:POPS (4:1) mixture simulations to be added.**

*POPG with additional NaCl* <sup>45</sup>.Simulation details by A. Peon. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.

*DPPG* Data in 298 K is available at <sup>48</sup> and in 314 K at. <sup>47</sup> <sup>46</sup>.Simulation details by F. Favela. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.

*POPC:POPG mixture with additional NaCl* <sup>47</sup>.Simulation details by A. Peon. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.

## S9.4 Berger

*POPE* Data is available at. <sup>35,36</sup> <sup>48</sup>.Simulation details by T. Piggot.

*DOPE* Data is available at. <sup>37,38</sup> <sup>49</sup>.Simulation details by T. Piggot.

*POPC:POPE, POPC:DOPE and DOPC:DOPE mixtures* Data is available at. <sup>89,90</sup> 300 K with v-rescale (tau=0.1 ps), 1 bar with PR semiisotropic (tau=4 ps, compressibility=4.5e-5 bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion <sup>50</sup>.Simulation details by Fuchs et al.

## S9.5 GROMOS 43A1-S3

*POPE* Data is available at. <sup>29</sup> <sup>51</sup>.Simulation details by T. Piggot.

## S9.6 OPLS-UA

*POPE* Data is available at. <sup>31</sup> <sup>52</sup>.Simulation details by T. Piggot.

*POPE with vdW interaction in H* Data is available at. <sup>30</sup> <sup>53</sup>.Simulation details by T. Piggot.

## S9.7 GROMOS-CKP and GROMOS-CKPM

*POPE* Data is available at. <sup>25</sup> <sup>54</sup>.Simulation details by T. Piggot.

*DOPE* Data is available at. <sup>28</sup> <sup>55</sup>.Simulation details by T. Piggot.

*DPPE* Data is available at. <sup>24</sup> <sup>56</sup>.Simulation details by T. Piggot.

## S9.8 Lipid17

## S9.9 ECC-LIPID POPG

In ECC-lipid models, electronic continuum correction (ECC) is applied to implicitly include the missing electronic polarizability into the force field description.<sup>94,95</sup> In practise, this is implemented by scaling the charges and Lennard-Jones  $\sigma$ s of headgroup, glycerol backbone, and carbonyl regions of Amber Lipid14/17 models are scaled by constant factors. Here, we follow the approach that previously improved ion binding to bilayers containing negatively charged PS lipids:<sup>95</sup> ECC-POPC parameters (scaling factors  $f_q=0.8$  and  $f_\sigma=0.89$  applied to Lipid14 POPC parameters)<sup>94</sup> were used for POPC and scaling factors of  $f_q=0.75$  and  $f_\sigma=0.89$  were applied to the charges and Lennard-Jones  $\sigma$ s of headgroup, glycerol backbone, and carbonyl regions of Amber Lipid17 POPG parameters. The Lipid17 parameters (described above) and initial configurations were taken from Ref.<sup>52</sup> with the correct dihedral type, and the resulting parameters are available from Ref. ? . ECC-ion parameters with the scaled charges,<sup>83-85</sup> downloaded from [bitbucket.org/hseara/ions/src/master/](https://bitbucket.org/hseara/ions/src/master/), were used in these simulations.

## References

- (1) Botan, A.; Favela-Rosales, F.; Fuchs, P. F. J.; Javanainen, M.; Kanduč, M.; Kulig, W.; Lamberg, A.; Loison, C.; Lyubartsev, A.; Miettinen, M. S. et al. Toward Atomistic Resolution Structure of Phosphatidylcholine Headgroup and Glycerol Backbone at Different Ambient Conditions. *J. Phys. Chem. B* **2015**, *119*, 15075–15088.
- (2) Antila, H. S.; Buslaev, P.; Favela-Rosales, F.; Mendes Ferreira, T.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S. et al. Headgroup Structure and Cation Binding in Phosphatidylserine Lipid Bilayers. *The Journal of Physical Chemistry B* **2019**, acs.jpcc.9b06091.
- (3) Seelig, J.; Gally, H. U. Investigation of phosphatidylethanolamine bilayers by deuterium and phosphorus-31 nuclear magnetic resonance. *Biochemistry* **1976**, *15*, 5199–5204.
- (4) Gally, H. U.; Pluschke, G.; Overath, P.; Seelig, J. Structure of Escherichia coli membranes. Glycerol auxotrophs as a tool for the analysis of the phospholipid head-group region by deuterium magnetic resonance. *Biochemistry* **1981**, *20*, 1826–1831.
- (5) Borle, F.; Seelig, J. Ca<sup>2+</sup> binding to phosphatidylglycerol bilayers as studied by differential scanning calorimetry and 2H- and 31P-nuclear magnetic resonance. *Chemistry and Physics of Lipids* **1985**, *36*, 263 – 283.
- (6) Wohlgemuth, R.; Waespe-Sarcevic, N.; Seelig, J. Bilayers of phosphatidylglycerol. A deuterium and phosphorus nuclear magnetic resonance study of the head-group region. *Biochemistry* **1980**, *19*, 3315–3321.
- (7) Scherer, P.; Seelig, J. Structure and dynamics of the phosphatidylcholine and the phosphatidylethanolamine head group in L-M fibroblasts as studied by deuterium nuclear magnetic resonance. *EMBO J.* **1987**, *6*.

- (8) Macdonald, P. M.; Seelig, J. Calcium binding to mixed phosphatidylglycerol-phosphatidylcholine bilayers as studied by deuterium nuclear magnetic resonance. *Biochemistry* **1987**, *26*, 1231–1240.
- (9) Ollila, O. S.; Pabst, G. Atomistic resolution structure and dynamics of lipid bilayers in simulations and experiments. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2016**, *1858*, 2512 – 2528.
- (10) Seelig, J.; MacDonald, P. M.; Scherer, P. G. Phospholipid head groups as sensors of electric charge in membranes. *Biochemistry* **1987**, *26*, 7535–7541.
- (11) Antila, H. S.; Buslaev, P.; Favela-Rosales, F.; Mendes Ferreira, T.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S. et al. Headgroup Structure and Cation Binding in Phosphatidylserine Lipid Bilayers. *The Journal of Physical Chemistry B* **0**, *0*, null.
- (12) Catte, A.; Girych, M.; Javanainen, M.; Loison, C.; Melcr, J.; Miettinen, M. S.; Monticelli, L.; Maatta, J.; Oganessian, V. S.; Ollila, O. H. S. et al. Molecular electrometer and binding of cations to phospholipid bilayers. *Phys. Chem. Chem. Phys.* **2016**, *18*, 32560–32569.
- (13) Pan, J.; Heberle, F. A.; Tristram-Nagle, S.; Szymanski, M.; Koepfinger, M.; Katsaras, J.; KuÅDerka, N. Molecular structures of fluid phase phosphatidylglycerol bilayers as determined by small angle neutron and X-ray scattering. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2012**, *1818*, 2135 – 2148.
- (14) Javanainen, M. Simulation of a POPE bilayer at 310K with the CHARMM36 force field. 2019; <https://doi.org/10.5281/zenodo.2641987>.
- (15) PEON, CHARMM36 POPE Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3237461>.



- (16) PEÅŞN, A. CHARMM36 POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2577454>.
- (17) Piggot, T. CHARMM36-UA POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane and POPE is called PEUA). 2018; <https://doi.org/10.5281/zenodo.1293774>.
- (18) Jämbeck, J. P. M.; Lyubartsev, A. P. An Extension and Further Validation of an All-Atomistic Force Field for Biological Membranes. *J. Chem. Theory Comput.* **2012**, *8*, 2938–2948.
- (19) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPE bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.495247>.
- (20) Piggot, T. Slipids POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane). 2018; <https://doi.org/10.5281/zenodo.1293813>.
- (21) Peon, A. SLIPID POPE Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3231342>.
- (22) Åqvist, J. Ion-water interaction potentials derived from free energy perturbation simulations. *J. Phys. Chem.* **1990**, *94*, 8021–8024.
- (23) PEÅŞN, A. SLIPID POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2578069>.
- (24) Piggot, T. GROMOS-CKP DPPE Simulations (versions 1 and 2) 342 K. 2018; <https://doi.org/10.5281/zenodo.1293957>.
- (25) Piggot, T. GROMOS-CKP POPE Simulations (versions 1 and 2) 313 K. 2018; <https://doi.org/10.5281/zenodo.1293932>.
- (26) PEON, A. GROMOS POPE Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3237754>.

- (27) PEÅŠN, A. Gromos POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2574491>.
- (28) Piggot, T. GROMOS-CKP DOPE Simulations (versions 1 and 2) 271 K. 2018; <https://doi.org/10.5281/zenodo.1293941>.
- (29) Piggot, T. GROMOS 43A1-S3 POPE Simulations (versions 1 and 2) 313 K (NOTE: anisotropic pressure coupling). 2018; <https://doi.org/10.5281/zenodo.1293762>.
- (30) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K with vdW on H atoms. 2018; <https://doi.org/10.5281/zenodo.1293853>.
- (31) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K. 2018; <https://doi.org/10.5281/zenodo.1293855>.
- (32) RÅšg, T.; OrÅĆowski, A.; Llorente, A.; Skotland, T.; SylvÅĎnne, T.; Kauhanen, D.; Ekroos, K.; Sandvig, K.; Vattulainen, I. Data including GROMACS input files for atomistic molecular dynamics simulations of mixed, asymmetric bilayers including molecular topologies, equilibrated structures, and force field for lipids compatible with OPLS-AA parameters. *Data in Brief* **2016**, 7, 1171 – 1174.
- (33) Javanainen, M. Simulation of a POPE bilayer, lipid model based on OPLS-aa by Rog et al. 2019; <https://doi.org/10.5281/zenodo.3571071>.
- (34) Milan Rodriguez, P.; Fuchs, P. F. MacRog pure POPE MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3725670>.
- (35) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293889>.
- (36) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293891>.

- (37) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293928>.
- (38) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293905>.
- (39) Gould, I.; Skjevik, A.; Dickson, C.; Madej, B.; Walker, R. Lipid17: A Comprehensive AMBER Force Field for the Simulation of Zwitterionic and Anionic Lipids. 2018; In preparation.
- (40) PEON, A. LIPID17 POPE Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3378970>.
- (41) PEÅŞN, A. LIPID17 POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2577305>.
- (42) Ollila, O. H. S. POPG lipid bilayer simulation at T298K ran with MODEL\_CHARMM\_GUI force field and Gromacs. 2017; <https://doi.org/10.5281/zenodo.1011096>.
- (43) PEÅŞN, A. CHARMM36 POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2573531>.
- (44) ANTONIO, CHARMM36 POPG Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3237463>.
- (45) Jämbeck, J. P. M.; Lyubartsev, A. P. Implicit inclusion of atomic polarization in modeling of partitioning between water and lipid bilayers. *Phys. Chem. Chem. Phys.* **2013**, *15*, 4677–4686.
- (46) Favela-Rosales, F. MD simulation trajectory of a fully hydrated POPG bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546133>.

- (47) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @314K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546136>.
- (48) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @298K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546135>.
- (49) PEĂȘN, A. SLIPID POPG Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3364460>.
- (50) PEĂȘN, A. SLIPID POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2633773>.
- (51) Smith, D. E.; Dang, L. X. Computer simulations of NaCl association in polarizable water. *J. Chem. Phys* **1994**, *100*, 3757–3766.
- (52) PEON, A. LIPID17 POPG Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3247659>.
- (53) PEĂȘN, A. LIPID17 POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.2573905>.
- (54) PEON, A. GROMOS POPG Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3266166>.
- (55) PEĂȘN, A. Gromos POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3257649>.
- (56) PEON, A. CHARMM36 POPC Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3247813>.
- (57) PEON, A. CHARMM36 POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3248689>.

- (58) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in water and Na<sup>+</sup> counter ions. 2020; <https://doi.org/10.5281/zenodo.3997116>.
- (59) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in 0.1 M CaCl<sub>2</sub> solution and Na<sup>+</sup> counter ions. 2020; <https://doi.org/10.5281/zenodo.4005515>.
- (60) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in 1 M CaCl<sub>2</sub> solution and Na<sup>+</sup> counter ions. 2020; <https://doi.org/10.5281/zenodo.3997135>.
- (61) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in water with Na<sup>+</sup> counter ions. 2020; <https://doi.org/10.5281/zenodo.3996952>.
- (62) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in 0.1 M CaCl<sub>2</sub> solution with Na<sup>+</sup> counter ions. 2020; <https://doi.org/10.5281/zenodo.3997019>.
- (63) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in 1 M CaCl<sub>2</sub> solution with Na<sup>+</sup> counterions. 2020; <https://doi.org/10.5281/zenodo.3997037>.
- (64) Papadopoulos, C.; Fuchs, P. F. CHARMM36 pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306800>.
- (65) Papadopoulos, C.; Fuchs, P. F. CHARMM36 POPC/POPE (50%-50%) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306821>.
- (66) Milan Rodriguez, P.; Fuchs, P. F. MacRog pure POPC MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3741793>.
- (67) Milan Rodriguez, P.; Fuchs, P. F. MacRog POPC/POPE 1:1 MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3725637>.

- (68) Favela-Rosales, F. MD simulation trajectory of a lipid bilayer: Pure POPC in water. SLIPIDS, Gromacs 4.6.3. 2016. 2016; <https://doi.org/10.5281/zenodo.166034>.
- (69) Javanainen, M. Simulation of POPC:POPE 1:1 membrane with the Slipids force field. 2020; <https://doi.org/10.5281/zenodo.3605386>.
- (70) PEON, A. GROMOS-CKP POPC Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3247435>.
- (71) PEON, A. GROMOS-CKP POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3266240>.
- (72) PEON, A. SLIPID POPC Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3235552>.
- (73) PeON, A. SLIPID POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K ). 2019; <https://doi.org/10.5281/zenodo.3240156>.
- (74) Jämbeck, J. P.; Lyubartsev, A. P. Another piece of the membrane puzzle: extending slipids further. *Journal of chemical theory and computation* **2012**, *9*, 774–784.
- (75) Dang, L. X.; Schenter, G. K.; Glezakou, V.-A.; Fulton, J. L. Molecular simulation analysis and X-ray absorption measurement of Ca<sup>2+</sup>, K<sup>+</sup> and Cl<sup>-</sup> ions in solution. *J. Phys. Chem. B* **2006**, *110*, 23644–54.
- (76) Javanainen, M. Simulations of POPC:POPG 1:1 membranes with varying levels of CaCl<sub>2</sub> using the Slipids force field. 2020; <https://doi.org/10.5281/zenodo.3613573>.
- (77) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na<sup>+</sup> counterions, 298K. 2019; <https://doi.org/10.5281/zenodo.3693681>.
- (78) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na<sup>+</sup> counterions and 100mM CaCl<sub>2</sub>, 298K. 2019; <https://doi.org/10.5281/zenodo.3833725>.

- (79) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na<sup>+</sup> counterions and 1000mM CaCl<sub>2</sub>, 298K. 2019; <https://doi.org/10.5281/zenodo.3874378>.
- (80) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions, 298K. 2019; <https://doi.org/10.5281/zenodo.3857816>.
- (81) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions and 100mM CaCl<sub>2</sub>, 298K. 2019; <https://doi.org/10.5281/zenodo.3871590>.
- (82) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions and 1000mM CaCl<sub>2</sub>, 298K. 2019; <https://doi.org/10.5281/zenodo.3864993>.
- (83) Pluhařová, E.; Fischer, H. E.; Mason, P. E.; Jungwirth, P. Hydration of the chloride ion in concentrated aqueous solutions using neutron scattering and molecular dynamics. *Mol. Phys.* **2014**, *112*, 1230–1240.
- (84) Kohagen, M.; Mason, P. E.; Jungwirth, P. Accounting for Electronic Polarization Effects in Aqueous Sodium Chloride via Molecular Dynamics Aided by Neutron Scattering. *J. Phys. Chem. B* **2016**, *120*, 1454–1460.
- (85) Martínek, T.; Duboué-Dijon, E.; Timr, Š.; Mason, P. E.; Baxová, K.; Fischer, H. E.; Schmidt, B.; Pluhařová, E.; Jungwirth, P. Calcium ions in aqueous solutions: Accurate force field description aided by ab initio molecular dynamics and neutron scattering. *J. Chem. Phys.* **2018**, *148*, 222813.
- (86) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions, 298K. 2020; <https://doi.org/10.5281/zenodo.3859339>.
- (87) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions and 100mM CaCl<sub>2</sub>, 298K. 2020; <https://doi.org/10.5281/zenodo.3855729>.

- (88) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na<sup>+</sup> counterions and 1000mM CaCl<sub>2</sub>, 298K. 2020; <https://doi.org/10.5281/zenodo.3862036>.
- (89) AmÄllie, B.; F.J., F. P. Berger pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402417>.
- (90) AmÄllie, B.; F.J., F. P. Berger POPC/POPE (50:50 ratio) MD simulation (300 K - 400ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402449>.
- (91) AmÄllie, B.; F.J., F. P. Berger POPC/DOPE (50:50 ratio) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402441>.
- (92) AmÄllie, B.; F.J., F. P. Berger pure DOPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402411>.
- (93) AmÄllie, B.; F.J., F. P. Berger DOPC/DOPE (50:50 ratio) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402437>.
- (94) Melcr, J.; Martinez-Seara, H.; Nencini, R.; Kolafa, J.; Jungwirth, P.; Ollila, O. H. S. Accurate Binding of Sodium and Calcium to a POPC Bilayer by Effective Inclusion of Electronic Polarization. *The Journal of Physical Chemistry B* **2018**, *122*, 4546–4557.
- (95) Melcr, J.; Ferreira, T.; Jungwirth, P.; Ollila, O. H. S. Improved Cation Binding to Lipid Bilayer with Negatively Charged POPS by Effective Inclusion of Electronic Polarization. [https://github.com/ohs011lila/ecc\\_lipids/blob/master/Manuscript/manuscript.pdf](https://github.com/ohs011lila/ecc_lipids/blob/master/Manuscript/manuscript.pdf), Submitted.