



# Supporting Information:

## NMRlipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PE and PG lipids

Pavel Buslaev,<sup>†</sup> Fernando Favela-Rosales,<sup>‡</sup> Patrick Fuchs,<sup>¶</sup> Matti Javanainen,<sup>§</sup>  
Jesper J. Madsen,<sup>||,⊥</sup> Josef Melcr,<sup>§, #</sup> Markus S. Miettinen,<sup>@</sup> O. H. Samuli  
Ollila,<sup>\*,△</sup> Chris G. Papadopoulos,<sup>∇</sup> Antonio Peón,<sup>††</sup> Thomas J. Piggot,<sup>‡‡</sup> and  
Pierre Poulain<sup>¶</sup>

<sup>†</sup>*University of Jyväskylä*

<sup>‡</sup>*Departamento de Investigación, Tecnológico Nacional de México, Campus Zacatecas  
Occidente, México*

<sup>¶</sup>*Paris, France*

<sup>§</sup>*Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences,  
Flemingovo nám. 542/2, CZ-16610 Prague 6, Czech Republic*

<sup>||</sup>*Department of Chemistry, The University of Chicago, Chicago, Illinois, United States of  
America*

<sup>⊥</sup>*Department of Global Health, College of Public Health, University of South Florida, Tampa,  
Florida, United States of America*

<sup>#</sup>*Groningen Biomolecular Sciences and Biotechnology Institute and The Zernike Institute  
for Advanced Materials, University of Groningen, 9747 AG Groningen, The Netherlands*

<sup>@</sup>*Department of Theory and Bio-Systems, Max Planck Institute of Colloids and Interfaces,  
14424 Potsdam, Germany*

<sup>△</sup>*Institute of Biotechnology, University of Helsinki*

<sup>∇</sup>*I2BC - University Paris Sud  
S2*

<sup>††</sup>*Spain*

<sup>‡‡</sup>*Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, United*

## S1 Simulated systems

### S1.1 CHARMM36

*POPE* [Simulation details by M. Javanainen.](#)

*POPE with additional NaCl* [Simulation details by A. Peon.](#)

*POPG* [Simulation details by Ollila.](#)

*POPG with additional NaCl* [Simulation details by A. Peon.](#)

*POPC:POPE mixtures* Data is available at.<sup>1,2</sup> 300 K with v-rescale ( $\tau=0.1$  ps), 1 bar with PR semiisotropic ( $\tau=4$  ps, compressibility= $4.5\text{e-}5$  bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [Full simulation details by Fuchs et al.](#)

*POPC:POPG mixture with additional calcium* [Simulation details by J. Madsen.](#)

*POPC:POPG mixture with additional NaCl* [Simulation details by A. Peon.](#)

### S1.2 CHARMM36ua

*POPE* Data is available at.<sup>3</sup> [Simulation details by T. Piggot.](#)

### S1.3 Slipids

*POPE* Data is available at.<sup>4</sup> [Simulation details by T. Piggot.](#)

*POPE with additional NaCl* [Simulation details by A. Peon.](#)

*DPPE* Data is available at.<sup>5</sup> [Simulation details by F. Favela.](#)

*POPG* Data is available at.<sup>6</sup> [Simulation details by F. Favela.](#)

*POPG with additional NaCl* [Simulation details by A. Peon.](#)

*DPPG* Data in 298 K is available at<sup>7</sup> and in 314 K at.<sup>8</sup> [Simulation details by F. Favela.](#)

*POPC:POPG mixture with additional NaCl* [Simulation details by A. Peon.](#)

## S1.4 Berger

*POPE* Data is available at.<sup>9,10</sup> [Simulation details by T. Piggot.](#)

*DOPE* Data is available at.<sup>11,12</sup> [Simulation details by T. Piggot.](#)

*POPC:POPE*, *POPC:DOPE* and *DOPC:DOPE* mixtures Data is available at.<sup>13,14</sup> 300 K with v-rescale ( $\tau=0.1$  ps), 1 bar with PR semiisotropic ( $\tau=4$  ps, compressibility= $4.5\text{e-}5$  bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [Simulation details by Fuchs et al.](#)

## S1.5 GROMOS 43A1-S3

*POPE* Data is available at.<sup>15</sup> [Simulation details by T. Piggot.](#)

## S1.6 OPLS-UA

*POPE* Data is available at.<sup>16</sup> [Simulation details by T. Piggot.](#)

*POPE with vdW interaction in H* Data is available at.<sup>17</sup> [Simulation details by T. Piggot.](#)

## S1.7 GROMOS-CKP and GROMOS-CKPM

*POPE* Data is available at.<sup>18</sup> [Simulation details by T. Piggot.](#)

*DOPE* Data is available at.<sup>19</sup> [Simulation details by T. Piggot.](#)

*DPPE* Data is available at.<sup>20</sup> [Simulation details by T. Piggot.](#)

## S2 R-PDLF and SDROSS experiments




Figure S1: (A) Chemical structure of POPE with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPE sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar sliced from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.

A, B etc. labels to be put in the figure.




Figure S2: (A) Chemical structure of POPG with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPG sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar sliced from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.




Figure S3: Simpson simlaton of S-DROSS curve of  $\beta$ -carbon of POPG.

# S3 Changes of PG headgroup order parameters upon addition of PC




Figure S4: Modulation of PG lipid headgroup order parameters with the increasing amount of PC in lipid bilayer from experiments<sup>21,22</sup> and simulations with different force fields.



## S4 Sodium binding to POPC simulations




Figure S5: Sodium (solid line) and chloride ion density profiles along membrane normal from different simulations with PC lipids.

Discussion about differences to the NMRlipids II to be discussed once we have the details on ions models.

## References

- (1) Papadopoulos, C.; Fuchs, P. F. CHARMM36 pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306800>.
- (2) Papadopoulos, C.; Fuchs, P. F. CHARMM36 POPC/POPE (50%-50%) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306821>.
- (3) Piggot, T. CHARMM36-UA POPE Simulations (versions 1 and 2) 310 K (NOTE:

- hexagonal membrane and POPE is called PEUA). 2018; <https://doi.org/10.5281/zenodo.1293774>.
- (4) Piggot, T. Slipids POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane). 2018; <https://doi.org/10.5281/zenodo.1293813>.
  - (5) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPE bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.495247>.
  - (6) Favela-Rosales, F. MD simulation trajectory of a fully hydrated POPG bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546133>.
  - (7) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @298K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546135>.
  - (8) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @314K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546136>.
  - (9) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293889>.
  - (10) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293891>.
  - (11) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293928>.
  - (12) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293905>.
  - (13) Amélie, B.; F.J., F. P. Berger pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402417>.

- (14) Amélie, B.; F.J., F. P. Berger POPC/POPE (50:50 ratio) MD simulation (300 K - 400ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402449>.
- (15) Piggot, T. GROMOS 43A1-S3 POPE Simulations (versions 1 and 2) 313 K (NOTE: anisotropic pressure coupling). 2018; <https://doi.org/10.5281/zenodo.1293762>.
- (16) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K. 2018; <https://doi.org/10.5281/zenodo.1293855>.
- (17) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K with vdW on H atoms. 2018; <https://doi.org/10.5281/zenodo.1293853>.
- (18) Piggot, T. GROMOS-CKP POPE Simulations (versions 1 and 2) 313 K. 2018; <https://doi.org/10.5281/zenodo.1293932>.
- (19) Piggot, T. GROMOS-CKP DOPE Simulations (versions 1 and 2) 271 K. 2018; <https://doi.org/10.5281/zenodo.1293941>.
- (20) Piggot, T. GROMOS-CKP DPPE Simulations (versions 1 and 2) 342 K. 2018; <https://doi.org/10.5281/zenodo.1293957>.
- (21) Borle, F.; Seelig, J. Ca<sup>2+</sup> binding to phosphatidylglycerol bilayers as studied by differential scanning calorimetry and <sup>2</sup>H- and <sup>31</sup>P-nuclear magnetic resonance. *Chemistry and Physics of Lipids* **1985**, *36*, 263 – 283.
- (22) Macdonald, P. M.; Seelig, J. Calcium binding to mixed phosphatidylglycerol-phosphatidylcholine bilayers as studied by deuterium nuclear magnetic resonance. *Biochemistry* **1987**, *26*, 1231–1240.