

**Supporting Information:**

**NMRlipids IV: Headgroup & glycerol backbone  
structures, and cation binding in bilayers with PE  
and PG lipids**

O. H. Samuli Ollila<sup>\*,†</sup>

*<sup>†</sup>Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech  
Republic, Prague 6, Czech Republic*

*<sup>‡</sup>Institute of Biotechnology, University of Helsinki*

E-mail: samuli.ollila@helsinki.fi

## S1 Simulated systems

### S1.1 CHARMM36

*POPC:POPE mixtures* Data is available at.<sup>1,2</sup> 300 K with v-rescale ( $\tau=0.1$  ps), 1 bar with PR semiisotropic ( $\tau=4$  ps, compressibility= $4.5\text{e-}5$  bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [Full simulation details by Fuchs et al.](#)

*POPC:POPG mixture with additional calcium* [Simulation details by J. Madsen.](#)

### S1.2 CHARMM36ua

*POPE* Data is available at.<sup>3</sup> [Simulation details by T. Piggot.](#)

### S1.3 Slipids

*POPE* Data is available at.<sup>4</sup> [Simulation details by T. Piggot.](#)

*DPPE* Data is available at.<sup>5</sup> [Simulation details by F. Favela.](#)

*POPG* Data is available at.<sup>6</sup> [Simulation details by F. Favela.](#)

*DPPG* Data in 298 K is available at<sup>7</sup> and in 314 K at.<sup>8</sup> [Simulation details by F. Favela.](#)

### S1.4 Berger

*POPE* Data is available at.<sup>9,10</sup> [Simulation details by T. Piggot.](#)

*DOPE* Data is available at.<sup>11,12</sup> [Simulation details by T. Piggot.](#)

*POPC:POPE, POPC:DOPE and DOPC:DOPE mixtures* Data is available at.<sup>13,14</sup> 300 K with v-rescale ( $\tau=0.1$  ps), 1 bar with PR semiisotropic ( $\tau=4$  ps, compressibility= $4.5\text{e-}5$  bar<sup>-1</sup>), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [Simulation details by Fuchs et al.](#)

## S1.5 GROMOS 43A1-S3

*POPE* Data is available at.<sup>15</sup> [Simulation details by T. Piggot.](#)

## S1.6 OPLS-UA

*POPE* Data is available at.<sup>16</sup> [Simulation details by T. Piggot.](#)

*POPE with vdW interaction in H* Data is available at.<sup>17</sup> [Simulation details by T. Piggot.](#)

## S1.7 GROMOS-CKP and GROMOS-CKPM

*POPE* Data is available at.<sup>18</sup> [Simulation details by T. Piggot.](#)

*DOPE* Data is available at.<sup>19</sup> [Simulation details by T. Piggot.](#)

*DPPE* Data is available at.<sup>20</sup> [Simulation details by T. Piggot.](#)

## S2 R-PDLF and SDROSS experiments




Figure S1: (A) Chemical structure of POPE with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPE sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar sliced from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.

A, B etc. labels to be put in the figure.

## References

- (1) Papadopoulos, C.; Fuchs, P. F. CHARMM36 pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306800>.
- (2) Papadopoulos, C.; Fuchs, P. F. CHARMM36 POPC/POPE (50%-50%) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306821>.
- (3) Piggot, T. CHARMM36-UA POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane and POPE is called PEUA). 2018; <https://doi.org/10.5281/zenodo.1293774>.
- (4) Piggot, T. Slipids POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane). 2018; <https://doi.org/10.5281/zenodo.1293813>.
- (5) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPE bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.495247>.
- (6) Favela-Rosales, F. MD simulation trajectory of a fully hydrated POPG bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546133>.
- (7) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @298K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546135>.
- (8) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @314K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546136>.
- (9) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293889>.
- (10) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293891>.

- (11) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293928>.
- (12) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293905>.
- (13) Amélie, B.; F.J., F. P. Berger pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402417>.
- (14) Amélie, B.; F.J., F. P. Berger POPC/POPE (50:50 ratio) MD simulation (300 K - 400ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402449>.
- (15) Piggot, T. GROMOS 43A1-S3 POPE Simulations (versions 1 and 2) 313 K (NOTE: anisotropic pressure coupling). 2018; <https://doi.org/10.5281/zenodo.1293762>.
- (16) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K. 2018; <https://doi.org/10.5281/zenodo.1293855>.
- (17) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K with vdW on H atoms. 2018; <https://doi.org/10.5281/zenodo.1293853>.
- (18) Piggot, T. GROMOS-CKP POPE Simulations (versions 1 and 2) 313 K. 2018; <https://doi.org/10.5281/zenodo.1293932>.
- (19) Piggot, T. GROMOS-CKP DOPE Simulations (versions 1 and 2) 271 K. 2018; <https://doi.org/10.5281/zenodo.1293941>.
- (20) Piggot, T. GROMOS-CKP DPPE Simulations (versions 1 and 2) 342 K. 2018; <https://doi.org/10.5281/zenodo.1293957>.