

Supporting Information:

NMRLipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PE and PG lipids

Pavel Buslaev,[†] Rebeca García Fandiño,[‡] Fernando Favela-Rosales,[§] Tiago Ferreira,^{||} Patrick Fuchs,[⊥] Ivan Gushchin,[#] Matti Javanainen,[@] Anne M. Kiirikki,[△] Jesper J. Madsen,^{∇,††} Josef Melcr,^{@,‡‡} Paula Milan Rodriguez,[⊥] Markus S. Miettinen,^{¶¶} O. H. Samuli Ollila,^{*,△} Chris G. Papadopoulos,^{§§} Antonio Peón,^{|||} Thomas J. Piggot,^{⊥⊥} and Ángel Piñeiro^{##}

[†]*University of Jyväskylä*

[‡]*Center for Research in Biological Chemistry and Molecular Materials (CiQUS),
Universidade de Santiago de Compostela, E-15782 Santiago de Compostela, Spain*

^{¶¶}*CIQUP, Centro de Investigaçãõ em Quãmica, Departamento de Quãmica e
Bioquãmica, Faculdade de Ciãncias, Universidade do Porto, Porto, Portugal*

[§]*Departamento de Ciencias Básicas, Tecnológico Nacional de México, Campus Zacatecas
Occidente, México*

^{||}*Halle, Germany*

[⊥]*Paris, France*

[#]*Moscow Institute of Physics and Technology, Dolgoprudny, Russia*

[@]*Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences,
Flemingovo nám. 542/2, CZ-16610 Prague 6, Czech Republic*

[△]*Institute of Biotechnology, University of Helsinki*

[∇]*Department of Chemistry, The University of Chicago, Chicago, Illinois, United States of
America*

^{††}*Department of Global Health, College of Public Health, University of South Florida,
Tampa, Florida, United States of America*

S1 R-PDLF and SDROSS experiments



Figure S1: (A) Chemical structure of POPE with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPE sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar sliced from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.

1.A, B etc. labels to be put in the figure.



Figure S2: (A) Chemical structure of POPG with the labeling of headgroup and glycerol backbone carbons. (B) INEPT spectra from POPG sample with the headgroup and glycerol backbone peaks labeled. (C) 2D R-PDLF spectra (D) Dipolar sliced from the 2D R-PDLF spectra with the resulting order parameters on top of figures. (E) Experimental S-DROSS curves giving signs of the order parameters.




Figure S3: Simpson simlaton of S-DROSS curve of β -carbon of POPG.

S2 Lipid ligand names in PDB used in the analysis of conformations of protein-bound lipids

PC: PLC, PX4, 6PL, LIO, HGX, PC7, PC8, P1O, 6O8, XP5, EGY, PLD, SBM, HXG, and PCW

PE: 8PE, PTY, 3PE, PEH, PEF, 6OE, 6O9, 9PE, PEV, 46E, SBJ, L9Q, PEK, EPH, ZPE, 9TL, 9Y0, 6OU, LOP, and PEE

PG: PGT, PGK, LHG, 44G, PGV, OZ2, D3D, PGW, DR9, P6L, PG8, H3T, and GOT

PS: PSF, PS6, Q3G, P5S, D39, PS2, 17F, and 8SP.

S3 Evaluation of simulations against NMR experiments

S3.1 Conformational ensembles of headgroup and glycerol backbone in PE and PG lipids

The quality of PE and PG headgroup conformational ensembles in different simulations against NMR experiments is evaluated in figures S4 and S5 using C-H bond order parameters as in our previous studies for PC and PS lipids.^{1,2} Conclusions are the same for all lipids: None of the force fields correctly captures the lipid headgroup conformational ensembles, but CHARMM36 gives results closest to experiments.

It should be noted that the PG headgroup is biologically abundant R enantiomer in all simulations, while our ^{13}C NMR experiments has a racemic mixture. Nevertheless, previous ^2H NMR experiments comparing results between different enantiomers concluded that the structural differences between these are minor.³




Figure S4: The headgroup and glycerol backbone order parameters of PE lipids from experiments (POPE and signs this work, DPPE from Ref. 4 and E.coliPE from Ref. 5) and simulations with different force fields.

2.This should be clarified as in NMRLipidsI and error bars should be added. Probably larger error bars for united atom models based on the report by Fuchs et al.

3.Lipid17 data should be updated to the one with correct dihedrals reported here

<https://github.com/NMRLipids/NMRLipidsIVPEandPG/issues/12#issuecomment-756641407>




Figure S5: The headgroup and glycerol backbone order parameters of PG lipids from experiments (POPG and signs from this work and from Ref. 6, DPPG with 100mM NaCl from Ref. 3, and E. Coli PG results from Ref. 5) and simulations with different force fields.

4. Lipid17 data should be updated to the one with correct dihedrals reported here

<https://github.com/NMRLipids/NMRLipidsIVPEandPG/issues/12#issuecomment-756641407>

S3.2 PC headgroup in mixtures with PE or PG lipids

Headgroup order parameters of PC lipids are unchanged upon addition of zwitterionic lipids or cholesterol in experiments, but increase upon addition of negatively charged PG or PS lipids because headgroup dipole tilts more parallel to the membrane plane after incorporation of negative charges into the membrane.^{7,10,11} The response of PC headgroup order parameters to the addition of PE or PG lipids from different simulations is compared with experiments in figure S6. None of the simulations reproduce neither the experimentally observed increase in PC headgroup order parameters with increasing amount of PG nor the related tilting of the headgroup more parallel with the membrane. Similar observations in our previous work for PS lipids were explained by the overestimated counterion binding affinity that neutralizes the effect of added negative charge.² All simulations except Berger-OPLS predict tilting of P-N headgroup outwards from the membrane and decrease of PC headgroup order parameters upon addition of PE lipids. These results are not in line with experiments where the PC headgroup order parameters are not affected by zwitterionic lipids.⁷ The good performance of Berger-OPLS simulations in here is surprising because headgroup conformational ensemble is not very close to experiments in this model and the response of headgroup order parameters to cholesterol was significantly overestimated by the Berger/Höltje force field in our previous work.¹

In conclusion, more accurate force fields are needed to correctly simulate the interactions between different headgroups.



Figure S6: Modulation of POPC headgroup order parameters with increasing amount of POPE (left) and POPG (right) in bilayer from experiments at 298 K^{7,8} and simulations with different force fields (temperatures listed in tables S3 and S4 are between 298-310 K). Signs are determined as discussed in Refs. 1,9.

S3.3 PG headgroup in mixtures with PC lipids

Changes in other than PC lipid headgroup with changing membrane composition are less extensively characterized in the literature. The β -carbon order parameter in PG headgroup increases mildly⁸ or is unchanged⁶ upon increasing amount of PC lipids (Fig. S7), but experimental data from α -carbon is not available. Also the tested force fields predict very small changes for the β -carbon order parameter, while the P-N vector tilt and its response to the increased amount of PC varies significantly between force fields in figure S7. Therefore, more experimental data and more accurate force fields are still required to resolve the PG conformational ensembles in mixtures with other lipids.



Figure S7: Modulation of PG lipid headgroup order parameters with the increasing amount of PC in lipid bilayer from experiments at 298 K^{6,8} and simulations with different force fields at 310 K.

S3.4 Calcium binding to POPC:POPG mixtures

The changes of headgroup order parameters in POPC:POPG mixtures upon addition of CaCl_2 between different simulations and experiments^{6,8} are compared in figures S8 (molar ratio 1:1) and S10 (molar ratio 4:1). The results are in line with our previous studies: most force fields overestimate the calcium binding,^{2,12} but CHARMM36 with the NBfix correction underestimates the binding affinity,² and the implicit inclusion of electronic polarizability using the electronic continuum correction (ECC) improves the results.^{13,14}




Figure S8: Modulation of headgroup order parameters of POPC (*left*) and POPG (*right*) in POPC:POPG (1:1) mixture upon addition of CaCl_2 in 298 K temperature from experiments^{6,8} and simulations. The β -carbon order parameter of POPC (dashed line on top left) is not directly measured but calculated from empirical relation $\Delta S_\beta = 0.43\Delta S_\alpha$.¹⁵ The changes with respect to the systems without CaCl_2 are shown for other data than for the α -carbon of POPG for which experimental order parameter is not available. Calcium density distributions are shown in figure S9.





Figure S9: Calcium ion density profiles along membrane normal from simulations of POPC:POPG (1:1) mixtures with different force fields. The changes in the order parameters upon addition of CaCl_2 are compared with experiments in figure S8.






Figure S11: Calcium ion density profiles along membrane normal from simulations of POPC:POPG (4:1) mixtures with different force fields.

S4 Dihedral angle distributions and the analysis of relative energies




Figure S12: Heavy atom dihedral angle distributions from CHARMM36 simulations that correctly capture the order parameter differences between the force fields.

S5 Changes in headgroup conformations upon addition of charged surfactants or CaCl_2




Figure S13: Changes in PC headgroup conformational ensembles upon increasing the amount of positive charge in bilayer, characterized by the heavy atom dihedral distributions, from CHARMM36 simulations.



Figure S14: Changes in POPC lipid17ecc dihedrals with increasing amount of CaCl_2 .



Figure S15: Changes in POPC CHARMM36 dihedrals with increasing amount of CaCl_2 .




Figure S16: Changes in POPG Slipids dihedrals with increasing amount of CaCl_2 .




Figure S17: Changes in POPG lipid17 dihedrals with increasing amount of CaCl_2 .

S6 Simulated systems

The simulated systems of pure PE and PG bilayers without additional ions are listed in Tables S1 and S2, and lipid mixtures with additional ions in Tables S3 and S4. the simulation data are indexed in a searchable database available at www.nmrlipids.fi, and in the NMRLipids/MATCH repository (github.com/NMRLipids/MATCH). The large set of MD simulation data was analysed using the development version of NMRLipids databank. Unique naming convention for lipid atoms in each force field was defined using the mapping files and analysis for all simulations indexed in NMRLipids databank manner were performed using Python codes.

The C–H bond order parameters were calculated directly from the carbon and hydrogen positions using the definition

$$S_{\text{CH}} = \frac{1}{2} \langle 3 \cos^2 \theta - 1 \rangle, \quad (1)$$

where θ is the angle between the C–H bond and the membrane normal (taken to align with z , with bilayer periodicity in the xy -plane). Angular brackets denote average over all sampled configurations. The order parameters were first calculated averaging over time separately for each lipid in the system. The average and the standard error of the mean were then calculated over different lipids. Python programs that use the MDAnalysis library^{16,17} used for all atom simulations are available in Ref. 18 (`scripts/calcOrderParameters.py`). For united atom simulations, the trajectories with hydrogens having ideal geometry were constructed first using either `buildH` program¹⁹ or (`scratch/opAAUA_prod.py`) in Ref. 18, and the order parameters were then calculated from these trajectories. This approach has been tested against trajectories with explicit hydrogens and the deviations in order parameters are small.^{19,20}

The number density profiles of ions were calculated using the `gmx density` tool of the GROMACS software package.²¹

Table S1: List of MD simulations with PE lipids.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
POPE	CHARMM36 [?]	0	144	5760	0	310	500	400	22
POPE	CHARMM36 [?]	0	500	25000	0	310	500	100	23
POPE	CHARMM36 [?]	0.11	500	25000	50	310	500	100	24
POPE	CHARMM36ua [?]	0	336	15254	0	310	2×200	2×100	25
DPPE	Slipids ²⁶	0	288	9386	0	336	200	100	27
POPE	Slipids ²⁶	0	336	?	0	310	2×200	2×100	28
POPE	Slipids ²⁶	0	500	25000	0	310	500	100	29
POPE	Slipids / Åqvist ^{26,30}	0.11	500	25000	50	310	500	100	31
DPPE	GROMOS-CKP [?]	0	128	3655	0	342	2×500	2×400	32
POPE	GROMOS-CKP [?]	0	128	3552	0	313	2×500	2×400	33
POPE	GROMOS-CKP [?]	0	500	25000	0	310	500	100	34
POPE	GROMOS-CKP [?]	0.11	500	25000	50	310	500	100	35
DOPE	GROMOS-CKP [?]	0	128	4789	0	271	2×500	2×400	36
POPE	GROMOS 43A1-S3 [?]	0	128	3552	0	313	2×200	2×100	37
POPE	OPLS-UA vdW on H [?]	0	128	3328	0	303	2×200	2×100	38
POPE	OPLS-UA [?]	0	128	3328	0	303	2×200	2×100	39
POPE	OPLS-MacRog ⁴⁰	0	144	5760	0	310	500	350	41
POPE	OPLS-MacRog ⁴⁰	0	128	5120	0	300	500	300	42
POPE	Berger-Vries [?]	0	128	3552	0	303	2×200	2×100	43
POPE	Berger-largeH [?]	0	128	3552	0	303	2×200	2×100	44
DOPE	Berger-Vries [?]	0	128	4789	0	271	2×200	2×100	45
DOPE	Berger-largeH [?]	0	128	4789	0	271	2×300	2×100	46
POPE	LIPID17 ⁴⁷	0	500	25000	50	310	500	100	48
POPE	LIPID17 ⁴⁷	0.11	500	25000	50	310	500	100	49

^aNumber of lipid molecules with largest mole fraction

^bNumber of water molecules

^cNumber of additional cations

^dSimulation temperature

^eTotal simulation time

^fTime used for analysis

^gReference for simulation files

5. Simulations with added NaCl are not currently used here, maybe should be removed from the table?

6. Citation for CHARMM36 PE?

7. Which ion model is used in ²⁴?

8. Citation for GROMOS-CKP?

9. Citation for GROMOS 43A1-S3?

10. Citation for OPLS-UA models?

11. Citations for Berger-* simulations?

12. LIPID17 simulations with correct dihedrals still coming

Table S2: List of MD simulations with PG lipids.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
POPG/K ⁺	CHARMM36 [?] 13.	0	118	4110	0	298	100	100	50
POPG	CHARMM36 [?]	0.11	500	25000	49	310	500	100	51
POPG	CHARMM36 [?]	0	500	25000	0	310	500	100	52
POPG/Na ⁺	Slipids / Åqvist ^{30,53}	0	288	10664	0	298	250	100	54
DPPG/Na ⁺	Slipids / Åqvist ^{30,53}	0	288	11232	0	314	200	100	55
DPPG/Na ⁺	Slipids / Åqvist ^{30,53}	0	288	11232	0	298	400	100	56
POPG	Slipids / Åqvist ^{30,53}	0	500	25000	0	310	500	100	57
POPG	Slipids / Åqvist ^{30,53}	0.11	500	25000	49	310	500	100	58
POPG	LIPID17 / Dang ^{47,59}	0	500	25000	0	310	500	100	60
POPG	LIPID17 [?]	0.11	500	25000	49	310	500	100	61
POPG	GROMOS-CKP [?]	0	500	25000	0	310	500	100	62
POPG	GROMOS-CKP [?]	0.11	500	25000	49	310	500	100	63

^aNumber of lipid molecules with largest mole fraction

^bNumber of water molecules

^cNumber of additional cations

^dSimulation temperature

^eTotal simulation time

^fTime used for analysis

^gReference for simulation files

14.Simulations with added NaCl are not currently used here, maybe should be removed from the table?

15.Citations and ion model for CHARMM36?

16.Lipid17 simulation with ions with correct dihedral potentials still coming?

17.Citation and ion model for GROMOS-CKP?

Table S3: List of MD simulations with PE and PG lipids mixed with PC.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	CaCl ₂ (M)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
POPC	CHARMM36 [?]	0	0	500	25000	0	310	500	100	64
POPC:POPG (7:3)	CHARMM36 [?]	0	0	350	25000	0	310	500	100	65
POPC:POPG (1:1)	CHARMM36 [?]	0	0	150:150	31500	0	298	500	400	66
POPC:POPG (1:1)	CHARMM36 [?]	0	0.1	150:150	31329	57	298	400	300	67
POPC:POPG (1:1)	CHARMM36 [?]	0	1.08	150:150	29766	578	298	500	400	68
POPC:POPG (4:1)	CHARMM36 [?]	0	0	350:88	26280	0	298	500	400	69
POPC:POPG (4:1)	CHARMM36 [?]	0	0.1	350:88	26280	47	298	500	400	70
POPC:POPG (4:1)	CHARMM36 [?]	0	1.0	350:88	24927	451	298	500	400	71
POPC	CHARMM36 [?]	0	0	256	8704	0	300	300	250	72
POPC:POPE (1:1)	CHARMM36 [?]	0	0	128	8704	0	300	300	250	73
POPC	OPLS-MacRog ⁴⁰	0	0	128	5120	0	300	500	300	74
POPC:POPE (1:1)	OPLS-MacRog ⁴⁰	0	0	128	5120	0	300	500	300	75
POPC	Slipid ²⁶	0	0	512	23943	0	298	170	100	76
POPC:POPE (1:1)	Slipid ²⁶	0	0	128	5120	0	298	500	300	77
POPC	GROMOS-CKP / ?? [?] ?	0	0	500	25000	0	310	500	100	78
POPC:POPG (7:3)	GROMOS-CKP / ?? [?] ?	0	0	350:150	25000	0	310	500	100	79
POPC	Slipid ²⁶	0	0	500	25000	0	310	500	100	80
POPC:POPG (7:3)	Slipid / Åqvist ^{26,30}	0	0	350:150	25000	0	310	500	100	81
POPC:POPG (1:1)	Slipid / Dang ^{26,59,82,83}	0	0	128:128	12800	0	298	500	400	84
POPC:POPG (1:1)	Slipid / Dang ^{26,59,82,83}	0	0.1	128:128	12800	23	298	500	400	84
POPC:POPG (1:1)	Slipid / Dang ^{26,59,82,83}	0	0.2	128:128	12800	46	298	1500	500	84
POPC:POPG (1:1)	Slipid / Dang ^{26,59,82,83}	0	0.5	128:128	12800	115	298	1500	500	84
POPC:POPG (1:1)	Slipid / Dang ^{26,59,82,83}	0	1.0	128:128	12800	230	298	1500	500	84

^aNumber of lipid molecules with largest mole fraction

^bNumber of water molecules

^cNumber of additional cations

^dSimulation temperature

^eTotal simulation time

^fTime used for analysis

^gReference for simulation files

18. Citation and ion model for GROMOS-CKP?

Table S4: List of MD simulations with PE and PG lipids mixed with PC.

lipid/counter-ions	force field for lipids / ions	NaCl (M)	CaCl ₂ (M)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
POPC:POPG (4:1)	Lipid17 / Dang ^{47,59,83}	0	0	350:88	26265	0	298	400	350	85
POPC:POPG (4:1)	Lipid17 / Dang ^{47,59,83}	0	0.1	350:88	26124	47	298	400	250	86
POPC:POPG (4:1)	Lipid17 / Dang ^{47,59,83}	0	1.0	350:88	24840	475	298	1200	200	87
POPC:POPG (1:1)	Lipid17 / Dang ^{47,59,83}	0	0	150:150	31572	0	298	320	200	88
POPC:POPG (1:1)	Lipid17 / Dang ^{47,59,83}	0	0.1	150:150	31401	57	298	718	198	89
POPC:POPG (1:1)	Lipid17 / Dang ^{47,59,83}	0	1.0	150:150	29865	569	298	720	200	90
POPC:POPG (4:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	0	350:88	26265	0	298	400	300	94
POPC:POPG (4:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	0.1	350:88	26124	47	298	400	300	95
POPC:POPG (4:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	1.0	350:88	24840	475	298	400	300	96
POPC:POPG (1:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	0	150:150	31572	0	298	347.8	333	97
POPC:POPG (1:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	0.1	150:150	29865	54	298	400	300	98
POPC:POPG (1:1)	Lipid17ecc / ECC-ions ⁹¹⁻⁹³	0	1.0	150:150	29865	569	298	600	400	99
POPC	Berger [?] 19.	0	0	256	10240	0	300	300	200	100
POPC:POPE (1:1)	Berger [?] 20.	0	0	128	11008	0	300	300	200	101
POPC:DOPE (1:1)	Berger [?] 21.	0	0	128	10240	0	300	300	200	102
DOPC	Berger [?] 22.	0	0	256	11008	0	300	300	200	103
DOPC:DOPE (1:1)	Berger [?] 23.	0	0	128	11008	0	300	300	200	104

^aNumber of lipid molecules with largest mole fraction

^bNumber of water molecules

^cNumber of additional cations

^dSimulation temperature

^eTotal simulation time

^fTime used for analysis

^gReference for simulation files

24. Citation and description for "Berger" model?

S6.1 CHARMM36

POPE **25.Simulation details by M. Javanainen.**

POPE with additional NaCl **26.Simulation details by A. Peon.**

POPG **27.Simulation details by Ollila.**

POPG with additional NaCl **28.Simulation details by A. Peon.**

POPC:POPE mixtures Data is available at.^{72,73} 300 K with v-rescale ($\tau=0.1$ ps), 1 bar with PR semiisotropic ($\tau=4$ ps, compressibility= $4.5\text{e-}5$ bar⁻¹), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion **29.Full simulation details by Fuchs et al.**

POPC:POPG mixture with additional calcium **30.Simulation details by A. Kiirikki.**

POPC and POPC:POPG (7:3) mixture **31.Simulation details by A. Peon.**

S6.2 CHARMM36ua

POPE Data is available at.²⁵ **32.Simulation details by T. Piggot.**

S6.3 Slipids

POPE Data is available at.²⁸ **33.Simulation details by T. Piggot.**

POPE with additional NaCl **34.Simulation details by A. Peon. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.**

DPPE with 288 lipids. The starting structure for simulation was constructed with the MEMBRANE BUILDER website:¹⁰⁵ 288 DPPE lipids and 9386 water molecules. The TIP3P¹⁰⁶ water model was used to solvate the system. Simulation was performed for 200 ns, and the last 100 ns were used for the analysis. Simulation was carried out within the NPT ensemble using the GROMACS 5.0.4 package.¹⁰⁷ Timestep of 2 fs was used with the leapfrog integrator. The Nosé–Hoover thermostat^{108,109} was used with reference temperature of 336 K and a relaxation time constant of 0.5 ps; lipids and water were coupled separately to the heat bath. Pressure was kept constant at 1.013 bar using a semi-isotropic Parrinello–Rahman

barostat¹¹⁰ with a time constant of 10.0 ps. Long-range electrostatic interactions were calculated using the PME method.^{111,112} A real space cut-off of 1.0 nm was employed with grid spacing of 0.12 nm in the reciprocal space. Lennard-Jones potentials were cut off at 1.4 nm, with a dispersion correction applied to both energy and pressure. All covalent bonds in lipids were constrained using the LINCS algorithm,¹¹³ whereas water molecules were constrained using SETTLE.¹¹⁴ Twin-range cutoffs, 1.0 nm and 1.6 nm, were used for the neighbor lists with the long-range neighbor list updated every 10 steps.

POPG with 288 lipids. The starting structure for simulation was constructed with the MEMBRANE BUILDER website:¹⁰⁵ 288 POPG lipids, 10664 water molecules and 288 Na ions. The TIP3P¹⁰⁶ water model was used to solvate the system and Ions are described by the parameters derived by Aqvist.³⁰ Simulation was performed for 250 ns, and the last 100 ns were used for the analysis. Same simulation conditions as DPPE with reference temperature of 298 K.

POPG with additional NaCl **35.Simulation details by A. Peon. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.**

DPPG with 288 lipids. The starting structure for simulation was constructed with the MEMBRANE BUILDER website:¹⁰⁵ 288 DPPG lipids, 11232 water molecules and 288 Na ions. The TIP3P¹⁰⁶ water model was used to solvate the system and Ions are described by the parameters derived by Aqvist.³⁰ For the 298 K temperature, simulation was performed for 400 ns, and the last 100 ns were used for the analysis. For the 314 K temperature, simulation was performed for 200 ns, and the last 100 ns were used for the analysis. Same simulation conditions as DPPE for both temperatures.

POPC:POPG mixture with additional NaCl **36.Simulation details by A. Peon. I have assumed that ion parameters are default Slipids, i.e., Åqvist, please correct if this is not true.**

POPC:POPG mixture with additional CaCl **37.Simulation details by M. Javanainen.**

S6.4 Berger

POPE Data is available at.^{43,44} [38.Simulation details by T. Piggot.](#)

DOPE Data is available at.^{45,46} [39.Simulation details by T. Piggot.](#)

POPC:POPE, *POPC:DOPE* and *DOPC:DOPE* mixtures Data is available at.^{100,101} 300 K with v-rescale ($\tau=0.1$ ps), 1 bar with PR semiisotropic ($\tau=4$ ps, compressibility= $4.5\text{e-}5$ bar⁻¹), PME order 4 and space 0.12, rcoulomb and rvdw 1.0, 128 lipids per leaflet, no ion [40.Simulation details by Fuchs et al.](#)

S6.5 GROMOS 43A1-S3

POPE Data is available at.³⁷ [41.Simulation details by T. Piggot.](#)

S6.6 OPLS-UA

POPE Data is available at.³⁹ [42.Simulation details by T. Piggot.](#)

POPE with vdW interaction in H Data is available at.³⁸ [43.Simulation details by T. Piggot.](#)

S6.7 GROMOS-CKP and GROMOS-CKPM

POPE Data is available at.³³ [44.Simulation details by T. Piggot.](#)

DOPE Data is available at.³⁶ [45.Simulation details by T. Piggot.](#)

DPPE Data is available at.³² [46.Simulation details by T. Piggot.](#)

POPG [47.Simulation details by A. Peon.](#)

POPC:POPG mixture [48.Simulation details by A. Peon.](#)

S6.8 OPLS-MacRog

POPE [49.Simulation details by M. Javanainen and P. Fuchs.](#)

POPC:POPE mixtures [50.Simulation details by P. Fuchs.](#)

S6.9 Lipid17

POPE 51.Simulation details by A. Peon.

POPG 52.Simulation details by A. Peon.

POPC:POPG 53.Simulation details by S. Virtanen or O. H. S. Ollila.

S6.10 Lipid17ecc

54.This is to be finished and POPC:POPG mixtures to be described In ECC-lipid models, electronic continuum correction (ECC) is applied to implicitly include the missing electronic polarizability into the force field description.^{13?} In practise, this is implemented by scaling the charges and Lennard-Jones σ s of headgroup, glycerol backbone, and carbonyl regions of Amber Lipid14/17 models are scaled by constant factors. Here, we follow the approach that previously improved ion binding to bilayers containing negatively charged PS lipids:[?] ECC-POPC parameters (scaling factors $f_q=0.8$ and $f_\sigma=0.89$ applied to Lipid14 POPC parameters)¹³ were used for POPC and scaling factors of $f_q=0.75$ and $f_\sigma=0.89$ were applied to the charges and Lennard-Jones σ s of headgroup, glycerol backbone, and carbonyl regions of Amber Lipid17 POPG parameters. The Lipid17 parameters (described above) and initial configurations were taken from Ref.⁶⁰ with the correct dihedral type, and the resulting parameters are available from Ref. ? . ECC-ion parameters with the scaled charges,⁹¹⁻⁹³ downloaded from bitbucket.org/hseara/ions/src/master/, were used in these simulations.

References

- (1) Botan, A.; Favela-Rosales, F.; Fuchs, P. F. J.; Javanainen, M.; Kanduč, M.; Kulig, W.; Lamberg, A.; Loison, C.; Lyubartsev, A.; Miettinen, M. S. et al. Toward Atomistic Resolution Structure of Phosphatidylcholine Headgroup and Glycerol Backbone at Different Ambient Conditions. *J. Phys. Chem. B* **2015**, *119*, 15075–15088.
- (2) Antila, H. S.; Buslaev, P.; Favela-Rosales, F.; Mendes Ferreira, T.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S. et al. Headgroup Structure and Cation Binding in Phosphatidylserine Lipid Bilayers. *The Journal of Physical Chemistry B* **2019**, acs.jpcc.9b06091.
- (3) Wohlgemuth, R.; Waespe-Sarcevic, N.; Seelig, J. Bilayers of phosphatidylglycerol. A deuterium and phosphorus nuclear magnetic resonance study of the head-group region. *Biochemistry* **1980**, *19*, 3315–3321.
- (4) Seelig, J.; Gally, H. U. Investigation of phosphatidylethanolamine bilayers by deuterium and phosphorus-31 nuclear magnetic resonance. *Biochemistry* **1976**, *15*, 5199–5204.
- (5) Gally, H. U.; Pluschke, G.; Overath, P.; Seelig, J. Structure of Escherichia coli membranes. Glycerol auxotrophs as a tool for the analysis of the phospholipid head-group region by deuterium magnetic resonance. *Biochemistry* **1981**, *20*, 1826–1831.
- (6) Borle, F.; Seelig, J. Ca²⁺ binding to phosphatidylglycerol bilayers as studied by differential scanning calorimetry and ²H- and ³¹P-nuclear magnetic resonance. *Chemistry and Physics of Lipids* **1985**, *36*, 263 – 283.
- (7) Scherer, P.; Seelig, J. Structure and dynamics of the phosphatidylcholine and the phosphatidylethanolamine head group in L-M fibroblasts as studied by deuterium nuclear magnetic resonance. *EMBO J.* **1987**, *6*.

- (8) Macdonald, P. M.; Seelig, J. Calcium binding to mixed phosphatidylglycerol-phosphatidylcholine bilayers as studied by deuterium nuclear magnetic resonance. *Biochemistry* **1987**, *26*, 1231–1240.
- (9) Ollila, O. S.; Pabst, G. Atomistic resolution structure and dynamics of lipid bilayers in simulations and experiments. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2016**, *1858*, 2512 – 2528.
- (10) Seelig, J.; MacDonald, P. M.; Scherer, P. G. Phospholipid head groups as sensors of electric charge in membranes. *Biochemistry* **1987**, *26*, 7535–7541.
- (11) Antila, H. S.; Buslaev, P.; Favela-Rosales, F.; Mendes Ferreira, T.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S. et al. Headgroup Structure and Cation Binding in Phosphatidylserine Lipid Bilayers. *The Journal of Physical Chemistry B* **0**, *0*, null.
- (12) Catte, A.; Girych, M.; Javanainen, M.; Loison, C.; Melcr, J.; Miettinen, M. S.; Monticelli, L.; Maatta, J.; Oganessian, V. S.; Ollila, O. H. S. et al. Molecular electrometer and binding of cations to phospholipid bilayers. *Phys. Chem. Chem. Phys.* **2016**, *18*, 32560–32569.
- (13) Melcr, J.; Martinez-Seara, H.; Nencini, R.; Kolafa, J.; Jungwirth, P.; Ollila, O. H. S. Accurate Binding of Sodium and Calcium to a POPC Bilayer by Effective Inclusion of Electronic Polarization. *The Journal of Physical Chemistry B* **2018**, *122*, 4546–4557.
- (14) Melcr, J.; Ferreira, T. M.; Jungwirth, P.; Ollila, O. H. S. Improved Cation Binding to Lipid Bilayers with Negatively Charged POPS by Effective Inclusion of Electronic Polarization. *Journal of Chemical Theory and Computation* **2020**, *16*, 738–748.
- (15) Akutsu, H.; Seelig, J. Interaction of metal ions with phosphatidylcholine bilayer membranes. *Biochemistry* **1981**, *20*, 7366–7373.

- (16) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. *Journal of Computational Chemistry* **2011**, *32*, 2319–2327.
- (17) Richard J. Gowers;; Max Linke;; Jonathan Barnoud;; Tyler J. E. Reddy;; Manuel N. Melo;; Sean L. Seyler;; Jan DomaÅŹski;; David L. Dotson;; SÅľbastien Buchoux;; Ian M. Kenney, et al. MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. Proceedings of the 15th Python in Science Conference. 2016; pp 98 – 105.
- (18) ohsOllila;; et al., MATCH GitHub repository. <https://github.com/NMRLipids/MATCH>.
- (19) Fuchs, P.; et al., BuildH GitHub repository. <https://github.com/patrickfuchs/buildH>.
- (20) Piggot, T. J.; Allison, J. R.; Sessions, R. B.; Essex, J. W. On the Calculation of Acyl Chain Order Parameters from Lipid Simulations. *J. Chem. Theory Comput.* **2017**, *13*, 5683–5696.
- (21) Abraham, M.; van der Spoel, D.; Lindahl, E.; Hess, B.; the GROMACS development team, GROMACS user manual version 5.0.7. 2015.
- (22) Javanainen, M. Simulation of a POPE bilayer at 310K with the CHARMM36 force field. 2019; <https://doi.org/10.5281/zenodo.2641987>.
- (23) PEON, CHARMM36 POPE Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3237461>.
- (24) PEÅŹN, A. CHARMM36 POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2577454>.

- (25) Piggot, T. CHARMM36-UA POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane and POPE is called PEUA). 2018; <https://doi.org/10.5281/zenodo.1293774>.
- (26) Jämbeck, J. P. M.; Lyubartsev, A. P. An Extension and Further Validation of an All-Atomistic Force Field for Biological Membranes. *J. Chem. Theory Comput.* **2012**, *8*, 2938–2948.
- (27) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPE bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.495247>.
- (28) Piggot, T. Slipids POPE Simulations (versions 1 and 2) 310 K (NOTE: hexagonal membrane). 2018; <https://doi.org/10.5281/zenodo.1293813>.
- (29) Peon, A. SLIPID POPE Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3231342>.
- (30) Åqvist, J. Ion-water interaction potentials derived from free energy perturbation simulations. *J. Phys. Chem.* **1990**, *94*, 8021–8024.
- (31) PEÅŞN, A. SLIPID POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2578069>.
- (32) Piggot, T. GROMOS-CKP DPPE Simulations (versions 1 and 2) 342 K. 2018; <https://doi.org/10.5281/zenodo.1293957>.
- (33) Piggot, T. GROMOS-CKP POPE Simulations (versions 1 and 2) 313 K. 2018; <https://doi.org/10.5281/zenodo.1293932>.
- (34) PEON, A. GROMOS POPE Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3237754>.
- (35) PEÅŞN, A. Gromos POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2574491>.

- (36) Piggot, T. GROMOS-CKP DOPE Simulations (versions 1 and 2) 271 K. 2018; <https://doi.org/10.5281/zenodo.1293941>.
- (37) Piggot, T. GROMOS 43A1-S3 POPE Simulations (versions 1 and 2) 313 K (NOTE: anisotropic pressure coupling). 2018; <https://doi.org/10.5281/zenodo.1293762>.
- (38) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K with vdW on H atoms. 2018; <https://doi.org/10.5281/zenodo.1293853>.
- (39) Piggot, T. OPLS-UA POPE Simulations (versions 1 and 2) 303 K. 2018; <https://doi.org/10.5281/zenodo.1293855>.
- (40) RÅsg, T.; OrÅĆowski, A.; Llorente, A.; Skotland, T.; SylvÅdnne, T.; Kauhanen, D.; Ekroos, K.; Sandvig, K.; Vattulainen, I. Data including GROMACS input files for atomistic molecular dynamics simulations of mixed, asymmetric bilayers including molecular topologies, equilibrated structures, and force field for lipids compatible with OPLS-AA parameters. *Data in Brief* **2016**, *7*, 1171 – 1174.
- (41) Javanainen, M. Simulation of a POPE bilayer, lipid model based on OPLS-aa by Rog et al. 2019; <https://doi.org/10.5281/zenodo.3571071>.
- (42) Milan Rodriguez, P.; Fuchs, P. F. MacRog pure POPE MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3725670>.
- (43) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293889>.
- (44) Piggot, T. Berger POPE Simulations (versions 1 and 2) 303 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293891>.
- (45) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - de Vries repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293928>.

- (46) Piggot, T. Berger DOPE Simulations (versions 1 and 2) 271 K - larger repulsive H. 2018; <https://doi.org/10.5281/zenodo.1293905>.
- (47) Gould, I.; Skjevik, A.; Dickson, C.; Madej, B.; Walker, R. Lipid17: A Comprehensive AMBER Force Field for the Simulation of Zwitterionic and Anionic Lipids. 2018; In preparation.
- (48) PEON, A. LIPID17 POPE Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3378970>.
- (49) PEÅŞN, A. LIPID17 POPE Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2577305>.
- (50) Ollila, O. H. S. POPG lipid bilayer simulation at T298K ran with MODEL_CHARMM_GUI force field and Gromacs. 2017; <https://doi.org/10.5281/zenodo.1011096>.
- (51) PEÅŞN, A. CHARMM36 POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2573531>.
- (52) ANTONIO, CHARMM36 POPG Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3237463>.
- (53) Jämbeck, J. P. M.; Lyubartsev, A. P. Implicit inclusion of atomic polarization in modeling of partitioning between water and lipid bilayers. *Phys. Chem. Chem. Phys.* **2013**, *15*, 4677–4686.
- (54) Favela-Rosales, F. MD simulation trajectory of a fully hydrated POPG bilayer: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546133>.
- (55) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @314K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546136>.

- (56) Favela-Rosales, F. MD simulation trajectory of a fully hydrated DPPG bilayer @298K: SLIPIDS, Gromacs 5.0.4. 2017. 2017; <https://doi.org/10.5281/zenodo.546135>.
- (57) PeÅŝn, A. LIPID17 POPG Bilayer Simulation (Last 100 ns, 310 K) using dihedral type 9. 2019; <https://doi.org/10.5281/zenodo.3832274>.
- (58) PEÅŦN, A. SLIPID POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2633773>.
- (59) Smith, D. E.; Dang, L. X. Computer simulations of NaCl association in polarizable water. *J. Chem. Phys* **1994**, *100*, 3757–3766.
- (60) PEON, A. LIPID17 POPG Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3247659>.
- (61) PEÅŦN, A. LIPID17 POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.2573905>.
- (62) PEON, A. GROMOS POPG Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3266166>.
- (63) PEÅŦN, A. Gromos POPG Bilayer Simulation (Last 100 ns, 150 mM NaCl, 310 K). 2019; <https://doi.org/10.5281/zenodo.3257649>.
- (64) PEON, A. CHARMM36 POPC Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3247813>.
- (65) PEON, A. CHARMM36 POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3248689>.
- (66) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in water and Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3997116>.

- (67) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in 0.1 M CaCL solution and Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.4005515>.
- (68) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 1:1 MD simulation with CHARMM36 in 1 M CaCL solution and Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3997135>.
- (69) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in water with Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3996952>.
- (70) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in 0.1 M CaCL₂ solution with Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3997019>.
- (71) Kiirikki, A. M.; Ollila, O. H. S. POPC:POPG 4:1 MD simulation with CHARMM36 in 1 M CaCL₂ solution with Na⁺ counterions. 2020; <https://doi.org/10.5281/zenodo.3997037>.
- (72) Papadopoulos, C.; Fuchs, P. F. CHARMM36 pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306800>.
- (73) Papadopoulos, C.; Fuchs, P. F. CHARMM36 POPC/POPE (50%-50%) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1306821>.
- (74) Milan Rodriguez, P.; Fuchs, P. F. MacRog pure POPC MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3741793>.
- (75) Milan Rodriguez, P.; Fuchs, P. F. MacRog POPC/POPE 1:1 MD simulation (300 K - 500ns - 1 bar). 2020; <https://doi.org/10.5281/zenodo.3725637>.
- (76) Favela-Rosales, F. MD simulation trajectory of a lipid bilayer: Pure POPC in water. SLIPIDS, Gromacs 4.6.3. 2016. 2016; <https://doi.org/10.5281/zenodo.166034>.

- (77) Javanainen, M. Simulation of POPC:POPE 1:1 membrane with the Slipids force field. 2020; <https://doi.org/10.5281/zenodo.3605386>.
- (78) PEON, A. GROMOS-CKP POPC Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3247435>.
- (79) PEON, A. GROMOS-CKP POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3266240>.
- (80) PEON, A. SLIPID POPC Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3235552>.
- (81) PeON, A. SLIPID POPC-POPG 7:3 Bilayer Simulation (Last 100 ns, 310 K). 2019; <https://doi.org/10.5281/zenodo.3240156>.
- (82) Jämbeck, J. P.; Lyubartsev, A. P. Another piece of the membrane puzzle: extending slipids further. *Journal of chemical theory and computation* **2012**, *9*, 774–784.
- (83) Dang, L. X.; Schenter, G. K.; Glezakou, V.-A.; Fulton, J. L. Molecular simulation analysis and X-ray absorption measurement of Ca²⁺, K⁺ and Cl⁻ ions in solution. *J. Phys. Chem. B* **2006**, *110*, 23644–54.
- (84) Javanainen, M. Simulations of POPC:POPG 1:1 membranes with varying levels of CaCl₂ using the Slipids force field. 2020; <https://doi.org/10.5281/zenodo.3613573>.
- (85) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na⁺ counterions, 298K. 2019; <https://doi.org/10.5281/zenodo.3693681>.
- (86) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na⁺ counterions and 100mM CaCl₂, 298K. 2019; <https://doi.org/10.5281/zenodo.3833725>.

- (87) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 80:20 MD simulation, Na⁺ counterions and 1000mM CaCl₂, 298K. 2019; <https://doi.org/10.5281/zenodo.3874378>.
- (88) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions, 298K. 2019; <https://doi.org/10.5281/zenodo.3857816>.
- (89) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions and 100mM CaCl₂, 298K. 2019; <https://doi.org/10.5281/zenodo.3871590>.
- (90) Virtanen, S.; Ollila, O. H. S. LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions and 1000mM CaCl₂, 298K. 2019; <https://doi.org/10.5281/zenodo.3864993>.
- (91) Pluhařová, E.; Fischer, H. E.; Mason, P. E.; Jungwirth, P. Hydration of the chloride ion in concentrated aqueous solutions using neutron scattering and molecular dynamics. *Mol. Phys.* **2014**, *112*, 1230–1240.
- (92) Kohagen, M.; Mason, P. E.; Jungwirth, P. Accounting for Electronic Polarization Effects in Aqueous Sodium Chloride via Molecular Dynamics Aided by Neutron Scattering. *J. Phys. Chem. B* **2016**, *120*, 1454–1460.
- (93) Martínek, T.; Duboué-Dijon, E.; Timr, Š.; Mason, P. E.; Baxová, K.; Fischer, H. E.; Schmidt, B.; Pluhařová, E.; Jungwirth, P. Calcium ions in aqueous solutions: Accurate force field description aided by ab initio molecular dynamics and neutron scattering. *J. Chem. Phys.* **2018**, *148*, 222813.
- (94) Kiirikki, A. M.; Ollila, O. H. S. Lipid17ecc POPC:POPG 4:1 MD simulation in water with Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3997154>.
- (95) Kiirikki, A. M.; Ollila, O. H. S. Lipid17ecc POPC:POPG 4:1 bilayer simulation in 0.1

- M CaCl₂ solution and Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.3997176>.
- (96) Kiirikki, A. M.; Ollila, O. H. S. Lipid17ecc POPC:POPG 4:1 bilayer simulation in 1 M CaCl₂ solution and Na⁺ counter ions. 2020; <https://doi.org/10.5281/zenodo.4492639>.
- (97) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions, 298K. 2020; <https://doi.org/10.5281/zenodo.3859339>.
- (98) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions and 100mM CaCl₂, 298K. 2020; <https://doi.org/10.5281/zenodo.3855729>.
- (99) Ollila, O. H. S.; Virtanen, I. S. ECC-LIPID17 POPC-POPG 50:50 MD simulation, Na⁺ counterions and 1000mM CaCl₂, 298K. 2020; <https://doi.org/10.5281/zenodo.3862036>.
- (100) AmÄllie, B.; F.J., F. P. Berger pure POPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402417>.
- (101) AmÄllie, B.; F.J., F. P. Berger POPC/POPE (50:50 ratio) MD simulation (300 K - 400ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402449>.
- (102) AmÄllie, B.; F.J., F. P. Berger POPC/DOPE (50:50 ratio) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402441>.
- (103) AmÄllie, B.; F.J., F. P. Berger pure DOPC MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402411>.
- (104) AmÄllie, B.; F.J., F. P. Berger DOPC/DOPE (50:50 ratio) MD simulation (300 K - 300ns - 1 bar). 2018; <https://doi.org/10.5281/zenodo.1402437>.

- (105) Ghahremanpour, M. M.; Arab, S. S.; Aghazadeh, S. B.; Zhang, J.; van der Spoel, D. MemBuilder: a web-based graphical interface to build heterogeneously mixed membrane bilayers for the GROMACS biomolecular simulation program. *Bioinformatics* **2013**, *30*, 439–441.
- (106) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **1983**, *79*, 926–935.
- (107) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2015**, *1*, 19–25.
- (108) Nose, S. A molecular dynamics method for simulations in the canonical ensemble. *Mol. Phys.* **1984**, *52*, 255–268.
- (109) Hoover, W. G. Canonical dynamics: Equilibrium phase-space distributions. *Phys. Rev. A* **1985**, *31*, 1695–1697.
- (110) Parrinello, M.; Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. *J. Appl. Phys.* **1981**, *52*, 7182–7190.
- (111) Darden, T.; York, D.; Pedersen, L. Particle mesh Ewald: An $N \cdot \log(N)$ method for Ewald sums in large systems. *J. Chem. Phys.* **1993**, *98*, 10089–10092.
- (112) Essman, U. L.; Perera, M. L.; Berkowitz, M. L.; Larden, T.; Lee, H.; Pedersen, L. G. A smooth particle mesh ewald potential. *J. Chem. Phys.* **1995**, *103*, 8577–8592.
- (113) Hess, B.; Bekker, H.; Berendsen, H. J. C.; Fraaije, J. G. E. M. LINCS: a linear constraint solver for molecular dynamics simulations. *J. Comput. Chem.* **1997**, *18*, 1463–1472.
- (114) Miyamoto, S.; Kollman, P. A. SETTLE: An analytical Version of the SHAKE and RATTLE Algorithm for Rigid Water Models. *J. Comput. Chem* **1992**, *13*, 952–962.