

# NMRLipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PS lipids

O. H. Samuli Ollila<sup>1,2,\*</sup> and NMRLipids collaboration <sup>1</sup>Authorship query to be sent soon.<sup>3</sup>

<sup>1</sup>*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic*

<sup>2</sup>*Institute of Biotechnology, University of Helsinki*

<sup>3</sup>*nmrlipids.blogspot.fi*

(Dated: June 21, 2018)

Primarily measured but also simulated NMR order parameters will be collected also for other than phosphatidylcholine (these are discussed in NMRLipids I) headgroup. The information will be used to understand structural differences between different lipid molecules in bilayers.

## INTRODUCTION

Phosphatidylserine (PS) is the most common negatively charged lipid in eukaryotic membranes. PS lipids compose 8.5% of total lipid weight of erythrocytes, but the abundance varies between different organelles up to 25-35% in plasma membrane [1–3]. Despite of the relatively low abundance, PS lipids are important signaling molecules. They interact with signaling proteins [2], regulate surface charge and protein localization [4], and induce protein aggregation [5, 6]. Some domains specifically interact PS lipids, while others are attracted by general electrostatics and the binding can be regulated by calcium [2]. Therefore, the structural details of lipid headgroups and the details of cation binding are crucial for the PS mediated signaling processes.

Previous experimental studies have concluded that PS headgroups have more rigid conformation than phosphocholines (PC) due to hydrogen bonding or electrostatic interactions [7, 8]. Multivalent cations and  $\text{Li}^+$  are able to form strong dehydrated molecular complexes with PS lipids, while monovalent ions interact more weakly with PS containing bilayers [9–16]. Dilution of bilayers with PC lipids makes PS headgroups less rigid and reduces propensity for the formation of strong complexes with multivalent ions [7, 8, 15, 16].

The structure of PS lipid headgroups and their interactions with ions have been studied with various experimental methods and theoretical techniques [15, 17? ]. However, the consensus has not been reached due to the difficulties to interpret the experimental data [? ] and the inaccuracies in simulation models at the headgroup region [18–20]. Some studies propose that the negatively charged lipids attract cations only due to the increase of local concentration in the vicinity of membranes and that the binding constant of cations is similar to zwitterionic and negatively charged lipids [21? , 22]. On the other hand, some studies propose specific binding of calcium directly to PS lipid headgroups [23? , 24]. The NMR data proposes that the PS headgroup is more rigid than PC, PE or PG headgroups, but more detailed interpretation has not been done.

Headgroup and glycerol backbone C-H bond order parameters calculated from MD simulations have been recently used to interpret the lipid structures in NMR experiments and to validate lipid structure and ion binding in simulations of PC

lipid bilayers [18–20, 25]. In this work we apply this approach to PS lipid headgroup in order to elucidate the structural details and ion binding to negatively charged lipids. The results are expected to elucidate also PS mediated signalling events because glycerol backbone and headgroup structure and behaviour are similar in model membranes and in bacteria [21, 26, 27].

## METHODS

### Solid state NMR experiments

The magnitude and signs of the C-H bond order parameters in headgroup and glycerol backbone were measured using natural abundance  $^{13}\text{C}$  solid state NMR spectroscopy as described previously [25, 28]. Shortly, the absolute values of the order parameters were determined from the dipolar splittings given by the indirect dimension of 2D R-PDFL experiment [29] and the signs were measured using S-DROSS experiments [30].

<sup>2</sup>Details of the used spectrometer and maybe some other details should be given.

<sup>3</sup>Sample preparation should be described.

<sup>4</sup>How is the peak assignment done?

### Molecular dynamics simulations

Molecular dynamics simulation data was collected using the Open Collaboration method [18]. The NMRLipids project blog (nmrlipids.blogspot.fi) and the GitHub repository (github.com/NMRLipids/NMRLipidsIVotherHGs) were used as the communication platforms. The simulated systems are listed in Table II and simulation details are given in the SI. The simulation data is also indexed in the searchable database (nmrlipids.fi), and in the NMRLipids/MATCH GitHub repository (https://github.com/NMRLipids/MATCH).

The C-H bond order parameters were calculated directly from the definition

$$S_{\text{CH}} = \frac{1}{2} \langle 3 \cos^2 \theta - 1 \rangle, \quad (1)$$

TABLE I: List of MD simulations without additional salt. CKPM refers to the version with Berger/Chiu  $\text{NH}_3$  charges compatible with Berger (i.e. the  $\text{NH}_3$  group having the same charges as in the  $\text{N}(\text{CH}_3)_3$  group of the PC lipids; 'M' stands for Mukhopadhyay after the first published Berger-based PS simulation that used these charges) and CKP refers to the version with more Gromos compatible version (i.e. the charges for the  $\text{NH}_3$  group taken from the lysine side-chain).

| lipid/counter-ions  | force field for lipids / ions | NaCl (mM) | $\text{CaCl}_2$ (mM) | $^a\text{N}_l$ | $^b\text{N}_w$ | $^c\text{N}_c$ | $^d\text{T}$ (K) | $^e t_{\text{sim}}$ (ns) | $^f t_{\text{anal}}$ (ns) | $^g$ files |
|---------------------|-------------------------------|-----------|----------------------|----------------|----------------|----------------|------------------|--------------------------|---------------------------|------------|
| DOPS/ $\text{Na}^+$ | CHARMM36 [31]                 | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [32]       |
| DOPS/ $\text{Na}^+$ | CHARMM36ua [?] ] 6.           | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [33]       |
| DOPS/ $\text{Na}^+$ | Slipids [34]                  | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [35]       |
| DOPS/ $\text{Na}^+$ | Slipids [34]                  | 0         | 0                    | 288            | 11232          | 0              | 303              | 200                      | 100                       | [36]       |
| DOPS/ $\text{Na}^+$ | Berger [37]                   | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [38]       |
| DOPS/ $\text{Na}^+$ | GROMOS-CKP1 [?] ] 7.          | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [39]       |
| DOPS/ $\text{Na}^+$ | GROMOS-CKP2 [?] ] 8.          | 0         | 0                    | 128            | 4480           | 0              | 303              | 500                      | 100                       | [40]       |
| DOPS/ $\text{Na}^+$ | lipid17 [41] / JC [42]        | 0         | 0                    | 128            | 4480           | 0              | 303              | 600                      | 100                       | [43]       |
| DOPS/ $\text{Na}^+$ | lipid17 [41] / ff99 [44]      | 0         | 0                    | 128            | 4480           | 0              | 303              | 600                      | 100                       | [45]       |
| POPS/ $\text{Na}^+$ | CHARMM36 [31]                 | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [46]       |
| POPS/ $\text{K}^+$  | CHARMM36 [31]                 | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [47]       |
| POPS/ $\text{Na}^+$ | CHARMM36ua [?] ] 9.           | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [48]       |
| POPS/ $\text{Na}^+$ | Slipids [34]                  | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [49]       |
| POPS/ $\text{Na}^+$ | Berger [?] ]                  | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [50]       |
| POPS/ $\text{Na}^+$ | MacRog [51]                   | 0         | 0                    | ?              | ??             | 0              | ?                | ?                        | ?                         | [?] ] 10.  |
| OPPS/ $\text{Na}^+$ | MacRog [51]                   | 0         | 0                    | 128            | 5120           | 0              | 298              | 200                      | 100                       | [52]       |
| POPS/ $\text{Na}^+$ | GROMOS-CKPM [?] ] 11.         | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [53]       |
| POPS/ $\text{Na}^+$ | GROMOS-CKP [?] ] 12.          | 0         | 0                    | 128            | 4480           | 0              | 298              | 500                      | 100                       | [54]       |
| POPS/ $\text{Na}^+$ | lipid17 [41] / JC [42]        | 0         | 0                    | 128            | 4480           | 0              | 298              | 600                      | 100                       | [55]       |
| POPS/ $\text{Na}^+$ | lipid17 [41] / ff99 [44]      | 0         | 0                    | 128            | 4480           | 0              | 298              | 600                      | 100                       | [56]       |

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

where  $\theta$  is the angle between the C-H bond and the membrane normal. Angular brackets point to the average over all sampled configurations. 29. Error estimation should be discussed.

The number density profiles were calculated using *gmx density* tool from Gromacs software package [63].

## RESULTS AND DISCUSSION

### Headgroup and glycerol backbone order parameters measured from POPS lipid bilayer

The INEPT and 2D R-PDLF experiments from POPS sample give well resolved spectras for all the carbons in headgroup and glycerol backbone region, except for  $g_3$  for which the resolution was not sufficient to determine the numerical value of the order parameter (Fig. 1). Slices of the R-PDLF spectra (Fig. 1 C)) show a single splitting for the  $\beta$ -carbon with the order parameter value of 0.12, and a superposition of a large and a very small splitting for the  $\alpha$ -carbon. The

larger splitting gives a order parameter value of 0.09, while the numerical value from the small splitting cannot resolved with the available resolution. Since only the absolute values of the PS headgroup order parameters were measured previously [7, 16], we used the S-DROSS experiment [?] ] to determine the signs of the order parameters. The S-DROSS slice for the  $\beta$ -carbon (Fig. 1 D)) clearly shows that the order parameter is negative, which is confirmed by SIMPSON simulations. The beginning of the S-DROSS slice suggests that the higher order parameter of the  $\alpha$ -carbon is positive and the deviation towards negative values with the longer  $T_1$  times suggests that the smaller order parameter is negative. This is confirmed by a SIMPSON simulation where the value of -0.02 was taken from  $^2\text{H}$  NMR experiment [16] for the smaller order parameter. The literature value was used because the resolution of our experiment was not sufficient to determine the small value of the order parameter. The S-DROSS curve from SIMPSON simulation with a positive value for the smaller order parameter (dashed grey in Fig. 1 D)) did not agree with the experiment, confirming the interpretation that the smaller order

TABLE II: List of MD simulations. The salt concentrations calculated as  $[\text{salt}] = N_c \times [\text{water}] / N_w$ , where  $[\text{water}] = 55.5$  M. CKPM refers to the version with Berger/Chiu  $\text{NH}_3$  charges compatible with Berger (i.e. the  $\text{NH}_3$  group having the same charges as in the  $\text{N}(\text{CH}_3)_3$  group of the PC lipids; 'M' stands for Mukhopadhyay after the first published Berger-based PS simulation that used these charges [? ]) and CKP refers to the version with more Gromos compatible version (i.e. the charges for the  $\text{NH}_3$  group taken from the lysine side-chain).

| lipid/counter-ions                            | force field for lipids / ions | NaCl (mM) | $\text{CaCl}_2$ (mM) | $^a N_l$ | $^b N_w$ | $^{13} N_c$ | $^d T$ (K) | $^e t_{\text{sim}}$ (ns) | $^f$ |
|---|-------------------------------|-----------|----------------------|----------|----------|-------------|------------|--------------------------|------|
| POPC:POPS (5:1)/ $\text{K}^+$                 | CHARMM36 [31, 57]             | 0         | 0                    | 110:22   | 4935     | 0           | 298        | 100                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | CHARMM36 [31, 57]             | 0         | 0                    | 250:50   | ?        | 0           | 298        | 200                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | CHARMM36 [31, 57]             | 0         | 0                    | 110:22   | 4620     | 0           | 298        | 500                      |      |
| POPC:POPS (5:1)/ $\text{Na}^+$                | CHARMM36 [31, 57]             | 0         | 0                    | 110:22   | 4620     | 0           | 298        | 500                      |      |
| POPC:POPS (1:1)/ $\text{K}^+$                 | CHARMM36 [31, 57]             | 0         | 0                    | 150:150  | ?        | 0           | 298        | 200                      |      |
| POPC:POPS (5:1)                               | CHARMM36 [31, 57, 61]         | 0         | 150 <b>17.</b>       | 250:50   | ?        | ?           | 298        | 200                      |      |
| POPC:POPS (5:1)                               | CHARMM36 [31, 57, 61]         | 0         | 1000 <b>19.</b>      | 250:50   | ?        | ?           | 298        | 200                      |      |
| POPC:POPS <sup>21</sup> . (5:1)/ $\text{K}^+$ | MacRog [51]                   | 0         | 0                    | 120:24   | 5760     | 0           | 298        | 200                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | MacRog [51]                   | 0         | 100                  | 120:24   | 5760     | 10          | 298        | 200                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | MacRog [51]                   | 0         | 300                  | 120:24   | 5760     | 31          | 298        | 200                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | MacRog [51]                   | 0         | 1000                 | 120:24   | 5760     | 104         | 298        | 200                      |      |
| POPC:POPS (5:1)/ $\text{K}^+$                 | MacRog [51]                   | 0         | 3000                 | 120:24   | 5760     | 311         | 298        | 200                      |      |

**27. MacRog simulations with KCl to be added**

**28. Berger simulations with NaCl and  $\text{CaCl}_2$  to be added**

<sup>a</sup>Number of lipid molecules with largest mole fraction

<sup>b</sup>Number of water molecules

<sup>c</sup>Number of additional cations

<sup>d</sup>Simulation temperature

<sup>e</sup>Total simulation time

<sup>f</sup>Time used for analysis

<sup>g</sup>Reference for simulation files

parameter is negative.

The headgroup and glycerol backbone order parameters of POPS measured in this work are in good agreement with the previously reported values from  $^2\text{H}$  NMR experiments of DOPS [7] (Fig. 2). The  $\beta$ -carbon order parameter is significantly more negative and  $\alpha$ -carbon experiences a significant forking in PS headgroup when compared with the values previously measured for POPC [28] (Fig. 2). These features have been interpreted to arise from a rigid PS headgroup conformation, stabilized by hydrogen bonds or electrostatic interactions [7, 8], but detailed structural interpretation is not available.

#### Headgroup and glycerol backbone in simulations of PS lipid bilayers without additional ions

The headgroup order parameters of DOPS and POPS bilayers from different simulation models are compared with the experimental data in Fig. 3. Subjective ranking of the quality of the models is shown in Fig. 4. The tested models perform generally less well than in the previous study for PC headgroup [18] and none of the models reproduce the experimental order parameters within experimental error bars. Therefore, the models cannot be straightforwardly used to interpret the structure of PS headgroup. However, the differences between PC and PS headgroups are partially reproduced by some of the models.

The two best performing models for the  $\alpha$  and  $\beta$ -carbons of PS, Slipids and CHARMM36, reproduce the large forking in the  $\alpha$ -carbon and the Slipids model gives also a good agreement with experiments for the  $\beta$ -carbon order parameter in both PC and PS headgroups (Fig. 3 and Ref. 18). Interestingly, the dihedral angle distributions of CHARMM36 and Slipids in Fig. 16 share significant similarities in the headgroup region. However, the experimental order parameters in the glycerol backbone region are not well reproduced by the Slipids model, which was also the case in for PC lipids [18]. This difference probably arises from the differences in the dihedral angle distributions of C1-C2-C3-O31 and C2-C3-O31-C31 in Figure 15, which are also illustrated in Figure 5.

**35. Also the discussion about POPS/OPPS issue with MacRog model should be added.**

#### Counterion binding to lipid bilayers containing PS lipids

Membranes containing PS lipids are always accompanied with counterions, which modulate electrostatic interactions between lipids and other biomolecules. Counterions are also suggested screen the repulsion between charged lipid headgroups in MD simulations and reduce the area per lipid of PS bilayers to be smaller than in PC bilayers [37, 64, 65]. The counterion density profiles along membrane normal show significant differences between simulation models (Fig. 6).




FIG. 1: (a) The headgroup region of the INEPT spectrum with headgroup and glycerol backbone carbons assigned. (b) 2D R-PDLF spectra for headgroup and glycerol backbone regions. (c) Slices for  $\alpha$  and  $\beta$  carbons. (d) Experimental SDRoss data (points) and SIMPSON simulations (lines). Order parameter values of -0.12 for the  $\beta$ -carbon, and 0.09 and -0.02 for the larger and smaller  $\alpha$ -carbon splittings were used in the SIMPSON calculations. The S-DRoss curve from SIMPSON simulation with positive value for the smaller order parameter (dashed grey).

30. This is preliminary figure, should be polished. 31. Should we show slices for all the analyzed carbons in (c)?

The strongest counterion binding, i.e., the lowest concentrations in bulk water, are observed in MacRog, Berger and Lipid17/JC simulations. CHARMM36, CHARMM36ua and Gromos-CKP models exhibit two local maxima in counterion density, while a single maxima is observed in the other models. 36. More detailed discussion may be possible after comparing monovalent ion binding to bilayers between CHARMM simulations and experiments. Area per lipid is in agreement with experiments [66] only in the Gromos-CKP models, while other models give significantly lower values (Fig. 6). The difference cannot be explained by the electrostatic screening of the headgroup repulsion due




FIG. 2: Headgroup and glycerol backbone order parameters of POPS measured in this work compared with values for DOPS ( $^2\text{H}$  NMR, 0.1M of NaCl) [7] and POPC ( $^{13}\text{C}$  NMR) [28] from literature. Signs for PS order parameters as measured in this work and signs for PC as measured in Refs [25? ].

32. There should be values in [15] which should be added.

to counterion binding because CHARMM36, CHARMM36ua and Slipid models give smaller area per lipid than Gromos-CKP models with similar counterion binding affinity.

To evaluate counterion binding in different simulation models against experimental data [15], we plot the headgroup order parameters measured from POPC:POPS 5:1 mixture as a function of different monovalent ions added to the buffer (Fig. 7). Experimental order parameter data for POPC headgroup in the mixture is available as a function of LiCl and KCl concentrations, while POPS headgroup order parameters are measured also as a function of NaCl. Lithium interacts more strongly with PS headgroups than other monovalent ions [11, 12, 14, 15, 67], as also observed for PC headgroups [68]. This is evident also in the changes of PS headgroup order parameters, which decrease with the addition of lithium but increase with the addition of sodium or potassium (Fig. 7). POPC headgroup order parameters exhibit a clear decrease as a function of LiCl concentration but only modest changes as a function of KCl concentration, indicating significant  $\text{Li}^+$  binding but only weak  $\text{Na}^+$  binding to the mixture when interpreted using the electrometer concept [69–71]. In simulations with the Berger model, the headgroup order parameter response of POPC to the added NaCl is similar to the




FIG. 3: Order parameters for PS headgroup and glycerol backbone from simulations with different models and experiments without  $CaCl_2$ . All DOPS data at 303 K, POPS at 298 K. Experimental data from [7] contain 0.1 M of NaCl. Signs are taken from experiments for POPS described in Supplementary Information. The vertical bars shown are not error bars, but demonstrate that we had at least two data sets; the ends of the bars mark the extreme values from the sets, and the dot marks their measurement-time-weighted average.

|              | $\beta$ | $\alpha$ | $g_3$ | $g_2$ | $g_1$ | $\Sigma$ |
|--------------|---------|----------|-------|-------|-------|----------|
| CHARMM 36    | M       | M F      | M     | M     | M F   | 8        |
| CHARMM 36-UA | M       | M        | M     | M     | M F   | 8        |
| GROMOS-CKP1  | M       | M F      | M F   | M     | M F   | 14       |
| GROMOS-CKP2  | M       | M F      | M F   | M     | M F   | 14       |
| Slipid       | M       | M        | M F   | M     | M F   | 14       |
| Berger       | M       | M F      | M F   | M     | M F   | 15       |

FIG. 4: Rough subjective ranking of force fields based on Figure 3. Here M indicates a magnitude problem, F a forking problem; letter size increases with problem severity. Color scheme: within experimental error (dark green), almost within experimental error (light green), clear deviation from experiments (light red), and major deviation from experiments (dark red). The  $\Sigma$ -column shows the total deviation of the force field, when individual carbons are given weights of 0 (matches experiment), 1, 2, and 4 (major deviation). For full details of the assessment, see Supplementary Information.

33.Issue about possible updates to this plot:

<https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/4>

34.Lipid17 and MacRog results should be added into this plot.




FIG. 5: Snapshots overlaid from different simulations for glycerol backbone region by Pavel Buslaev.

experiments of LiCl, indicating overestimated binding affinity of sodium, in line with the results for PC bilayers [19]. Indeed, the sodium density profile shows a significant binding peak in the Berger model (Fig. 8). Potassium binding in the MacRog simulation is significantly weaker (Fig. 8) and the headgroup order parameter changes are also in better agreement with simulations (Fig. 7). 37.Discussion about Lipid17 to be written when we have the density profiles. All the tested models overestimate the changes of POPS headgroup order parameters as a function of monovalent ions (Fig. 7), suggesting that model development is necessary to interpret the PS headgroup-ion interactions from MD simulations.






FIG. 6: Counterion densities of POPS lipid bilayer along the membrane normal from simulations with different force fields.

### Headgroup structure in PS and PC mixtures

Dilution of PS lipid bilayers with PC lipids reduces propensity for the formation of strong complexes with multivalent ions and is proposed to make PS headgroups less rigid [7, 8, 15, 16]. Therefore, the intermolecular interactions at the headgroup region seems to determine important physical properties of mixed lipid bilayers. The headgroup order parameters of POPC increase in experiments with increasing amount of POPS in the mixed lipid bilayer (Fig. 9) [27]. This behaviour is generally observed when negatively charged lipids or sur-

factants are mixed with PC lipids [27, 72] and can be understood by the tilting of lipid headgroup more parallel to the membrane plane according to the electrometer concept [71]. The headgroup order parameters of PS lipids shift closer to zero when bilayer is diluted with PC lipids in experiments (Fig. 9) [7, 15, 27], which is interpreted to indicate reduced rigidity [7, 8].

The increase of POPC headgroup order parameters with the increasing amount of negatively charged POPS lipid is reproduced in MacRog simulations with potassium counterions, but not in Berger simulations with sodium or in CHARMM36 simulations with potassium or sodium counterions (Fig. 9). The observations can be explained using the electrometer concept. The Berger simulation exhibits very strong sodium binding (Fig. 8), which surpasses the effect of negatively charged lipids as the amount of counterions increase with increasing amount of PS. In CHARMM36 simulations, the counterion binding neutralizes the effect of PS and the headgroup order parameters are not changed with increasing amount of PS. Finally, the weak binding of potassium in the MacRog simulations enables the increase of order parameters with the added amount of PS (Figs. 9 and 8).

Oppositely to experiments, the headgroup order parameter of POPS shift away from zero in CHARMM36 simulations when bilayer is diluted with POPC (Fig. 9). In lipid14/17 simulations, the POPS order parameter shift closer to zero when bilayer is diluted with POPC, but the numerical values of order parameters are too far from experiments to enable interpretation of the experimental data. Therefore, we conclude that the force field development is necessary before MD simulations can be used to interpret the interactions between PC and PS headgroups.

### Ca<sup>2+</sup> binding affinity in bilayers with negatively charged PS lipids

The headgroup order parameters of PC lipids decrease proportionally to the bound positive charge in to a bilayer [19, 71] and can be therefore used to measure the ion binding affinity. This molecular electrometer concept can be also applied to lipid bilayers with mixtures of PC and negatively charged lipids [15, 73, 74] (see Fig. 12).

The headgroup order parameter changes of POPC and POPS from POPC:POPS (5:1) mixtures are shown in Fig. 10 as a function of Ca<sup>2+</sup> concentration from different simulations and experiments [15]. The ion density distributions from the simulations are shown in Fig. 11. The results suggest that Ca<sup>2+</sup> ions clearly overbind in simulations with MacRog and Berger models, as expected from previous results for PC lipid bilayers [19]. It should be noted, however, that the lowest concentration (100mM) gives a good agreement with experiments. **43. Should be analyze/discuss this further? Binding with ~100 mM is saturated in both Berger and MacRog simulations. Maybe this is realistic? It should be noted that Berger simulation do not have counterions.. Surprisingly, the calcium binding seems to be too weak CHARMM36 simulations. This**




FIG. 7: Changes of the PC (left) and PS (right) headgroup order parameters as a function of added NaCl, KCl and LiCl from POPC:POPS (5:1) mixture. The experimental data is from Ref. 15. The values from counterion-only systems are set as a zero point of y-axis. To correctly illustrate the significant forking of the  $\alpha$ -carbon order parameter in PS headgroup (bottom, right), the y-axis is transferred with the same value for both order parameters such that the lower order parameter value is at zero.

38.CHARM36 results for this plot would be highly useful.

is due to the NBfix interaction parameters from Ref. 61, incorporated in the parameters from CHARMM-GUI at the time of running the simulations (January 2018). The binding of calcium to pure POPC bilayer is also too weak with these parameters as shown in Figs. 18 and 19. 44.The discussion is to be finished when we have all the data in the plot.

Also the order parameters of PS headgroup from POPC:POPS (5:1) mixture are shown in Fig. 10 as a function of  $\text{CaCl}_2$  concentration. In experiments the order parameters exhibit a strong dependence of  $\text{CaCl}_2$  with small concentrations with a rapid saturation around 50 mM. The changes of PS headgroup order parameters with added  $\text{CaCl}_2$  are overestimated in all tested simulation models. Furthermore, the changes of the headgroup order parameters do not qualitatively agree with experiments. This is in contrast to previ-

ous results for PC headgroup [19], where qualitatively correct response to bound ions was observed despite of significant discrepancies in the headgroup structure without additional ions.

## CONCLUSIONS




FIG. 8: Counterion density distributions from PC:PS mixtures.

39.Lipid 17 is to be added.

## SUPPLEMENTARY INFORMATION

### Simulated systems

#### CHARMM36

51.To be written by Piggot, Madsen and Ollila

#### CHARMM36ua

52.To be written by Piggot

#### Slipids

53.To be written by Piggot and Favela

### Berger

54.To be written by Piggot and Ollila Simulations with sodium were taken directly from Ref. ? and simulations with calcium directly from 17. Simulation of POPC at 310 K was taken directly from Ref. 75.

### GROMOS-CKP

55.To be written by Piggot

### Lipid17

56.To be written by Kav and Miettinen

### MacRog

57.To be written by Javanainen and Piggot

### Cation binding affinity to lipid bilayers with different amount of charge

Before using the headgroup order parameters to compare ion binding affinity between simulations and experiments, it is important to quantify the response of the order parameters to the bound charge in simulations. The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [72] In Fig. 14. The figure shows that the order parameters are too sensitive to bound charge in Lipid14 model, while CHARMM36 is in better agreement with experiments. This has to be taken into account when analysin the binding affinities.

### Difference between POPC and OPPS in MacRog model

#### Dihedrals

#### Dihedrals

The experimental results show essentially no changes in the order parameters as a function of added NaCl, while significant changes are observed in simulations. However, the minimum buffer concentration of NaCl in the experimental was 100mM [67]. Therefore, we cannot exclude the possibility that the NaCl induced changes were already saturated with 100mM NaCl concentration, which was the case for CaCl<sub>2</sub> in Fig. 10.






FIG. 9: Changes of PC (left panel) and PS (right panel) headgroup order parameters from POPC:POPS mixtures with increasing amount of POPS. Experimental results of POPC are taken from Ref. 27 (signs are determined as discussed in [18, 20]). Experimental values for POPS in pure bilayer and in mixture are measured in this work and in Ref. 15 at 298K, respectively. Since the experimental data of POPS in pure and diluted mixture come from different experimental sets (13C NMR in this work and 2H NMR from Ref. 15), the experimental change of the order parameter is less accurate than in typical measurements where same technique is used in all conditions, see discussion about qualitative and quantitative accuracy in Ref. 20. For POPC (left panel) the zero point of y-axis is set to the value of pure bilayer. For  $\beta$ -carbon of POPS (right panel, top) the zero point of y-axis is set to the value from POPC:POPS (5:1) mixture. For  $\alpha$ -carbon of POPS (right panel, bottom) the y-axis is transferred with the same value for both order parameters such that the lower order parameter value from POPC:POPS (5:1) mixture is at zero to correctly illustrate the significant forking.

40. Simulation of CHARMM36 at 298K should be maybe rerun with Gromacs 5.

41. Simulation of pure POPC at 298K with Lipid14 would be useful for this plot (only at 303 K is available from NMRLipids I)

42. MacRog simulations of pure POPS with potassium counterions only would be useful for this and other plots.

#### Details of the rough subjective force field ranking (Fig. 4)

The assessment was based fully on the Fig. 3. First, for each carbon (the columns in Fig. 3) in each force field (the rows), we looked separately at deviations in magnitude and forking.

**Magnitude** deviations, i.e., how close to the experimentally obtained C–H order parameters (OPs) the force-field-produced OPs were. For each carbon, the following 5-step scale was used:

**0 ( )**: More than half of all the calculated OPs (that is, of all different hydrogens in all different lipids) were within the *subjective sweet spots* (SSP, blue-shaded areas in Fig. 3).

**1 (m)**: All the calculated OPs were < 0.03 units away

from the SSP.

**2 (M)**: All the calculated OPs were < 0.05 units away from the SSP.

**3 (M)**: All the calculated OPs were < 0.10 units away from the SSP.

**4 (M)**: Some of the calculated OPs were > 0.10 units away from the SSP.

**Forking** deviations, i.e., how well the difference in order parameters of two hydrogens attached to a given carbon matched that obtained experimentally. Note that this is not relevant for  $\beta$  and  $g_2$ , which have only one hydrogen. For the  $\alpha$  carbon, for which a considerable forking of 0.105 is experimentally seen, the following 5-step scale was used:




FIG. 10: Changes of POPC (left) and POPS (right) headgroup order parameters in POPC:POPS (5:1) mixture as a function  $CaCl_2$  concentration. Experimental data is taken from 15. The values from counterion-only systems are set as a zero point of y-axis. To correctly illustrate the significant forking of the  $\alpha$ -carbon order parameter in PS headgroup (bottom, right), the y-axis is transferred with the same value for both order parameters such that the lower order parameter value is at zero.

45. Information about the counterions in different simulations should be added

46. Upcoming simulations with original CHARMM36 have been mentioned in the blog:

<http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1520090718976#c5569269391707740056>

47. Upcoming Lipid17 simulations have been mentioned in the blog

<http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1515177306419#c994825612316235467>

**0 ( ):** The distance  $D$  between the dots (that mark the measurement-time-weighted averages in Fig. 3) was  $0.08 < D < 0.13$  units for all the calculated OPs (that is, for all different lipids).

**1 (F):**  $(0.06 < D < 0.08)$  OR  $(0.13 < D < 0.15)$ .

**2 (F):**  $(0.04 < D < 0.06)$  OR  $(0.15 < D < 0.17)$ .

**3 (F):**  $(0.02 < D < 0.04)$  OR  $(0.17 < D < 0.19)$ .

**4 (F):**  $(D < 0.02)$  OR  $(0.19 < D)$ .

For the  $g_3$  carbon, for which no forking is indicated by experiments, the following 5-step scale was used:

**0 ( ):**  $D < 0.02$ .

**1 (F):**  $0.02 < D < 0.04$ .

**2 (F):**  $0.04 < D < 0.06$ .

**3 (F):**  $0.06 < D < 0.08$ .

**4 (F):**  $0.08 < D$ .

For the  $g_1$  carbon, for which a considerable forking of 0.13 is experimentally seen, the following 5-step scale was used:

**0 ( ):**  $0.11 < D < 0.15$ .

**1 (F):**  $(0.09 < D < 0.11)$  OR  $(0.15 < D < 0.17)$ .

**2 (F):**  $(0.07 < D < 0.09)$  OR  $(0.17 < D < 0.19)$ .




FIG. 11: Ca<sup>2+</sup> density profiles from simulations.

48. The CHARMM results are mass densities, numbers should be used.

49. Should we include also counterions into the plot?

50. Not all the data from MacRog is included.

3 (**F**): ( $0.05 < D < 0.07$ ) OR ( $0.19 < D < 0.21$ ).

4 (**F**): ( $D < 0.05$ ) OR ( $0.21 < D$ ).

Based on these assessments of magnitude and forking deviations, each carbon was then assigned to one of the following groups: "within experimental error" (magnitude and forking deviations both on step 0 of the scales described above), "almost within experimental error" (sum of the magnitude and forking deviation steps 1 or 2), "clear deviation from experiments" (sum of magnitude and forking deviation steps from 3 to 5), and "major deviation from experiments" (sum of magnitude and forking deviation steps from 6 to 8). These groups are indicated by colors in Fig. 4. (Note that for  $\beta$  and  $g_2$ , for which there can be no forking, the corresponding group assignment limits were: 0, 1, 2, and 3.)




FIG. 12: The change of PC headgroup order parameters as a function of CaCl<sub>2</sub> measured from bilayers containing different amount of negatively charged lipids. The values are taken from 2H NMR experiments reported in the literature (DPPC [69], POPC [70], POPC:POPS (5:1) [15], POPC:POPG mixtures with 0.1M NaCl [74] and POPC:POPG (1:1) without NaCl [73]). As expected, the decrease of order parameters with the added CaCl<sub>2</sub> is more pronounced for systems with larger fraction of negatively charged lipids, indicating larger amount of bound cations.

Finally, the total ability of the force field to describe the headgroup and glycerol structure was estimated. To this end, the groups were given the following weights: 0 (within experimental error), 1 (almost within experimental error), 2 (clear deviation from experiments), 4 (major deviation from experiments), and the weights of the five carbons were summed up. The sum, given in the  $\Sigma$ -column of Fig. 3, was then used to (roughly and subjectively, as should be clear from the above description) rank the force fields.




FIG. 13: The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [72].




FIG. 14: Headgroup order parameters from POPS and OPPS simulations with MacRog model.

\* samuli.ollila@helsinki.fi

- [1] M. A. Lemmon, Nat. Rev. Mol. Cell Biol. **9**, 99 (2008).
- [2] P. A. Leventis and S. Grinstein, Annual Review of Biophysics **39**, 407 (2010).
- [3] L. Li, X. Shi, X. Guo, H. Li, and C. Xu, Trends in Biochemical Sciences **39**, 130 (2014), ISSN 0968-0004.
- [4] T. Yeung, G. E. Gilbert, J. Shi, J. Silvius, A. Kapus, and S. Grinstein, Science **319**, 210 (2008).
- [5] H. Zhao, E. K. J. Tuominen, and P. K. J. Kinnunen, Biochemistry **43**, 10302 (2004).
- [6] G. P. Gorbenko and P. K. Kinnunen, Chemistry and Physics of Lipids **141**, 72 (2006).
- [7] J. L. Browning and J. Seelig, Biochemistry **19**, 1262 (1980).
- [8] G. Büldt and R. Wohlgenuth, The Journal of Membrane Biology **58**, 81 (1981), ISSN 1432-1424, URL <http://dx.doi.org/10.1007/BF01870972>.
- [9] H. Hauser, E. Finer, and A. Darke, Biochemical and Biophysical Research Communications **76**, 267 (1977), ISSN 0006-291X, URL <http://www.sciencedirect.com/science/article/pii/0006291X77907215>.
- [10] R. J. Kurland, Biochemical and Biophysical Research Communications **88**, 927 (1979), ISSN 0006-291X, URL <http://www.sciencedirect.com/science/article/pii/0006291X79914979>.
- [11] H. Hauser and G. G. Shipley, Biochemistry **22**, 2171 (1983).
- [12] H. Hauser and G. Shipley, Biochimica et Biophysica Acta (BBA) - Biomembranes **813**, 343 (1985), ISSN 0005-2736, URL <http://www.sciencedirect.com/science/article/pii/0005273685902512>.
- [13] G. W. Feigenson, Biochemistry **25**, 5819 (1986).
- [14] J. Mattai, H. Hauser, R. A. Demel, and G. G. Shipley, Biochemistry **28**, 2322 (1989).
- [15] M. Roux and M. Bloom, Biochemistry **29**, 7077 (1990).
- [16] M. Roux and M. Bloom, Biophys. J. **60**, 38 (1991).
- [17] A. Melcrová, S. Pokorna, S. Pullanchery, M. Kohagen, P. Jurkiewicz, M. Hof, P. Jungwirth, P. S. Cremer, and L. Cwiklik, Sci. Reports **6**, 38035 (2016).
- [18] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduč, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., J. Phys. Chem. B **119**, 15075 (2015).
- [19] A. Catte, M. Giryh, M. Javanainen, C. Loison, J. Melcr, M. S. Miettinen, L. Monticelli, J. Maatta, V. S. Oganessian, O. H. S. Ollila, et al., Phys. Chem. Chem. Phys. **18**, 32560 (2016).
- [20] O. S. Ollila and G. Pabst, Biochimica et Biophysica Acta (BBA) - Biomembranes **1858**, 2512 (2016).
- [21] J. Seelig, Cell Biology International Reports **14**, 353 (1990), ISSN 0309-1651, URL <http://www.sciencedirect.com/science/article/pii/030916519091204H>.
- [22] C. G. Sinn, M. Antonietti, and R. Dimova, Colloids and Surfaces A: Physicochemical and Engineering Aspects **282-283**, 410 (2006), a Collection of Papers in Honor of Professor Ivan B. Ivanov (Laboratory of Chemical Physics and Engineering, University of Sofia) Celebrating his Contributions to Colloid and Surface Science on the Occasion of his 70th Birthday.
- [23] P. T. Vernier, M. J. Ziegler, and R. Dimova, Langmuir **25**, 1020 (2009).
- [24] J. M. Boettcher, R. L. Davis-Harrison, M. C. Clay, A. J. Nieuwkoop, Y. Z. Ohkubo, E. Tajkhorshid, J. H. Morrissey, and C. M. Rienstra, Biochemistry **50**, 2264 (2011).
- [25] T. M. Ferreira, R. Sood, R. Bärenwald, G. Carlström, D. Topgaard, K. Saalwächter, P. K. J. Kinnunen, and O. H. S. Ollila,




FIG. 15: Dihedral angle distributions of bonds from phosphate to acyl chain carbonyls from different simulation models.






FIG. 16: Dihedral angle distributions of bonds from phosphate to headgroup from different simulation models.

- Langmuir **32**, 6524 (2016).
- [26] H. U. Gally, G. Pluschke, P. Overath, and J. Seelig, *Biochemistry* **20**, 1826 (1981).
- [27] P. Scherer and J. Seelig, *EMBO J.* **6** (1987).
- [28] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
- [29] S. V. Dvinskikh, H. Zimmermann, A. Maliniak, and D. Sandstrom, *J. Magn. Reson.* **168**, 194 (2004).
- [30] J. D. Gross, D. E. Warschawski, and R. G. Griffin, *J. Am. Chem. Soc.* **119**, 796 (1997).
- [31] R. M. Venable, Y. Luo, K. Gawrisch, B. Roux, and R. W. Pastor, *The Journal of Physical Chemistry B* **117**, 10183 (2013).
- [32] T. Piggot, *CHARMM36 DOPS simulations (versions 1 and 2) 303 K 1.0 nm LJ switching* (2017), URL <https://doi.org/10.5281/zenodo.1129411>.
- [33] T. Piggot, *CHARMM36-UA DOPS simulations (versions 1 and 2) 303 K 1.0 nm LJ switching* (2017), URL <https://doi.org/10.5281/zenodo.1129456>.
- [34] J. P. M. Jämbek and A. P. Lyubartsev, *Phys. Chem. Chem. Phys.* **15**, 4677 (2013).
- [35] T. Piggot, *Slipids DOPS simulations (versions 1 and 2) 303 K 1.0 nm cut-off with LJ-PME* (2017), URL <https://doi.org/10.5281/zenodo.1129439>.
- [36] F. Favela-Rosales, *MD simulation trajectory of a fully hydrated DOPS bilayer: SLIPIDS, Gromacs 5.0.4*. 2017. (2017), URL <https://doi.org/10.5281/zenodo.495510>.
- [37] P. Mukhopadhyay, L. Monticelli, and D. P. Tieleman, *Biophysical Journal* **86**, 1601 (2004).
- [38] T. Piggot, *Berger DOPS simulations (versions 1 and 2) 303 K 1.0 nm cut-off* (2017), URL <https://doi.org/10.5281/zenodo.1129419>.
- [39] T. Piggot, *GROMOS-CKP DOPS simulations (versions 1 and 2) 303 K with Berger/Chiu NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129429>.
- [40] T. Piggot, *GROMOS-CKP DOPS simulations (versions 1 and 2) 303 K with GROMOS NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129447>.
- [41] I. Gould, A. Skjevik, C. Dickson, B. Madej, and R. Walker, *Lipid17: A comprehensive amber force field for the simulation of zwitterionic and anionic lipids* (2018), in preparation.
- [42] I. S. Joung and T. E. Cheatham, *The Journal of Physical Chemistry B* **112**, 9020 (2008).
- [43] B. Kav and M. S. Miettinen, *Molecular dynamics simulation trajectory of an anionic lipid bilayer: 100 mol% DOPS with Na+ counterions using Joung-Cheatham Ions* (2018), B.K acknowledges financial support from International Max Planck Research School on Multiscale Bio-Systems, URL <https://doi.org/10.5281/zenodo.1134871>.
- [44] J. Åqvist, *J. Phys. Chem.* **94**, 8021 (1990).
- [45] B. Kav and M. S. Miettinen, *Molecular dynamics simulation trajectory of an anionic lipid bilayer: 100 mol% DOPS with Na+ counterions using ff99 Ions* (2018), B.K acknowledges financial support from International Max Planck Research School on Multiscale Bio-Systems, URL <https://doi.org/10.5281/zenodo.1135142>.
- [46] T. Piggot, *CHARMM36 POPS simulations (versions 1 and 2) 298 K 1.0 nm LJ switching* (2017), URL <https://doi.org/10.5281/zenodo.1129415>.
- [47] T. Piggot, *CHARMM36 POPS simulations (versions 1 and 2)*




FIG. 17: Order parameters of PS headgroup as a function of added NaCl measured from DMPC:DMPS (3:1) mixture [67].




FIG. 18: The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [72].




FIG. 19: The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [72].

- 298 K 1.0 nm LJ switching with K ions (2018), URL <https://doi.org/10.5281/zenodo.1182654>.
- [48] T. Piggot, *CHARMM36-UA POPS simulations (versions 1 and 2) 298 K 1.0 nm LJ switching* (2017), URL <https://doi.org/10.5281/zenodo.1129458>.
- [49] T. Piggot, *Slipids POPS simulations (versions 1 and 2) 298 K 1.0 nm cut-off with LJ-PME* (2017), URL <https://doi.org/10.5281/zenodo.1129441>.
- [50] T. Piggot, *Berger POPS simulations (versions 1 and 2) 298 K 1.0 nm cut-off* (2017), URL <https://doi.org/10.5281/zenodo.1129425>.
- [51] A. Maciejewski, M. Pasenkiewicz-Gierula, O. Cramariuc, I. Vattulainen, and T. Róg, *J. Phys. Chem. B* **118**, 4571 (2014).
- [52] M. Javanainen, *Simulation of a pops bilayer* (2017), URL <https://doi.org/10.5281/zenodo.1120287>.
- [53] T. Piggot, *GROMOS-CKP POPS simulations (versions 1 and 2) 298 K with Berger/Chiu NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129431>.
- [54] T. Piggot, *GROMOS-CKP POPS simulations (versions 1 and 2) 298 K with GROMOS NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129435>.
- [55] M. S. Miettinen and B. Kav, *Molecular dynamics simulation trajectory of an anionic lipid bilayer: 100 mol% POPS with Na+ counterions using Joung-Cheatham Ions* (2018), B.K. acknowledges financial support from International Max Planck Research School on Multiscale Bio-Systems., URL <https://doi.org/10.5281/zenodo.1148495>.
- [56] M. S. Miettinen and B. Kav, *Molecular dynamics simulation trajectory of an anionic lipid bilayer: 100 mol% POPS*

- with Na<sup>+</sup> counterions using ff99 ions (2018), B.K. acknowledges financial support from International Max Planck Research School on Multiscale Bio-Systems, URL <https://doi.org/10.5281/zenodo.1134869>.
- [57] J. B. Klauda, R. M. Venable, J. A. Freites, J. W. O'Connor, D. J. Tobias, C. Mondragon-Ramirez, I. Vorobyov, A. D. MacKerell Jr, and R. W. Pastor, *J. Phys. Chem. B* **114**, 7830 (2010).
- [58] O. H. S. Ollila, *POPS+83%popc lipid bilayer simulation at T298K ran CHARMM\_GUI force field and Gromacs* (2017), URL <https://doi.org/10.5281/zenodo.1011104>.
- [59] T. Piggot, *CHARMM36 POPS/POPC simulations (versions 1 and 2) 298 K 1.0 nm LJ switching with K ions* (2018), URL <https://doi.org/10.5281/zenodo.1182658>.
- [60] T. Piggot, *CHARMM36 POPS/POPC simulations (versions 1 and 2) 298 K 1.0 nm LJ switching with Na ions* (2018), URL <https://doi.org/10.5281/zenodo.1182665>.
- [61] S. Kim, D. Patel, S. Park, J. Slusky, J. Klauda, G. Widmalm, and W. Im, *Biophysical Journal* **111**, 1750 (2016), ISSN 0006-3495, URL <http://www.sciencedirect.com/science/article/pii/S0006349516307615>.
- [62] M. Javanainen, *Simulations of popc/pops membranes with cacl2*. (2017), URL <https://doi.org/10.5281/zenodo.897467>.
- [63] M. Abraham, D. van der Spoel, E. Lindahl, B. Hess, and the GROMACS development team, *GROMACS user manual version 5.0.7* (2015), URL [www.gromacs.org](http://www.gromacs.org).
- [64] S. A. Pandit and M. L. Berkowitz, *Biophysical Journal* **82**, 1818 (2002).
- [65] U. R. Pedersen, C. Leidy, P. Westh, and G. H. Peters, *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1758**, 573 (2006).
- [66] J. Pan, X. Cheng, L. Monticelli, F. A. Heberle, N. Kucerka, D. P. Tieleman, and J. Katsaras, *Soft Matter* **10**, 3716 (2014).
- [67] M. Roux and J.-M. Neumann, *FEBS Letters* **199**, 33 (1986).
- [68] G. Cevc, *Biochim. Biophys. Acta - Rev. Biomemb.* **1031**, 311 (1990).
- [69] H. Akutsu and J. Seelig, *Biochemistry* **20**, 7366 (1981).
- [70] C. Altenbach and J. Seelig, *Biochemistry* **23**, 3913 (1984).
- [71] J. Seelig, P. M. MacDonald, and P. G. Scherer, *Biochemistry* **26**, 7535 (1987).
- [72] P. G. Scherer and J. Seelig, *Biochemistry* **28**, 7720 (1989).
- [73] F. Borle and J. Seelig, *Chemistry and Physics of Lipids* **36**, 263 (1985).
- [74] P. M. Macdonald and J. Seelig, *Biochemistry* **26**, 1231 (1987).
- [75] S. Ollila, M. T. Hyvönen, and I. Vattulainen, *J. Phys. Chem. B* **111**, 3139 (2007).

### ToDo

- |  |    |  |   |
|--|----|--|---|
| 1. Authorship query to be sent soon. . . . .   | P. | 9. Correct citation for CHARMMua DOPS . . . . .  | 2 |
| 2. Details of the used spectrometer and maybe some other details should be given. . . . .        | 1  | 10. Data to be added by Piggot . . . . .   | 2 |
| 3. Sample preparation should be described. . . . .   | 1  | 11. Correct citation(s) for CKP. . . . .   | 2 |
| 4. How is the peak assignment done? . . . . .  | 1  | 12. Correct citation(s) for CKP. . . . .   | 2 |
| 5. Should confirm that the amounts of water in experiments matched those in simulations. . . . . | 2  | 29. Error estimation should be discussed. . . . .  | 2 |
| 6. Correct citation for CHARMMua DOPS . . . . .  | 2  | 13. Should confirm that the amounts of water in experiments matched those in simulations. . . . .  | 3 |
| 7. Correct citation(s) for CKP. . . . .  | 2  | 14. Equilibration? . . . . .   | 3 |
| 8. Correct citation(s) for CKP. . . . .  | 2  | 15. Trajectories and further details to be added by J. Madsen . . . . .  | 3 |
|  |    | 16. Trajectories and further details to be added by J. Madsen . . . . .  | 3 |
|  |    | 17. Concentration to be checked . . . . .  | 3 |
|  |    | 18. Trajectories and further details to be added by J. Madsen . . . . .  | 3 |
|  |    | 19. Concentration to be checked . . . . .  | 3 |
|  |    | 20. Trajectories and further details to be added by J. Madsen . . . . .  | 3 |
|  |    | 21. This is also probably OPPS? These should be corrected in this table as well. . . . .   | 3 |
|  |    | 22. Equilibration? . . . . .   | 3 |
|  |    | 23. Equilibration? . . . . .   | 3 |
|  |    | 24. Equilibration? . . . . .   | 3 |
|  |    | 25. Equilibration? . . . . .   | 3 |
|  |    | 26. Equilibration? . . . . .   | 3 |
|  |    | 27. MacRog simulations with KCl to be added . . . . .  | 3 |
|  |    | 28. Berger simulations with NaCl and CaCl to be added . . . . .  | 3 |
|  |    | 35. Also the discussion about POPS/OPPS issue with MacRog model should be added. . . . .   | 3 |
|  |    | 30. This is preliminary figure, should be polished. . . . .  | 4 |
|  |    | 31. Should we show slices for all the analyzed carbons in (c)? . . . . .   | 4 |
|  |    | 36. More detailed discussion may be possible after comparing monovalent ion binding to bilayers between CHARMM simulations and experiments. . . . .  | 4 |
|  |    | 32. There should be values in [15] which should be added. . . . .  | 4 |
|  |    | 33. Issue about possible updates to this plot: <a href="https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/4">https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/4</a> . . . . .                               | 5 |
|  |    | 34. Lipid17 and MacRog results should be added into this plot. . . . .   | 5 |
|  |    | 37. Discussion about Lipid17 to be written when we have the density profiles. . . . .  | 5 |
|  |    | 43. Should be analyze/discuss this further? Binding with ~100 mM is saturated in both Berger and MacRog simulations. Maybe this is realistic? It should be noted that Berger simulation do not have counterions. . . . . | 6 |
|  |    | 38. CHARMM36 results for this plot would be highly useful. . . . .   | 7 |
|  |    | 44. The discussion is to be finished when we have all the data in the plot. . . . .  | 7 |
|  |    | 39. Lipid 17 is to be added. . . . .   | 8 |
|  |    | 51. To be written by Piggot, Madsen and Ollila . . . . .   | 8 |
|  |    | 52. To be written by Piggot . . . . .  | 8 |
|  |    | 53. To be written by Piggot and Favela . . . . .   | 8 |
|  |    | 54. To be wiritten by Piggot and Ollila . . . . .  | 8 |
|  |    | 55. To be written by Piggot . . . . .  | 8 |

|  |    |  |    |
|--|----|--|----|
| 56. To be written by Kav and Miettinen . . . . .   | 8  | 47. Upcoming Lipid17 simulations   |    |
| 57. To be written by Javanainen and Piggot . . . . .   | 8  | have been mentioned in the blog  |    |
| 40. Simulation of CHARMM36 at 298K should be   |    | <a href="http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1515177306419#c99482561">http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-</a>         |    |
| maybe rerun with Gromacs 5. . . . .  | 9  | <a href="http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1515177306419#c99482561">current-status-and.html?showComment=1515177306419#c99482561</a> |    |
| 41. Simulation of pure POPC at 298K with Lipid14   |    |  |    |
| would be useful for this plot (only at 303 K is available  |    |  |    |
| from NMRLipids I) . . . . .  | 9  |  |    |
| 42. MacRog simulations of pure POPS with potassium   |    | 48. The CHARMM results are mass densities, numbers   |    |
| counterions only would be useful for this and other plots.   | 9  | should be used. . . . .  | 11 |
| 45. Information about the cuonterions in different sim-  |    |  |    |
| ulations should be added . . . . .   | 10 |  |    |
| 46. Upcoming simulations with original   |    | 49. Should we include also counterions into the plot? .  | 11 |
| CHARMM36 have been mentioned in the blog:  |    |  |    |
| <a href="http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1520090718976#c556926039707740056">http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-</a>                   |    |  |    |
| <a href="http://nmrlipids.blogspot.com/2017/12/nmrlipids-iv-current-status-and.html?showComment=1520090718976#c556926039707740056">current-status-and.html?showComment=1520090718976#c556926039707740056</a> |    | 50. Not all data from MacRog is included. . . . .  | 11 |