

NMRlipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PS lipids

O. H. Samuli Ollila^{1,2,*}

¹*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic*

²*Institute of Biotechnology, University of Helsinki*

(Dated: February 7, 2018)

Primarily measured but also simulated NMR order parameters will be collected also for other than phosphatidylcholine (these are discussed in NMRlipids I) headgroup. The information will be used to understand structural differences between different lipid molecules in bilayers.

INTRODUCTION

Phosphatidylserine (PS) is the most common negatively charged lipid in eukaryotic membranes. PS lipids compose 8.5% of total lipid weight of erythrocytes, but the abundance varies between different organelles up to 25-35% in plasma membrane [1–3]. Despite of the relatively low abundance, PS lipids are important signaling molecules. They interact with signaling proteins [2], regulate surface charge and protein localization [4], and induce protein aggregation [? ?]. Some domains specifically interact PS lipids, while others are attracted by general electrostatics and the binding can be regulated by calcium [2]. Therefore, it is crucial to resolve the structural details of PS headgroup and calcium binding in order to fully understand PS mediated signaling processes.

The structure of PS lipid headgroups and their interactions with ions have been studied with various experimental methods and theoretical techniques [5? , 6]. However, the consensus has not been reached due to the difficulties to interpret the experimental data [?] and the inaccuracies in simulation models at the headgroup region [7–9]. Some studies propose specific binding of calcium directly to PS lipid headgroups [?], while others propose that only the local concentration of cations in the vicinity of negatively charged membranes containing PS lipids and that the binding constant of cations is similar to zwitterionic and negatively charged lipids [10?]. The NMR data proposes that the PS headgroup is more rigid than PC, PE or PG headgroups, but more detailed interpretation has not been done.

Headgroup and glycerol backbone C-H bond order parameters calculated from MD simulations have been recently used to interpret the lipid structures in NMR experiments and to validate lipid structure and ion binding in simulations of PC lipid bilayers [7–9, 11]. In this work we apply this approach to PS lipid headgroup in order to elucidate the structural details and ion binding to negatively charged lipids. The results are expected to elucidate also PS mediated signalling events because glycerol backbone and headgroup structure and behaviour are similar in model membranes and in bacteria [10, 12, 13].

PC lipid headgroup order parameters can be used to measure ion binding affinity, because their magnitude is proportional to the amount of bound charge in bilayer [8, 14]. The molecular electrometer concept can be used also for bilayers con-




FIG. 1: Chemical structures and labels for the headgroup carbons.

taining PC lipids mixed with charged lipids [5, 15, 16]. This is demonstrated in Fig. 14, showing the changes of PC headgroup order parameters as a function of CaCl_2 concentration in the presence of different amounts of negatively charged PS or PG lipids.

The decrease of order parameters with CaCl_2 is more pronounced for systems with more negatively charged lipids. Order parameters reach the values of pure PC bilayer close to CaCl_2 concentrations of $\sim 50\text{--}300\text{mM}$. At this point the Ca^{2+} binding presumably fully cancels the charge from negative lipids and overcharging occurs above these concentrations. The interpretation of this data and some other results has been that [10]

“(i) Ca^{2+} binds to neutral lipids (phosphatidylcholine, phosphatidylethanolamine) and negatively charged lipids (phosphatidylglycerol) with approximately the same binding constant of $K = 10\text{--}20\text{ M}^{-1}$;

(ii) the free Ca^{2+} concentration at the membrane interface is distinctly enhanced if the membrane carries a negative surface charge, either due to protein or to lipid;

(iii) increased inter-facial Ca^{2+} also means increased amounts of bound Ca^{2+} at neutral and charged lipids;

(iv) the actual binding step can be described by a Langmuir adsorption isotherm with a 1 lipid:1 Ca^{2+} stoichiometry, provided the interfacial concentration C_M , is used to describe the chemical binding equilibrium.”

METHODS

Solid state NMR experiments

The experimental protocol is the same used in Ref. 11.

1. Some basic details should be given.

Molecular dynamics simulations

Molecular dynamics simulation data was collected with Open Collaboration method. The simulated systems are listed in Table I. The simulation details are in SI or in references in the table.

RESULTS AND DISCUSSION

Headgroup and glycerol backbone order parameters measured from POPS lipid bilayer

Figs. 2 and 3 summarize the experimental NMR results for POPS bilayer sample. 13. New figure, probably combining Figs. 2 and 3, should be done. This discussion to be finished after this.

The headgroup and glycerol backbone order parameters of POPS measured in this work are compared to the literature values of DOPS [37] and POPC [38] in Fig. 4. Our results for POPS are in good agreement with the previously reported values for DOPS measured with ^2H NMR. Significant differences are observed between PC and PS lipids, especially at the headgroup region. Previous discussions in the literature have concluded that glycerol backbone structure is largely similar in PC, PE, PG and PS lipids [12]. The headgroup region was found to be similar in PC, PE and PG lipid (assuming that the signs of PE and PG order parameters are the same as in PC), while the PS headgroup was suggested to be more rigid [39, 40]. The detailed structural differences between the headgroups is, however, not known.

Headgroup and glycerol backbone in simulations of PS lipid bilayers without additional ions

The headgroup order parameters of DOPS and POPS bilayers from different simulation models are compared with the experimental data in Fig. 5. In line with the previous study for PC lipids [7], the glycerol backbone order parameters in CHARMM36 and Berger force fields roughly agree with the experimental data, but significant discrepancies are observed




FIG. 2: (a) The headgroup region of the INEPT spectrum where alpha and beta are identified. (b) The R-PDLF slices for alpha and beta showing one single splitting for beta (which gives an order parameter equal to 0.12), and for alpha a superposition of a large splitting (order parameter equal to 0.09) and a very small splitting which cannot be resolved with the available resolution. (c) Points with error bars are the experimental S-DROSS data. The thick lines are SIMPSON simulations. The S-DROSS slice for beta clearly shows that the order parameter is negative, which is confirmed by SIMPSON simulations using the order parameter value of -0.12. The S-DROSS slice for alpha suggests that the higher order parameter is positive and the deviation towards negative values in the longer t_1 times suggests that the smaller order parameter is negative. This is confirmed by SIMPSON simulation using value of 0.09 for the larger alpha order parameter and the value of -0.02 for smaller (black curve). The value for the smaller alpha order parameter for SIMPSON calculation was taken from Fig 3 in Ref. 36, because resolution in ^{13}C NMR experiments was not high enough to determine numerical value for this. The S-DROSS curve from SIMPSON simulation with positive value for the smaller order parameter gave did not agree with experiments (dashed grey), confirming the interpretation that the smaller order parameter is negative.

14. Maybe we should combine this with 3

for Slipids and CKP models. The glycerol backbone struc-

TABLE I: List of MD simulations. The salt concentrations calculated as $[\text{salt}] = N_c \times [\text{water}] / N_w$, where $[\text{water}] = 55.5$ M. CKP1 refers to the version with Berger/chiu NH3 charges compatible with Berger and CKP2 to the version with more Gromos compatible version.

lipid/counter-ions	force field for lipids / ions	NaCl (mM)	CaCl ₂ (mM)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
DOPS/Na ⁺	CHARMM36 [?]] 2.	0	0	128	4480	0	303	500	100	[17]
DOPS/Na ⁺	CHARMM36ua [?]] 3.	0	0	128	4480	0	303	500	100	[18]
DOPS/Na ⁺	Slipids [19]	0	0	128	4480	0	303	500	100	[20]
DOPS/Na ⁺	Slipids [19]	0	0	288	11232	0	303	200	100	[21]
DOPS/Na ⁺	Berger [22]	0	0	128	4480	0	303	500	100	[23]
DOPS/Na ⁺	GROMOS-CKP1 [?]] 4.	0	0	128	4480	0	303	500	100	[24]
DOPS/Na ⁺	GROMOS-CKP2 [?]] 5.	0	0	128	4480	0	303	500	100	[25]
DOPS/Na ⁺	lipid17 [?]]	0	0	?	?	?	?	?	?	[?]]
POPS/Na ⁺	CHARMM36 [?]] 6.	0	0	128	4480	0	298	500	100	[26]
POPS/Na ⁺	CHARMM36ua [?]] 7.	0	0	128	4480	0	298	500	100	[27]
POPS/Na ⁺	Slipids [19]	0	0	128	4480	0	298	500	100	[28]
POPS/Na ⁺	Berger [?]]	0	0	128	4480	0	298	500	100	[29]
POPS/Na ⁺	MacRog [30]	0	0	128	5120	0	298	200	? 8.	[31]
POPS/Na ⁺	GROMOS-CKP1 [?]] 9.	0	0	128	4480	0	298	500	100	[32]
POPS/Na ⁺	GROMOS-CKP2 [?]] 10.	0	0	128	4480	0	298	500	100	[33]
POPS/Na ⁺	lipid17 [?]]	0	0	?	?	?	?	?	?	?
POPC:POPS (5:1)/K ⁺	CHARMM36 [34?]] 11.	0	0	110:22	4935	0	298	100	100 12.	[35]

^aNumber of lipid molecules with largest mole fraction

^bNumber of water molecules

^cNumber of additional cations

^dSimulation temperature

^eTotal simulation time

^fTime used for analysis

^gReference for simulation files

tures from CHARMM36 and Slipids simulations visualized in Fig. 7 reveal the missing structures in Slipids model, which probably lead to the incorrect order parameters. For the headgroup α and β carbon order parameters of PS lipids, the tested models perform less well than for PC headgroup in previous study [7]. None of the tested models give a satisfactory agreement with experiments for the headgroup order parameters of PS lipids. Based on the subjective ranking shown in Fig. 6, the CHARMM36 is the best performing model for both lipids. However, the total deviation from the experiments for the PS lipids is significantly higher for PS lipids (8) than for PC (3) [7]. Therefore the interpretation of structural differences between PC and PS lipids from current MD simulation models is very challenging.

23. Dihedral angle distributions in Fig. 15 should be analyzed from different models and included in the discussion. Maybe also figures similar to 7.

In contrast to PC lipids discussed previously [7, 8], systems with negatively charged lipids always contains counterions. Since the ion binding changes the headgroup order parameters [8, 14], the counterion distribution have to be taken into account also when additional ions are not present. The Na⁺ density profiles from different simulation are shown in Fig. 8.

Headgroup structure in PS and PC mixtures

Lipid mixtures containing PC and PS lipids are present in biological systems. Also the Ca²⁺ binding affinity to bilayers containing PS lipids is measured from mixtures with PC, because pure PS bilayers exhibit instant transition to the ordered phase with the addition of the ions. Therefore, the characterization of mutual interactions between PC and PS lipid headgroups is necessary in order to understand PS lipids in biological environments.

Fig. 9 shows the headgroup order parameters of POPC and POPS lipids in mixtures with various mole fractions from simulations and experiments [5, 13]. The headgroup order parameters of POPC increase with the increasing amount of negatively charged POPS, as expected from the the electrometer concept [14]. Notably, the concept seems to be generally valid, at least qualitatively, also for other lipid mixed with PC lipids; the PC lipid headgroup order parameters increase when mixed with negatively charged lipids (PS, PI, CL, PA and PG) and remains almost unchanged when mixed with neutral lipids (PE and SM) [13] or cholesterol [38]. This is, however, not reproduced in the simulation data in Fig. 9. The order parameter for β -carbon of POPC decreases and almost negligible increase is seen in α order parameter with the addition of PS lipid in CHARMM36 simulation. This may indicate incorrect




FIG. 3: R-PDFL slices for glycerol backbone carbons.

15. We need nicer figure for this data. Maybe combine with 2

16. What are the top figures actually?

interactions between PC and PS headgroups in simulations. On the other hand, the previously reported overestimated ion binding affinity may also play a role, as the amount of counter-




FIG. 4: Headgroup and glycerol backbone order parameters of POPS measured in this work compared with values for DOPS (^2H NMR, 0.1M of NaCl) [37] and POPC (^{13}C NMR) [38] from literature. Signs for PS order parameters as measured in this work and signs for PC as measured in Refs [11?].

rions is increasing with PS. **24. Data from other force fields to be collected before discussion can be finalized.**

Also the headgroup order parameters of POPS mixed with varying amounts of POPC from simulations and experiments [5, 15] are shown in Fig. 9. The β -carbon order parameter of POPS slightly decreases and positive order parameter of α -carbon slightly increases with increasing amount of PS lipids. This may indicate increasing order of the head-group. It should be, however, noted that the experimental data for pure POPS and POPC/POPS mixture come from different experimental sets, ^{13}C NMR in this work and ^2H NMR from Ref. 5, respectively. Therefore the accuracy of the order parameter change is not as high as typically in the measurements of order parameter changes, see discussion about qualitative and quantitative accuracy in Ref. 9. The changes of PS headgroup order parameters are not reproduced by the tested simulation models. The β -carbon order parameter increase with increasing amount of PS in CHARMM36 and MacRog simulations in contrast to the experimental data. The smaller α -carbon order parameter increase in both simulation models with increasing amount of PS, while it is almost unchanged in experiments. The larger α -carbon order parameter increase in MacRog and decrease in CHARMM36 with increasing amount of PS, both model exhibiting a poor agreement with experiments. Significant improvement in the MD simulation models are needed to interpret the PS headgroup structures and mutual interactions with PC lipids.

Ca^{2+} binding affinity in bilayers with negatively charged PS lipids

Before using headgroup order parameters to compare ion binding affinity between simulations and experiments, the re-




FIG. 5: Order parameters for PS headgroup and glycerol backbone from simulations with different models and experiments without CaCl_2 . Experimental data from [37] contains 0.1M of NaCl. Signs are taken from experiments for POPS described in Supplementary Information. The vertical bars shown for the computational values are not error bars, but demonstrate that for these systems we had at least two data sets; the ends of the bars mark the extreme values from the sets, and the dot marks their measurement-time-weighted average.

17. Check and report all the counterions. There may be some simulation data sets with both Na+ and K+ ions?
18. CHARM36-UA point for alpha in wrong place?
19. MacRog POPS data missing. The equilibration yet to be checked, thought.
20. In table I: "CKP1 refers to the version with Berger/chiu NH3 charges compatible with Berger and CKP2 to the version with more Gromos compatible version." Is this correct also in this figure?
21. Lipid17 results should be added.
22. We should also add CHARM36 POPS ran with K counterions:
<https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/1#issuecomment-308570874>

	β	α	g_3	g_2	g_1	Σ
CHARMM 36	M	M F	M	M	M F	8
CHARMM 36-UA	M	M	M	M	M F	8
GROMOS-CKP1	M	M F	M F	M	M F	14
GROMOS-CKP2	M	M F	M F	M	M F	14
Slipid	M	M	M F	M	M F	14
Berger	M	M F	M F	M	M F	15

FIG. 6: Rough subjective ranking of force fields based on Figure 5. Here M indicates a magnitude problem, F a forking problem; letter size increases with problem severity. Color scheme: within experimental error (dark green), almost within experimental error (light green), clear deviation from experiments (light red), and major deviation from experiments (dark red). The Σ -column shows the total deviation of the force field, when individual carbons are given weights of 0 (matches experiment), 1, 2, and 4 (major deviation). For full details of the assessment, see Supplementary Information.




FIG. 7: Snapshots overlaid from different simulations for glycerol backbone region by Pavel Buslaev.

sponse of the order parameters to bound charge has to be quantified. The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [41]. In Fig. 10. The figure shows that the order parameters are too sensitive to bound charge in Lipid14 model, while CHARM36 is in better agreement with experiments. This has to be taken into account when analysing the binding affinities.

The headgroup order parameter changes of POPC and POPS from POPC:POPS (5:1) bilayers as a function of Ca^{2+} concentration from different simulations and experiments [5] are shown in Fig. 11. As shown previously, the changes of PC headgroup order parameters can be used to measure the amount of bound ions to the bilayer according to the electrometer concept [8, 14, 42]. This would suggest that the




FIG. 8: Sodium density of POPS lipid bilayer along the membrane normal from simulations with different force fields.

Ca²⁺ binding is overestimated in MacRog simulation and underestimated in CHARMM36 simulation. It should be however noted, that MacRog simulation with low concentrations gives a good agreement with experiments. The result from CHARMM simulation is somewhat surprising because previous study suggested that Ca²⁺ ions overbind to pure PC lipid bilayers [8]. 27.The role of counterions and equilibration should be somehow checked and discussed. The density profiles of Ca²⁺ from the simulations are shown in Fig. 12. Overall, it seems that calcium clearly overbinds in MacRog simulations as expected based on previous results [8]. However, the calcium binding in CHARMM36 simulations seems to be surprisingly low. Thi

Also dependence of β -carbon of PG on CaCl₂ concentration is compared with experiments [15] in Fig. ?? . Absolute value of the order parameter is too large without ions, but rapid decrease due to addition of CaCl₂ is observed in agreement with experiments for systems with 1:1 mixture of POPC and POPG. In addition, absolute value in systems with CaCl₂

is in agreement with experiments. However, system with 4:1 mixture of POPC and POPG behaves differently, but experimental data is not available for comparison for this mixture.

32.More simulation data for systems with PS lipids and CaCl₂ would be probably useful

Also the experimental order parameters of POPS headgroup as a function of CaCl₂ concentration are shown in Fig. ?? . The order parameter changes with added CaCl₂ are overestimated in both simulation models. However, the changes can be considered qualitatively correct for both segments in CHARMM36 model and in β segment in MacRog model.

33.These should be compared to simulations for potential structural interpretation of the changes.

CONCLUSIONS




FIG. 9: Headgroup order parameters from PC:PS mixtures from different simulation models and experiments. Left panel shows the PC headgroup order parameters (experimental results from Ref. 13, signs are determined as discussed in [7, 9]). Right panel shows PS headgroup order parameters (experimental result for pure POPS measured in this work at 298K, experimental result for mixture from Ref. 5 at 298K). Counterions in experiments are sodium, while potassium is used in simulations. Using sodium in simulations do not have a significant effect.

25. Simulation of CHARMM36 at 298K should be maybe rerun with Gromacs 5.

26. We need results also from other than CHARMM36 force field.

SUPPLEMENTARY INFORMATION

Simulated systems

PC lipid headgroup response to different mixtures in experiments

As shown in Fig. 13, order parameters of PC headgroup behave in various lipid mixtures as expected from the electrometer concept [13, 14], i.e., order parameters increase when anionic lipids are mixed with PC and decrease with cationic surfactants. The changes with the addition of neutral lipids is significantly smaller.

Cation binding affinity to lipid bilayers with different amount of charge

Dihedrals

Details of the rough subjective force field ranking (Fig. 6)

The assessment was based fully on the Fig. 5. First, for each carbon (the columns in Fig. 5) in each force field (the rows), we looked separately at deviations in magnitude and forking.

Magnitude deviations, i.e., how close to the experimentally obtained C-H order parameters (OPs) the force-field-produced OPs were. For each carbon, the following 5-step scale was used:

0 (): More than half of all the calculated OPs (that is, of all different hydrogens in all different lipids) were within the *subjective sweet spots* (SSP, blue-shaded areas in Fig. 5).

1 (m): All the calculated OPs were < 0.03 units away from the SSP.

2 (M): All the calculated OPs were < 0.05 units away from the SSP.

3 (M): All the calculated OPs were < 0.10 units away




FIG. 10: The response of headgroup order parameters to the fixed amount of cationic surfactants in POPC bilayer is compared between simulations and experiments [41].

from the SSP.

4 (M): Some of the calculated OPs were > 0.10 units away from the SSP.

Forking deviations, i.e., how well the difference in order parameters of two hydrogens attached to a given carbon matched that obtained experimentally. Note that this is not relevant for β and g_2 , which have only one hydrogen. For the α carbon, for which a considerable forking of 0.105 is experimentally seen, the following 5-step scale was used:

0 (): The distance D between the dots (that mark the measurement-time-weighted averages in Fig. 5) was $0.08 < D < 0.13$ units for all the calculated OPs (that is, for all different lipids).

1 (F): $(0.06 < D < 0.08)$ OR $(0.13 < D < 0.15)$.

2 (F): $(0.04 < D < 0.06)$ OR $(0.15 < D < 0.17)$.

3 (F): $(0.02 < D < 0.04)$ OR $(0.17 < D < 0.19)$.

4 (F): $(D < 0.02)$ OR $(0.19 < D)$.

For the g_3 carbon, for which no forking is indicated by experiments, the following 5-step scale was used:

0 (): $D < 0.02$.

1 (F): $0.02 < D < 0.04$.

2 (F): $0.04 < D < 0.06$.

3 (F): $0.06 < D < 0.08$.

4 (F): $0.08 < D$.

For the g_1 carbon, for which a considerable forking of 0.13 is experimentally seen, the following 5-step scale was used:

0 (): $0.11 < D < 0.15$.

1 (F): $(0.09 < D < 0.11)$ OR $(0.15 < D < 0.17)$.

2 (F): $(0.07 < D < 0.09)$ OR $(0.17 < D < 0.19)$.

3 (F): $(0.05 < D < 0.07)$ OR $(0.19 < D < 0.21)$.

4 (F): $(D < 0.05)$ OR $(0.21 < D)$.

Based on these assessments of magnitude and forking deviations, each carbon was then assigned to one of the following groups: "within experimental error" (magnitude and forking deviations both on step 0 of the scales described above), "almost within experimental error" (sum of the magnitude and forking deviation steps 1 or 2), "clear deviation from experiments" (sum of magnitude and forking deviation steps from 3 to 5), and "major deviation from experiments" (sum of magnitude and forking deviation steps from 6 to 8). These groups are indicated by colors in Fig. 4. (Note that for β and g_2 , for which there can be no forking, the corresponding group assignment limits were: 0, 1, 2, and 3.)

Finally, the total ability of the force field to describe the headgroup and glycerol structure was estimated. To this end, the groups were given the following weights: 0 (within experimental error), 1 (almost within experimental error), 2 (clear deviation from experiments), 4 (major deviation from experiments), and the weights of the five carbons were summed up. The sum, given in the Σ -column of Fig. 5, was then used to (roughly and subjectively, as should be clear from the above description) rank the force fields.




FIG. 11: POPC (left) and POPS (right) headgroup order parameters in lipid bilayer with POPC:POPS (5:1) mixture as a function CaCl_2 concentration from experiments [5] and different simulations.

* samuli.ollila@helsinki.fi

- [1] M. A. Lemmon, Nat. Rev. Mol. Cell Biol. **9**, 99 (2008).
- [2] P. A. Leventis and S. Grinstein, Annual Review of Biophysics **39**, 407 (2010).
- [3] L. Li, X. Shi, X. Guo, H. Li, and C. Xu, Trends in Biochemical Sciences **39**, 130 (2014), ISSN 0968-0004.
- [4] T. Yeung, G. E. Gilbert, J. Shi, J. Silvius, A. Kapus, and S. Grinstein, Science **319**, 210 (2008).
- [5] M. Roux and M. Bloom, Biochemistry **29**, 7077 (1990).
- [6] A. Melcrová, S. Pokorna, S. Pullanchery, M. Kohagen, P. Jurkiewicz, M. Hof, P. Jungwirth, P. S. Cremer, and L. Cwiklik, Sci. Reports **6**, 38035 (2016).
- [7] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduč, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., J. Phys. Chem. B **119**, 15075 (2015).
- [8] A. Catte, M. Girych, M. Javanainen, C. Loison, J. Melcr, M. S. Miettinen, L. Monticelli, J. Maatta, V. S. Oganessian, O. H. S. Ollila, et al., Phys. Chem. Chem. Phys. **18**, 32560 (2016).
- [9] O. S. Ollila and G. Pabst, Biochimica et Biophysica Acta (BBA) - Biomembranes **1858**, 2512 (2016).
- [10] J. Seelig, Cell Biology International Reports **14**, 353 (1990), ISSN 0309-1651, URL <http://www.sciencedirect.com/science/article/pii/030916519091204H>.
- [11] T. M. Ferreira, R. Sood, R. Bärenwald, G. Carlström, D. Topgaard, K. Saalwächter, P. K. J. Kinnunen, and O. H. S. Ollila, Langmuir **32**, 6524 (2016).
- [12] H. U. Gally, G. Pluschke, P. Overath, and J. Seelig, Biochemistry **20**, 1826 (1981).
- [13] P. Scherer and J. Seelig, EMBO J. **6** (1987).
- [14] J. Seelig, P. M. MacDonald, and P. G. Scherer, Biochemistry **26**, 7535 (1987).
- [15] F. Borle and J. Seelig, Chemistry and Physics of Lipids **36**, 263 (1985).
- [16] P. M. Macdonald and J. Seelig, Biochemistry **26**, 1231 (1987).
- [17] T. Piggot, CHARMM36 DOPS simulations (versions 1 and 2) 303 K 1.0 nm LJ switching (2017), URL <https://doi.org/10.5281/zenodo.1129411>.
- [18] T. Piggot, CHARMM36-UA DOPS simulations (versions 1 and 2) 303 K 1.0 nm LJ switching (2017), URL <https://doi.org/10.5281/zenodo.1129456>.
- [19] J. P. M. Jämbeck and A. P. Lyubartsev, Phys. Chem. Chem. Phys. **15**, 4677 (2013).
- [20] T. Piggot, Slipids DOPS simulations (versions 1 and 2) 303 K 1.0 nm cut-off with LJ-PME (2017), URL <https://doi.org/10.5281/zenodo.1129411>.




FIG. 12: Ca²⁺ density profiles from simulations.

28. Ca²⁺ binding in CHARMM is lower than to neutral membrane. The reason should be found out (probably equilibration). 29. These are now mass densities, numbers would be probably better. 30. Counterions should be also included. 31. Not all the data from MacRog is included.

org/10.5281/zenodo.1129439.

- [21] F. Favela-Rosales, *MD simulation trajectory of a fully hydrated DOPS bilayer: SLIPIDS, Gromacs 5.0.4*. 2017. (2017), URL <https://doi.org/10.5281/zenodo.495510>.
- [22] P. Mukhopadhyay, L. Monticelli, and D. P. Tieleman, *Biophysical Journal* **86**, 1601 (2004).
- [23] T. Piggot, *Berger DOPS simulations (versions 1 and 2) 303 K 1.0 nm cut-off* (2017), URL <https://doi.org/10.5281/zenodo.1129419>.
- [24] T. Piggot, *GROMOS-CKP DOPS simulations (versions 1 and 2) 303 K with Berger/Chiu NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129429>.
- [25] T. Piggot, *GROMOS-CKP DOPS simulations (versions 1 and 2) 303 K with GROMOS NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129447>.
- [26] T. Piggot, *CHARMM36 POPS simulations (versions 1 and 2)*




FIG. 13: PC headgroup order parameters from experiments of mixtures with PE, PS, PG and cholesterol [13, 38, 41]. Signs are determined as discussed in [7, 9].

298 K 1.0 nm LJ switching (2017), URL <https://doi.org/10.5281/zenodo.1129415>.

- [27] T. Piggot, *CHARMM36-UA POPS simulations (versions 1 and 2) 298 K 1.0 nm LJ switching* (2017), URL <https://doi.org/10.5281/zenodo.1129458>.
- [28] T. Piggot, *Slipids POPS simulations (versions 1 and 2) 298 K 1.0 nm cut-off with LJ-PME* (2017), URL <https://doi.org/10.5281/zenodo.1129441>.
- [29] T. Piggot, *Berger POPS simulations (versions 1 and 2) 298 K 1.0 nm cut-off* (2017), URL <https://doi.org/10.5281/zenodo.1129425>.
- [30] A. Maciejewski, M. Pasenkiewicz-Gierula, O. Cramariuc, I. Vattulainen, and T. Róg, *J. Phys. Chem. B* **118**, 4571 (2014).
- [31] M. Javanainen, *Simulation of a pops bilayer* (2017), URL <https://doi.org/10.5281/zenodo.1120287>.
- [32] T. Piggot, *GROMOS-CKP POPS simulations (versions 1 and 2) 298 K with Berger/Chiu NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129431>.
- [33] T. Piggot, *GROMOS-CKP POPS simulations (versions 1 and 2) 298 K with GROMOS NH3 charges and PME* (2017), URL <https://doi.org/10.5281/zenodo.1129435>.
- [34] J. B. Klauda, R. M. Venable, J. A. Freites, J. W. O'Connor, D. J. Tobias, C. Mondragon-Ramirez, I. Vorobyov, A. D. MacKerell Jr, and R. W. Pastor, *J. Phys. Chem. B* **114**, 7830 (2010).
- [35] O. H. S. Ollila, *POPS+83%popc lipid bilayer simulation at T298K ran CHARMM_GUI force field and Gromacs*




FIG. 14: The change of PC headgroup order parameters in the presence of different amount of negatively charged lipids respect to the values without added CaCl_2 . The original data is the same as in Fig. ?? . Pure DPPC data from [42], pure POPC data from [43], POPC:POPS mixture data from [5], POPC:POPG mixture data with 0.1M NaCl from [16] and POPC:POPG mixture data without NaCl from [15].

- (2017), URL <https://doi.org/10.5281/zenodo.1011104>.
- [36] M. Roux and M. Bloom, *Biophys. J.* **60**, 38 (1991).
- [37] J. L. Browning and J. Seelig, *Biochemistry* **19**, 1262 (1980).
- [38] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
- [39] R. Wohlgemuth, N. Waespe-Sarcevic, and J. Seelig, *Biochemistry* **19**, 3315 (1980).
- [40] G. Büldt and R. Wohlgemuth, *The Journal of Membrane Biology* **58**, 81 (1981), ISSN 1432-1424, URL <http://dx.doi.org/10.1007/BF01870972>.
- [41] P. G. Scherer and J. Seelig, *Biochemistry* **28**, 7720 (1989).
- [42] H. Akutsu and J. Seelig, *Biochemistry* **20**, 7366 (1981).
- [43] C. Altenbach and J. Seelig, *Biochemistry* **23**, 3913 (1984).

ToDo

1. Some basic details should be given. 2
13. New figure, probably combining Figs. 2 and 3, should be done. This discussion to be finished after this. 2
14. Maybe we should combine this with 3 2
2. Correct citation for CHARMM DOPS 3
3. Correct citation for CHARMMua DOPS 3
4. Correct citation(s) for CKP. 3
5. Correct citation(s) for CKP. 3
6. Correct citation for CHARMM POPS 3
7. Correct citation for CHARMMua DOPS 3
8. Equilibration? 3
9. Correct citation(s) for CKP. 3
10. Correct citation(s) for CKP. 3
11. Correct citation for CHARMM POPS 3
12. Equilibration? 3
23. Dihedral angle distributions in Fig. 15 should be analyzed from different models and included in the discussion. Maybe also figures similar to 7. 3
15. We need nicer figure for this data. Maybe combine with 2 4
16. What are the top figures actually? 4
24. Data from other force fields to be collected before discussion can be finalized. 4
17. Check and report all the counterions. There may be some simulation data sets with both Na^+ and K^+ ions? 5
18. CHARMM36-UA point for alpha in wrong place? 5
19. MacRog POPS data missing. The equilibration yet to be checked, thought. 5
20. In table I: "CKP1 refers to the version with Berger/chiu NH_3 charges compatible with Berger and CKP2 to the version with more Gromos compatible version." Is this correct also in this figure? 5
21. Lipid17 results should be added. 5
22. We should also add CHARMM36 POPS ran with K^+ counterions: <https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/1#issue308570874> 5
27. The role of counterions and equilibration should be somehow checked and discussed. 6
32. More simulation data for systems with PS lipids and CaCl_2 would be probably useful 6
33. These should be compared to simulations for potential structural interpretation of the changes. 6
25. Simulation of CHARMM36 at 298K should be maybe rerun with Gromacs 5. 7
26. We need results also from other than CHARMM36 force field. 7
28. Ca^{2+} binding in CHARMM is lower than to neutral membrane. The reason should be found out (probably equilibration). 10
29. These are now mass densities, numbers would be probably better. 10




FIG. 15: Experimental results for sign measurement for POPS sample

30. Counterions should be also included. 10 31. Not all the data from MacRog is included. 10