

# NMRLipids IV: Headgroup & glycerol backbone structures, and cation binding in bilayers with PE, PG and PS lipids

O. H. Samuli Ollila<sup>1,2, \*</sup>

<sup>1</sup>*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic*

<sup>2</sup>*Institute of Biotechnology, University of Helsinki*

(Dated: August 11, 2017)

Primarily measured but also simulated NMR order parameters will be collected also for other than phosphatidylcholine (these are discussed in NMRLipids I) headgroup. The information will be used to understand structural differences between different lipid molecules in bilayers.

## INTRODUCTION

In NMRLipids I and II project we were looking for a MD model which would correctly reproduce headgroup and glycerol backbone structures and cation binding for PC lipid bilayers [1, 2]. Here we extend the same goal for other than PC lipids. Currently the focus is on PE, PG and PS bilayers and their mixtures with PC. Experimental data with different amounts of added salt is now collected and presented in this manuscript. Simulations for bilayers containing PG and PS with low salt conditions are ran with parameters from CHARMM GUI and compared to experiments. Some order parameters from simulations seems to be off from experimental values. Simulation data from other models would be highly useful to see if some of the existing models would reproduce the experimental order parameters and to analyze different conformations predicted by different models respect to experiments.

## EXPERIMENTAL GLYCEROL BACKBONE AND HEADGROUP ORDER PARAMETERS FOR PE, PG AND PS LIPIDS

Absolute values of experimental order parameters for different lipid headgroups are collected in Fig. 1. Signs are measured only for PC as far as I know, thus only absolute values are used for now.

Based on superficial reading, the conclusions in the literature are roughly

- 1) glycerol backbone structures are largely similar irrespectively of the headgroup [8],
  - 2) glycerol backbone and headgroup structure and behaviour are similar in model membranes and in bacteria [8–10],
  - 3) headgroup structures are similar in PC, PE and PG lipids, while headgroup is more rigid in PS lipids [6, 11].
- Extensive discussion about structural details of PE, PG or PS headgroups do not exists (as far as I know), In contrast to PC lipids (see [1] and references therein).




FIG. 1: Absolute values of order parameters for headgroup and glycerol backbone with different headgroups from experiments. POPC values are from [3], DOPS from [4] contains 0.1M of NaCl, POPG from [5] contains 10nM PIPES, DPPG from [6] contains 10mM PIPES and 100mM NaCl, DPPE from [7], E.coliPE and E.coliPG are from [8].

## GLYCEROL BACKBONE AND HEADGROUP ORDER PARAMETERS FOR PE, PG AND PS LIPIDS IN SIMULATIONS

Several simulations containing PE, PG and PS lipids have been published [? ], 1.List should be completed however, glycerol backbone and headgroup order parameters are not compared to the experiments (based on superficial reading of literature).

### PE headgroup

Absolute values of order parameters between Slipids simulations [? ] and experiments for DPPE are shown in Fig. 2. Glycerol backbone order parameters in Slipids are off from experiments, as systematically observed in this and previous studies [? ]. Alpha order parameter is too low, but beta shows apparent agreement with experiments. However, the sign of beta order parameter is positive, in contrast to PC where negative sign was measured. Thus, the the beta order parameter agrees with experiment with the assumption that its sign is




FIG. 2: Order parameters for DPPE headgroup and glycerol backbone from simulations with Slipids [?] and experiments (DPPE from [7] and E.coliPE from [8]). Absolute values are shown, because signs are not known experimentally.

2.Experimental signs of the order parameters would be highly useful.

opposite than for PC.

#### PS headgroup

Experimental order parameters for bilayers with PS headgroups are available in literature only for DOPC.

Comparison between experiments and simulations with CHARMM GUI parameters for PS lipids in Fig. 4 suggest that the model overestimates magnitudes in  $\beta$  and  $g_3$  order parameters and forking in  $\alpha$ .

#### PG headgroup

Comparison between experiments and simulations with CHARMM GUI parameters for PG lipids in Fig. 5 suggest that the model overestimates forking in  $\gamma$  and  $\alpha$  order parameters, and magnitudes of  $\beta$  and  $g_1$  order parameters.

#### CA<sup>2+</sup> BINDING IN BILAYERS WITH NEGATIVELY CHARGED PG AND PS LIPIDS

PC lipid headgroup order parameters can be used to measure ion binding affinity, because their magnitude is linearly proportional to the amount of bound charge in bilayer [2, 13]. This molecular electrometer concept can be used also for bilayers containing PC lipids mixed with charged lipids [5, 12, 14]. This is demonstrated in Figs 6, 7 and 8, showing order parameters for PC headgroup  $\alpha$  and  $\beta$  carbons as a function of CaCl<sub>2</sub> concentration in the presence of different amounts of negatively charged PS or PG lipids.




FIG. 3: Order parameters for DOPS headgroup and glycerol backbone from simulations with different models and experiments without CaCl<sub>2</sub>. Experimental data from [4] contains 0.1M of NaCl. Signs are not known experimentally. They chosen to give best agreement simulations.

3.Experimental signs of the order parameters would be highly useful.

PC headgroup order parameters increase when negatively charged PS or PG are added to PC bilayer in the absence of added CaCl<sub>2</sub>, as expected based on electrometer concept [13] (see Fig. 7). Further, the order parameters decrease with the addition of CaCl<sub>2</sub> and the decrease is more pronounced for systems with more negatively charged lipids (see Fig. 8). At CaCl<sub>2</sub> concentrations (~ 50-300mM) where order parameters reach the values for pure PC, the Ca<sup>2+</sup> binding presumably fully cancels the charge from negative lipids and overcharging occurs above these concentrations. The interpretation of this data and some other results has been that [10]

- ”(i) Ca<sup>2+</sup> binds to neutral lipids (phosphatidylcholine, phosphatidylethanolamine) and negatively charged lipids (phosphatidylglycerol) with approximately the same binding constant of  $K = 10\text{-}20\text{ M}^{-1}$ ;
- (ii) the free Ca<sup>2+</sup> concentration at the membrane




FIG. 4: Order parameters for POPS (top) and DOPS (bottom) headgroup and glycerol backbone from simulations with different models and experiments without  $\text{CaCl}_2$  (DOPS from [4] contains 0.1M of NaCl, POPC:POPS mixture from [12]). Absolute values are shown for experimental data, because signs are not known. Simulations values are  $-S_{CH}$

4. More simulation data for lipids with different headgroups to be collected

5. Some simulations contain potassium as counterions, while some sodium. All experiments here contain some amount of sodium salt. The best ion concentrations for comparison should be figured out.

6. Experimental signs of the order parameters would highly useful.

7. Why there is difference between CHARMM36 simulation results from POPS:POPC mixture and pure POPS? Discussion in <https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/1>

interface is distinctly enhanced if the membrane carries a negative surface charge, either due to protein or to lipid;

(iii) increased inter-facial  $\text{Ca}^{2+}$  also means increased amounts of bound  $\text{Ca}^{2+}$  at neutral and charged lipids;

(iv) the actual binding step can be described by a Langmuir adsorption isotherm with a 1 lipid:1  $\text{Ca}^{2+}$  stoichiometry, provided the interfacial con-




FIG. 5: Order parameters for PG headgroup and glycerol backbone from simulations with CHARMM GUI parameters and experiments without  $\text{CaCl}_2$  (POPG from [5] contains 10mM of PIPES, DPPG from [6] contains 10mM PIPES and 100mM  $\text{CaCl}_2$ , E.Coli PG results from [8]). Absolute values are shown, because signs are not known for experimental data. Glycerol backbone order parameters are negative in simulations, while  $\alpha$ ,  $\beta$  and smaller  $\gamma$  are positive.

8. More simulation data for lipids with different headgroups to be collected

9. CHARMM GUI simulation contains only counter ions as potassium. All experiments here contain some amount of sodium salt. The best ion concentrations for comparison should be figured out. 10. Experimental signs of the order parameters would highly useful.

centration  $C_M$ , is used to describe the chemical binding equilibrium.”

Also the experimental order parameters for PS and PG headgroups as a function of  $\text{CaCl}_2$  concentration are shown in Fig. 9. 12. These should be compared to simulations for potential structural interpretation of the changes.

## CA<sup>2+</sup> BINDING IN BILAYERS WITH NEGATIVELY CHARGED PG AND PS LIPIDS IN SIMULATIONS

Comparison of  $\text{Ca}^{2+}$  binding in PG between CHARMM36 simulations and experiments [5] is shown in Fig. 10. The decrease of  $\alpha$  order parameter is in agreement with experiments, while decrease of  $\beta$  order parameter is overestimated. The result is very similar to the results with PC in NMRLipids II publication [2]. It should be, however, noted that the  $\beta$ -order parameters are not actually measured for PG, but they are calculated from empirical relation  $\Delta S_\beta = 0.43 \Delta S_\alpha$  [15]. Anyway, the data presented in NMRLipids II project and in Fig. 10 together suggest that Calcium binding is similarly overestimated by CHARMM36 model in pure POPC bilayers and mixtures with POPG. The good agreement of  $\alpha$  carbon would be explained by too weak dependence of its order parameter of bound charge 14. Should we check this against cationic surfactant experiments [17]?




FIG. 6: PC headgroup order parameters as a function of  $\text{CaCl}_2$  concentration from experiments containing charged lipids. Pure DPPC data from [15], pure POPC data from [16], POPC:POPG mixture data from [12], POPC:POPG mixture data with 0.1M NaCl from [14] and POPC:POPG mixture data without NaCl from [5].

11. Check the NaCl concentrations in the samples.

Also dependence of  $\beta$ -carbon of PG on  $\text{CaCl}_2$  concentration is compared with experiments [5] in Fig. 9. Absolute value of the order parameter is too large without ions, but rapid decrease due to addition of  $\text{CaCl}_2$  is observed in agreement with experiments for systems with 1:1 mixture of POPC and POPG. In addition, absolute value in systems with  $\text{CaCl}_2$  is in agreement with experiments. However, system with 4:1 mixture of POPC and POPG behaves differently, but experimental data is not available for comparison for this mixture.

15. More simulation data for systems with negatively charged lipids and  $\text{CaCl}_2$  to be collected

## CONCLUSIONS




FIG. 7: Figure 6 zoomed to smaller concentrations.

## SUPPLEMENTARY INFORMATION

\* samuli.ollila@helsinki.fi

- [1] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduć, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., *J. Phys. Chem. B* **119**, 15075 (2015).
- [2] A. Catte, M. Grych, M. Javanainen, C. Loison, J. Melcr, M. S. Miettinen, L. Monticelli, J. Maatta, V. S. Oganessian, O. H. S. Ollila, et al., *Phys. Chem. Chem. Phys.* **18**, 32560 (2016).
- [3] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
- [4] J. L. Browning and J. Seelig, *Biochemistry* **19**, 1262 (1980).
- [5] F. Borle and J. Seelig, *Chemistry and Physics of Lipids* **36**, 263 (1985).
- [6] R. Wohlgemuth, N. Waespe-Sarcevic, and J. Seelig, *Biochemistry* **19**, 3315 (1980).
- [7] J. Seelig and H. U. Gally, *Biochemistry* **15**, 5199 (1976).
- [8] H. U. Gally, G. Pluschke, P. Overath, and J. Seelig, *Biochemistry* **20**, 1826 (1981).
- [9] P. Scherer and J. Seelig, *EMBO J.* **6** (1987).




FIG. 8: The change of PC headgroup order parameters in the presence of different amount of negatively charged lipids respect to the values without added  $CaCl_2$ . The original data is the same as in Fig. 6.




FIG. 9: PG and PS order parameters a function  $CaCl_2$  concentration taken from [5] and [12], respectively.

13. Get the small concentration data from the inserts

- [10] J. Seelig, Cell Biology International Reports **14**, 353 (1990), ISSN 0309-1651, URL <http://www.sciencedirect.com/science/article/pii/030916519091204H>.
- [11] G. Büldt and R. Wohlgemuth, The Journal of Membrane Biology **58**, 81 (1981), ISSN 1432-1424, URL <http://dx.doi.org/10.1007/BF01870972>.
- [12] M. Roux and M. Bloom, Biochemistry **29**, 7077 (1990).
- [13] J. Seelig, P. M. MacDonald, and P. G. Scherer, Biochemistry **26**, 7535 (1987).
- [14] P. M. Macdonald and J. Seelig, Biochemistry **26**, 1231 (1987).
- [15] H. Akutsu and J. Seelig, Biochemistry **20**, 7366 (1981).
- [16] C. Altenbach and J. Seelig, Biochemistry **23**, 3913 (1984).
- [17] P. G. Scherer and J. Seelig, Biochemistry **28**, 7720 (1989).

## ToDo

- 1. List should be completed . . . . . 1
- 2. Experimental signs of the order parameters would be highly useful. . . . . 2

- 3. Experimental signs of the order parameters would highly useful. . . . . 2
- 4. More simulation data for lipids with different headgroups to be collected . . . . . 3
- 5. Some simulations contain potassium as counterions, while some sodium. All experiments here contain some amount of sodium salt. The best ion concentrations for comparison should be figured out. . . . . 3
- 6. Experimental signs of the order parameters would highly useful. . . . . 3
- 7. Why there is difference between CHARMM36 simulation results from POPS:POPC mixture and pure POPS? Discussion in <https://github.com/NMRLipids/NMRLipidsIVotherHGs/issues/1> 3
- 8. More simulation data for lipids with different headgroups to be collected . . . . . 3
- 9. CHARMM GUI simulation contains only counterions as potassium. All experiments here contain some amount of sodium salt. The best ion concentrations for comparison should be figured out. . . . . 3




FIG. 10: PG order parameters as a function  $CaCl_2$  concentration from experiments [5] and CHARMM36 simulations. Note that beta order parameter is calculated from empirical relation  $\Delta S_{\beta} = 0.43\Delta S_{\alpha}$  [15], not actually measured.

- 10. Experimental signs of the order parameters would highly useful. . . . . 3
- 12. These should be compared to simulations for potential structural interpretation of the changes. . . . . 3
- 14. Should we check this against cationic surfactant experiments [17]? . . . . . 3
- 11. Check the NaCl concentrations in the samples. . . . . 4
- 15. More simulation data for systems with negatively charged lipids and  $CaCl_2$  to be collected . . . . . 4
- 13. Get the small concentration data from the inserts . . . . . 5