

NMRlipids III: Lipid–Cholesterol Interactions in Atomistic Molecular Dynamics Simulations

Fernando Favela-Rosales,[†] Peter Heftberger,[‡] Matti Javanainen,^{*,¶,§} Jesper J.
Madsen,^{||,⊥} Josef Melcr,[¶] Markus Miettinen,[#] O. H. Samuli Ollila,^{*,¶,§} Georg
Pabst,^{‡,®} and Thomas Piggot^{△,∇}

[†]*Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN,
Apartado Postal 14-740, 07000 México D.F., México*

[‡]*Institute of Molecular Biosciences, Biophysics Division, NAWI Graz, University of Graz,
Graz 8010, Austria*

[¶]*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech
Republic, Prague 6, Czech Republic*

[§]*Institute of Biotechnology, University of Helsinki*

^{||}*Department of Global Health, College of Public Health*

[⊥]*University of South Florida*

[#]*MPI*

[®]*BioTechMed-Graz, Graz 8010, Austria*

[△]*School of Chemistry, University of Southampton, Southampton SO17 1BJ, United
Kingdom*

[∇]*Chemical Biological and Radiological Sciences, Defence Science and Technology
Laboratory, Porton Down, Salisbury, Wiltshire SP4 0JQ, United Kingdom*

E-mail: matti.javanainen@helsinki.fi; samuli.ollila@helsinki.fi

Abstract

Cholesterol is a central building block in biomembranes, where it induces orientational order, slows down diffusion, and thereby renders the membrane stiffer. Molecular dynamics simulations have played a crucial role in resolving these effects and the underlying molecular picture, yet it has recently become evident that different simulation models predict quantitatively different behavior. Such limitations are easily neglected, as the field rapidly progresses towards simulations of complex membranes mimicking the *in vivo* conditions. Still, for their validity, it is crucial that the interactions between the fundamental building blocks of biomembranes, such as phospholipids and cholesterol, are modelled accurately. Here, we perform a systematic comparison of the ability of commonly used force fields to describe the structure and dynamics of binary mixtures of phosphatidylcholine and cholesterol. Comparison of lipid bilayer simulations with deuterium order parameters and lateral diffusion coefficients from NMR as well as form factors from X-ray scattering reveals that none of the existing force field outperforms the others across the physiologically relevant cholesterol concentrations. The Amber-compatible force fields show the smallest deviations from experiment with Slipids and Lipid17 models excelling at low and high cholesterol concentrations, respectively. The accompanied X-ray scattering data set measured for this work will foster future force field development and refinement.

1 Introduction

2. Check that the uses of “model” and “force field” are consistent

Cellular membranes contain an incredibly complex mixture of lipid molecules¹ which are unevenly distributed in the membrane plane and across its leaflets.²⁻⁴ A key player driving the lateral heterogeneity is cholesterol (CHOL), which is present at concentrations between ~ 10 mol-% (endoplasmic reticulum) and up to ~ 50 mol-% (plasma membrane).² CHOL has the unique ability to order neighbouring lipids and thus induce the liquid ordered (L_o) phase in model membranes.⁵⁻⁸ In the cellular setting, the interaction between phospholipids and

CHOL is associated with the formation of lipid rafts and nanodomains.^{9,10} This heterogeneity can then further regulate protein distribution¹¹ or conformation¹² *via* the hydrophobic mismatch mechanism,¹³ in addition to the direct modulation of protein function.^{14,15}

While detailed experimental information of the interactions among phospholipids and CHOL is relatively sparse, atomistic resolution molecular dynamics (MD) simulations have been widely applied to provide a detailed view of the lateral organization of lipid bilayers as well as lipid-CHOL interactions.^{8,16,17} However, the trustworthiness of any prediction from an MD simulation relies on the quality of the used model. Fortunately, in addition to constant refinement of existing parameters, atomistic *force fields* have also seen the expansion to other classes of biomolecules, including lipids, sugars, and nucleic acids.^{18,19} The traditional protein force fields CHARMM,²⁰ AMBER,²¹ and OPLS^{22,23} now have growing libraries of compatible lipid molecules, including CHOL, in the forms of CHARMM36,^{24,25} Lipid17/Slipids,²⁶⁻³¹ and the model by Maciejewski and Rog (here “MacRog”),³²⁻³⁵ respectively. Moreover, OPLS3e²³ contains parameters for lipids, but CHOL is currently not included.

Thanks to the compatibility with protein force fields, the lipid models can be used in studies of membrane proteins or protein-membrane interactions in addition to sole lipid membranes. Notably, simulations using CHARMM36, Lipid17, and Slipids are now readily set up using CHARMM-GUI for multiple simulation engines.^{36,37} Despite this ease-of-use, the choice of a force field for an MD simulation can be challenging, as each model comes with its own set of pros and cons. Unfortunately, careful comparisons of force field quality, especially for capturing the features of lipid mixtures, are scarce in the literature. This largely results from the lack of availability of systematic, comprehensive, and quantitative datasets from both experiments and simulations.

We have previously performed such comparisons for the structure of phosphatidylcholine (PC) lipids,³⁸ for PC-ion interactions,³⁹ for phosphatidylserine (PS) lipids,⁴⁰ and for phosphatidylethanolamine (PE) and phosphatidylglycerol (PG) lipids.⁴¹ Here, we extend this work to mixtures of PC and CHOL. We compare the ability of popular force fields to reproduce effects of CHOL

on the structure and dynamics of a PC lipid bilayer, and thereby set guidelines for future efforts to validate intermolecular interactions in binary and more complex systems. Moreover, if some of the studied force fields satisfactorily reproduce experimental behavior at least in some range of CHOL concentrations, the molecular level picture provided by it can be assumed to mimic reality in this concentration regime.

Our earlier work has demonstrated that the structural ensembles of lipids from an MD simulation can be directly compared to those from solid state NMR experiments.^{38–41} This approach based on order parameters is quantitative, has small error estimates, is highly reproducible, and does not suffer from perturbation of probes used in many other techniques. Thus, we compare our simulation results with different force fields to the systematic NMR data set of PC C–H bond order parameters with varying CHOL concentrations.⁴² In addition, we measured form factors for a range of PC–CHOL mixtures using small angle X-ray scattering, and used them to subsequently model the electron density profiles across the lipid bilayer. The form factor and electron density profiles can be compared with the same profiles calculated from the simulation data. Whereas NMR provides data on the conformations of individual lipids, form factors from scattering captures the details of the structure of the entire lipid bilayer.⁴³

One limitation in comparing simulations of lipid mixtures with experimental data is that the various experiments are performed at different concentrations, and thus mimicking all compositions in simulations is tedious. Here, we demonstrate that if the simulations and experiments sample multiple CHOL concentrations, the order parameters and electron density profiles can be readily interpolated to intermediate CHOL concentrations. This enables the comparison of data sets that are not extracted at the exact same CHOL concentrations, as is usually the case, especially if multiple techniques are combined in the comparison.

We also evaluated whether the simulation force fields reproduce the dependence of lateral diffusion coefficients on cholesterol measured using pulsed field gradient (PFG) NMR.^{44,45} This comparison is enabled by the recent theoretical work that allows for the elimination of

finite-size effects in MD simulations and thus a quantitative comparison with experiment.^{46,47} With the structural and dynamic comparisons established, we estimated the quality of each force field at different CHOL concentrations. We then analyzed the molecular organization in the best-performing force field at low and high CHOL range, as we expect them to best reflect this organization in reality.

2 Methods

2.1 X-ray Scattering Experiments

We measured small angle X-ray scattering (SAXS) data on multilamellar vesicles (MLVs) composed of POPC and CHOL with the latter present at 0–50 mol-% with 5 mol-% increments. The data were obtained at the EMBL BioSAXS beamline (Hamburg) using 20 keV photons at $T = 300$ K and analyzed in terms of the SDP-GAP model described in Refs. 48 and 49. All lipids were purchased from Avanti Polar Lipids (Alabaster, AL, USA) as dry powders and were used without any further purification. All other chemicals were obtained in pro analysis quality from Lactan (Graz, Austria). Fully hydrated MLVs were prepared using rapid solvent exchange as detailed previously.^{50,51} This avoids the precipitation of cholesterol crystallites at high concentration⁵² yielding non-phase separated samples up to 50 mol-% cholesterol content.

The data from MLVs contain the structure factor (the crystalline lattice) and form factor in a convoluted fashion, yet by fitting the scattered intensity data we obtain both contributions. In this work, only the form factors are discussed. The electron density profiles have been modelled in terms of the SDP model,^{53–55} *i.e.* volume distribution functions are modelled by individual Gaussians or error functions. CHOL is particularly challenging to describe within the SDP model, and therefore form factors are often compared between simulation and experiment⁵⁶ instead of density profiles, or CHOL density is modeled together with that of CH₂ groups.⁵⁷ Here, cholesterol is accounted for by two Gaussians, as proposed

by Jianjun Pan (private communication), and used in Refs. 51 and 58.

2.2 Molecular Dynamics Simulations

We performed MD simulations of a pure POPC membrane as well as five POPC/CHOL mixtures with CHOL content ranging from 11 to 47 mol-%. In order to eliminate the finite size effects due to periodic boundary conditions from lateral diffusion coefficients of lipids, we performed all simulations in three sizes. The small (64 POPC molecules) and medium (256 POPC molecules) systems were only used in the diffusion analysis, whereas all other analyses were performed on the large membranes (1024 POPC molecules). The number of POPC molecules was kept constant across the different CHOL concentrations. All membranes were solvated by 50 waters per lipid (POPC or CHOL). The simulations were performed using four commonly used force fields, namely CHARMM36,^{24,25} Amber-compatible Slipids^{28–30} with its 2020 update,³¹ Amber-compatible Lipid17,^{26,27} and OPLS-aa-compatible MacRog^{33–35} models. All simulations were 1 μ s long, totaling 72 μ s, and performed using GROMACS version 2020 or 2021.⁵⁹ The used simulation parameters are provided in Table S1, and the simulation data are available at DOI: 10.5281/zenodo.7035350 (CHARMM36), DOI: 10.5281/zenodo.7022749 (Slipids), DOI: 10.5281/zenodo.6992065 (Lipid17), and DOI: 10.5281/zenodo.7061800 (MacRog).

3.Run Lipid21 for revision?

Table 1: Details of the simulation systems provided for (small/medium/large) system sizes). The box dimensions in the membrane plane (x/y) and normal to the membrane (z) are provided for the Slipids simulations, and the values vary slightly between the force fields.

[CHOL]	POPC	CHOL	Water	x/y (nm)	z (nm)
0 mol-%	64/256/1024	0/0/0	3200/12800/51200	4.4/9.0/18.1	8.9/8.6/8.5
11 mol-%	64/256/1024	8/32/128	3600/14400/57600	4.5/9.4/18.1	9.4/9.1/9.3
20 mol-%	64/256/1024	16/64/256	4000/16000/64000	4.5/9.2/18.3	10.2/9.9/10.0
29 mol-%	64/256/1024	26/104/416	4500/18000/72000	4.6/9.2/18.5	10.8/10.8/10.7
38 mol-%	64/256/1024	40/160/640	5200/20800/83200	4.8/9.5/19.2	11.3/11.4/11.2
47 mol-%	64/256/1024	56/224/896	6000/24000/96000	5.0/10.1/20.0	11.8/11.6/11.7

2.3 Simulation Analyses

Electron Density Profiles and Bilayer Dimensions: The form factors and electron density profiles were calculated using the `form_factor.py` script in the NMRlipids Databank (<https://github.com/NMRLipids/Databank/>) after assigning all atom types with valence electron counts. The thicknesses were extracted by the `findpeaks` function in Matlab after smoothening the electron density data. Area per phospholipid was obtained by dividing the area of the bilayer by the number of phospholipids in one leaflet.

Order Parameters: The C–H order parameters were calculated as

$$S_{\text{CD}} = \left\langle \frac{3 \cos^2 \theta - 1}{2} \right\rangle, \quad (1)$$

where θ is the angle between the C–H vector and the z axis normal to the bilayer plane. In the POPC acyl chains, the values for the 2 (3) hydrogens of CH₂ (CH₃) groups were averaged, since these groups rotate freely and thus the order parameters are identical for both (all) hydrogens in experiments. In simulations, the values are also very similar, justifying this approach. An exception to this are the hydrogens bound to the 2nd carbon in the oleate chain; they lack rotational averaging in both both simulations and experiments, and were thus treated separately in our analyses. The C–H order parameters for the POPC head group were calculated similarly, but values for both hydrogens are shown separately for all CH₂ groups, *i.e.* no averaging was performed.

Lateral Diffusion Coefficients: The lateral diffusion coefficients D_{PBC} from simulations performed using periodic boundary conditions (PBC) were extracted from mean squared displacement (MSD) data calculated for lipid centers of mass after eliminating the drift of their host leaflet with the `gmx msd` tool. The MSD data were fit with a straight line in the

lag time (Δ) interval between 10 and 100 ns as

$$\text{MSD} = 4D_{\text{PBC}}\Delta. \quad (2)$$

The diffusion coefficients extracted from the three simulation box sizes were fitted with

$$D_{\text{PBC}} \approx D_{\infty} + \frac{k_{\text{B}}T}{4\pi\mu_{\text{m}}h} \frac{\ln [L / (L_{\text{SD}} + 1.565H)] - 1.713}{1 + H/L_{\text{SD}}}, \quad (3)$$

where D_{∞} is the lateral diffusion coefficient in an infinite system, h is the hydrodynamic thickness of the membrane, $k_{\text{B}}T$ is the Boltzmann constant, T is the temperature, μ_{m} and μ_{f} are shear viscosities of the membrane and the fluid (water), respectively. H is half the thickness of the water layer and $L_{\text{SD}} = \frac{h\mu_{\text{m}}}{2\mu_{\text{f}}}$ the Saffmann–Delbrück length.⁴⁷ The inter-leaflet friction coefficient does not appear in Eq. (3) as we expect it to be infinite, which was found to be a valid assumption for lipid bilayers.⁴⁷ The water viscosity value of 0.3228 mPa·s was interpolated to 298 K from the values for CHARMM TIP3P in Ref. 60 and used for all simulations (CHARMM TIP3P and normal TIP3P differ by $\sim 2\text{--}3\%$). The final simulation box dimensions were used for L and H . For h and H ($= (L_z - h)/2$), the average values from the three systems sizes were used.

More and Less Ordered Populations: The angles (θ) between the C–H vectors of the last six carbons of the palmitate chain were extracted and the data across all CHOL concentrations was accumulated. These data were fit with two Gaussians centered at 0,

$$P(\theta) = a_{\text{less}} \exp\left(-\frac{\theta}{2\sigma_{\text{less}}}\right) + a_{\text{more}} \exp\left(-\frac{\theta}{2\sigma_{\text{more}}}\right), \quad (4)$$

where the two standard deviations (σ) correspond to the more and less ordered components. After these σ values were established for a given force field, the data at all CHOL concentrations were independently fitted by Eq. (4) with the fixed σ values. These fits provided the

pre-factors, whose ratio $a_{\text{more}}/(a_{\text{more}} + a_{\text{less}})$ corresponds to the fraction of the more ordered component in the system.

This two-Gaussian approach assumes a “two-population” scheme, where the more and less ordered populations are present (at non-zero CHOL concentration), and that the addition of CHOL only affects their ratio, but not their natures. In an alternative picture of a “collective transition”, there would be a single Gaussian, whose standard deviation σ would decrease as a function of CHOL concentration as the membrane gets more ordered.

Quantitative Quality Measure for Lipid Bilayer Structure: The deviations (in %) from experimental values across CHOL concentrations were used to quantify the quality of the lipid force fields. The 1st and 2nd form factor minima locations, the electron density profiles (as a function of normal distance from bilayer center), the oleate and palmitate chain order parameters (as a function of carbon atoms in the acyl chains) and diffusion coefficients were interpolated to create continuous 2D matrices for the order parameters and density profiles, and a continuous 1D vector for the diffusion coefficients and form factor minima. For the 2D matrices, the absolute values of the differences of the matrices from simulations and experiments were calculated and averaged over the carbon atoms in the acyl chain (order parameters) or across the simulation system (density profiles). This resulted in 1D deviation vectors as a function of CHOL concentration. For diffusion coefficients and form factor minima, the absolute value of the difference of the interpolated 1D vectors of simulation and experimental data was calculated to provide deviation as a function of CHOL concentration. All these 1D vectors were normalized by dividing them by the experimental values to provide relative deviations in %.

3 Results and Discussion

4.Results: volmaps of chains around cholesterol?

5.Results: SI: RDFs of POPC chains and CHOL around CHOL

3.1 Cholesterol Condensing Effect is Manifested Differently in Different Force Fields

Lipid bilayer dimensions can be experimentally accessed by measuring X-ray scattering form factor, which is related to the electron density along membrane normal *via* a Fourier transform.^{43,49,57,61?} The form factor can be translated to an electron density profile, and area per lipid and bilayer thickness values can be extracted using the scattering density profile (SDP) model or its combination with MD simulations.^{49,54,57,61?},⁶² Sophisticated approaches for single component lipid bilayers are available but the data for multi-component systems are more difficult to interpret, and the used approaches involve severe approximations.^{49,57,61?}

Thus, we compare here the form factors and the total electron density profiles which do not require the de-convolution of the electron density profiles into densities of the different molecular species or their segments. For reference, all scattering intensities from experiment are shown in Figs. S1 and S2, the form factor profiles from experiment and simulation in Fig. S3 and the density profiles from experiment and simulation in Fig. S5.

As the absolute and relative lobe amplitudes depend on simulation box size (Fig. ??), the minima in the form factor profiles provide a robust comparison between simulation and experiment. Thus, we first traced the first two minima in the form factors, and they are plotted in Fig. S4. As simulations and experiments were performed at somewhat different CHOL concentrations, we interpolated the minima to the entire studied range of CHOL concentrations. The plots in Fig. S4 highlight that the addition of CHOL first shifts the first minima to smaller wave vector values in the experiment, which is reasonably well captured by the simulation force fields. CHARMM36 seems to be off more than the other three force fields. Above ~ 25 mol-% of CHOL, the location of the minimum shifts to larger wave vector values in the experiment, which is curiously not captured by any of the force fields. The

experiment shows a steady shift of the second minimum to smaller wave vector values, and this is reproduced by all simulation force fields. Slipids and Lipid17 are generally in best agreement with the experiment, whereas MacRog and CHARMM36 show poorer agreement.

Moving on to the electron density profiles extracted from the form factors, we generated two-dimensional maps for the density profiles using interpolation of the density profiles in Fig. S5 (see Methods). These maps are shown in Fig. 1 for both simulations (top row) and experiment (bottom row, left column).




Figure 1: Electron density profiles, thickness and area per lipid as a function of CHOL concentration. Top: Electron density maps for the simulations using four different force fields. Bottom left: Electron density map for the experiment. The scaling is uniform between all simulations and experiment. The original electron density profiles are shown in Fig. S5. The effect of system size on the density profiles in simulations is demonstrated in Fig. S12 in the SI. Bottom center: Bilayer thickness. Thickness is defined as twice the distance from the peak in electron density to the membrane core. Experimental data are extracted in a similar manner from electron density profiles obtained with X-ray scattering. Bottom right: area per lipid measured by dividing the total membrane area by the number of phospholipids. The experimental estimates, interpolated to 298 K, are taken from Refs. 63 (58.8 \AA^2), 64 (67.5 \AA^2), and 65 (63.5 \AA^2). The NMR data is measured in Ref. 66 using either ^{13}C (58.9 \AA^2) or ^2H (58.3 \AA^2) NMR. The size-dependency of area per lipid is shown in Fig. S11.

Overall, all electron density profiles share the same features across all CHOL concentrations; a high-density band corresponding to the tightly-packed interfacial region containing electron-rich phosphorus, a low-density region at the core of the membrane occupied by the disordered acyl chains, and intermediate density in the rest of the lipid regions as well as the aqueous phase. However, a more detailed look at the profiles in Fig. 1 reveals differences between the force fields. CHOL renders the membrane thicker, resulting in the spreading of the high-density bands with increasing amount of CHOL. In CHARMM36, Slipids, and MacRog, there is clear saturation of this effect at around 30 mol-% CHOL. In Lipid17, however, the membrane gets thicker until the maximum studied CHOL concentration of 47 mol-%. In experiment, the effect of CHOL is generally weaker, and some saturation is evident at high CHOL concentrations. These trends are visible in the bottom middle panel in Fig. 1, which shows the effect of CHOL on membrane thickness extracted from the density peak positions. Fig. 1 also demonstrates that CHARMM36 has the sharpest low- and high-density bands indicating smaller membrane fluctuations that would smear the bands, and the same is true for MacRog at higher cholesterol concentrations. The less sharp bands for Lipid17 and especially Slipids profiles indicate that the membranes are more flexible with these force fields.

Comparison against experiments (bottom left panel in Fig. 1) indicates that the membrane structure is more uniform in the experiment, as the bands are sharper. This is also evident in the original electron density profiles in Fig. S5. System size plays a role here, as demonstrated by the density maps calculated for the CHARMM36 simulations in three sizes, shown in Fig. S12. The larger the system, the larger the bilayer undulations are, which can smear the features of the density profiles. Still, even in the smallest system, the band intensities are not as localized as in the experiment. This difference likely partially stems from the features of the used SDP model, but it could also indicate different elastic properties between simulation and experiment.

Comparison of the thickness values (bottom middle panel in Fig. 1) demonstrates that the

Lipid17 bilayers are too thin at low CHOL concentrations, whereas other models agree well with experiment in this regime. Still, all simulation models overshoot the experiment at high CHOL concentrations. Experiment and all simulation models except Lipid17 demonstrate saturation after 30 mol-%.

The membrane-condensing effect of CHOL is well documented, yet there is some ambiguity surrounding the term.⁶⁷ The cross-sectional area of a CHOL molecule is smaller than that of a phospholipid, so an increase in CHOL concentration is expected to lower the average *area per lipid* (including CHOL). However, condensation could also refer to a decrease in the *area per phospholipid* upon the addition of cholesterol,⁶⁷ or to CHOL having a diminishing partial area, meaning that a certain amount of CHOL could be added to a phospholipid bilayer without affecting its total area.⁶⁸ We thus compared the behavior of the area per (POPC) lipid (APL) as a function of CHOL concentration in the bottom right panel of Fig. 1. The dependence of APL on CHOL concentration follows the trends in thickness inversely in general, yet provide curious differences between the force fields. Lipid17 has the largest APL across the entire CHOL contraction range, and the addition of CHOL always leads to an increase in the membrane area. MacRog also has a large APL for pure POPC, but the partial area of CHOL is negative until 30 mol-% concentration indicating a particularly strong condensation effect. The profiles for Slipids and CHARMM36 are very similar, and the partial area of CHOL is small or even zero until a concentration of 20 mol-%. Importantly, the observed differences in CHOL condensing effect take place at the physiologically relevant CHOL concentration range.² Experimental data are derived from models, so they can vary greatly, as evidenced by the very different values obtained for POPC by scattering and NMR experiments performed by different teams (See Fig. 1 and its caption). Still, all the simulation force fields provide areas per lipid that fall within the experimental spread for pure POPC.

3.2 Acyl Chain Ordering Varies Greatly Between the Force Fields

Order parameters for C–H bond vectors measured with ^{13}C or ^2H NMR techniques, provide indirect information about the structural sampling of individual molecules.⁴³ Moreover, these parameters are readily extracted from the simulation. An increase in acyl chain order parameters with added cholesterol in simulations and experiments can be explained by an increased fraction of *anti* conformations in the acyl chains,^{42?} which is suggested to play critical role in the phase behaviour of PC–cholesterol mixtures.⁶ Consequently, the correct cholesterol ordering effect is expected to be a necessary condition for a model used to understand lipid–cholesterol phase behaviour.

Here, as the NMR experiments⁴² were again performed at different CHOL concentrations than simulations (and X-ray scattering measurements), we generated order parameter maps as a function of acyl chain carbon number and cholesterol concentration for both simulations and experiments, and subtracted these maps to obtain the deviation of simulation from experiment. These maps are shown for the palmitate (top row) and oleate (bottom row) chains of POPC in Fig. 2, whereas the original order parameter profiles are shown in Figs. S6 and ??.

These original maps in Figs. S6 and ?? demonstrate that the addition of cholesterol substantially increases the acyl chain C–H order parameters for all simulation models and experiments, which can be explained by increased *anti* conformations in acyl chains.^{42?} The deviations mapped in Fig. 2 provide an intuitive way for a quantitative comparison of the behaviors of the force fields. Here, deviations in the range $[-0.02, 0.02]$ are considered to fall within the error estimate and are thus shown in white. Blue indicates that the order parameters are too low in the simulation (their absolute value is larger and the chains are too ordered). Overall, the simulation models behave reasonably well at low CHOL concentrations, but deviate significantly from experiment at higher CHOL concentrations.

In CHARMM36, the both the palmitate and oleate chains get too ordered upon increasing CHOL concentration, yet the oleate C9 carbon next to the double bond is too disordered in




Figure 2: POPC acyl chain order parameter deviation from experiments. Data are shown for palmitate (top row) and oleate (bottom row) and for the four force fields (columns). Negative values indicate that order is too high ($-S_{CD}$ values too negative) in the simulations. The values that are within the estimated error range of 0.02 from experiment are considered to be “correct” and coloured in white. The order parameters calculated for the two or three identical carbons were very similar and thus averaged over in simulations, as they have identical values in experiments. For the C2 carbon of the oleate chain, the order parameters are forked in simulations and experiments. We thus calculated the differences for both the larger and smaller values, and averaged this deviation for the C2 column of oleate.

the simulation. The oleate chain shows best agreement with experiment in Slipids with only slight excess ordering at higher CHOL concentrations. The excess ordering is more severe for the palmitate chain. Still, the major discrepancy between Slipids and experiment is the drastically too disordered C2 and C3 carbons of the palmitate chain. This effect was not present in the original Slipids model,³⁸ indicating that it was introduced in the recent reparametrization that improved the head group and glycerol backbone structures of Slipids.³¹ Lipid17 provides the best overall agreement with experiment, as no segments deviate significantly from experiment at any CHOL concentrations. At low CHOL concentrations, the middle of the oleate chain is too disordered, which agrees with the large APL observed in Fig. 1. At large CHOL concentrations, the oleate chain agrees well with experiment, whereas the palmitate chain has a slightly too ordered C9–C12 segment, yet the deviation is significantly

smaller than in the other force fields. MacRog behaves reasonably well at low CHOL concentrations, yet the C2–C5 carbons in palmitate are slightly too disordered, whereas the C2 of oleate is somewhat too ordered. However, at larger CHOL concentrations the chains become overall way too ordered, leading to the largest overall deviations from experiment.

The head group order parameters are not largely affected by CHOL concentration in experiments,⁴² which has been discussed in our earlier work.³⁸ The four force fields evaluated here also reproduce this behavior to a great degree (Fig. S8), so the limitations observed in the absence of CHOL³⁸ largely persist across all studied CHOL concentrations. Moreover, it has now become standard to consider them in the parameterization and validation of lipid force fields.^{31,69–71}

7. Cholesterol order parameters? Into the deviation metric?

3.3 The Force Fields Predict Very Different Lateral Mobilities

Apart from the ordering effect on the bilayer structure, CHOL is also known to make them stiffer and less dynamic.^{8,44,45} The comparison between lateral diffusion coefficients extracted from simulation and experiment has been limited due to a box-size dependence observed in simulations performed using periodic boundary conditions.^{46,47,72} Here, we tackle this issue by performing simulations with three system sizes, which allows the extrapolation of the results to an infinite system with the theoretical description developed by Vögele and Hummer.^{47,47} Importantly, reference values are available for POPC/CHOL mixtures from ¹H pulsed field gradient NMR diffusion measurements on label-free macroscopically aligned bilayers. The PBC-corrected lateral diffusion coefficients are shown in the top panel of Fig. 3 together with experimental values. The size-dependence of these values extracted from simulations together with the fit of Eq. (3) are shown in Fig. S9, whereas the CHOL-dependence of these values in systems with different sizes are shown in Fig. S10.

The lipid models again show significantly different behavior. CHARMM36 and Lipid17 provide drastically too large diffusion coefficients at low CHOL concentrations, yet they con-




Figure 3: **Dynamic properties of the POPC/cholesterol mixtures.** Top: Lateral diffusion coefficients corrected for finite-size effects using Eq. (??). Experimental data are taken from NMR measurements on well-hydrated samples.^{44,45} The hollow circles show data extracted for Slipids using a shorter Lennard-Jones cutoff (see text). Bottom: Shear viscosities obtained from the finite-size correction. Eq. (??). The size-dependence of lateral diffusion and the fits used to obtain the PBC-corrected diffusion coefficients and shear viscosities are shown in Figs. S9 and S10.

verge towards experimental estimates upon the addition of CHOL. This similar behavior is somewhat surprising, as they behave very differently in terms of both area per POPC (Fig. 1) and acyl chain ordering (Fig. 2). For CHARMM36, the excessive ordering observed at high CHOL concentrations counterbalances the too fast dynamics observed at low CHOL concentrations. However, the behavior of Lipid17 cannot be explained by the order parameters, as they agree with experiment relatively well across all studied CHOL concentrations. The area per POPC is exaggerated at all CHOL concentrations, yet the deviation from experiment decreases towards larger CHOL concentrations. Thus, it seems that CHARMM36 dynamics correlate with chain ordering, whereas those of Lipid17 go hand in hand with area per POPC. With MacRog, the dynamics of POPC are already slightly too slow compared to experiment, and the excessive ordering at large CHOL concentrations leads to a larger underestimation

of the experimental values. Curiously, Slipids provides an essentially quantitative agreement with experiment across the studied CHOL concentrations, and thus significantly outperforms other force fields in terms of lateral dynamics.

Slipids has the largest Lennard-Jones (LJ) cutoff of the used force fields of 1.4 nm (see Table S1), and this cutoff is known to affect dynamic properties.⁷³ Therefore, we repeated simulations at 0 mol-% and 47 mol-% CHOL for Slipids but with a cutoff of 0.9 nm corresponding to that of Lipid17, and the PBC-corrected diffusion coefficient values are shown in the top panel of Fig. 3 as hollow green circles. The decrease in cutoff indeed leads to a minor increase in the lateral diffusion coefficient. However, the LJ cutoff alone does not explain the differences between force fields, indicating that they truly spark from differences in the force field interaction parameters.

As the results in Fig. S10 demonstrate, the PBC correction can qualitatively change the trends observed for diffusion coefficients. This results from the fact that the size of the PBC correction, Eq. (3), depends on membrane viscosity, which further depends on CHOL concentration. For example, Lipid17 and CHARMM36 underestimate the experimental values in all system sizes used here and the effect of CHOL seems to be well reproduced. However, the corrected values significantly overshoot the experiment at low CHOL concentration, and CHOL induces a more drastic slowdown in simulations so that the simulations and experiment almost agree at 47 mol-% CHOL. With MacRog, even the corrected values fall below the experimental ones, yet the slowdown effect of CHOL again seems stronger than in experiment. Slipids is a clear outlier, as the CHOL-dependence seems too weak with finite system sizes, yet after accounting for PBC effects, the agreement with experiment is excellent. These results highlight that little can be said about CHOL-dependence of lateral diffusion coefficients without performing multiple simulations required for the PBC correction at multiple CHOL concentrations. When fine-tuning interaction parameters, this can lead to a massive number of simulations.

The PBC correction, Eq. (3) also provides the shear viscosity of the membrane as a fitting

parameter. Instead of shear viscosity, the surface viscosity is sometimes reported, yet this is simply the product of shear viscosity and membrane thickness (available in Fig. 1). The shear viscosity values are in line with the lateral diffusion coefficients.

3.4 Cholesterol Induces Order in a Non-collective Manner

CHOL could induce order in a lipid membrane in two ways; it could either increase the average order of all phospholipids (“collective transition”), or increase the prevalence of the ordered lipids (“two state model”). To clarify this effect, we extracted the C–H angle distributions with respect to the xy plane for the last six carbons of the palmitate chain from all force fields. We first accumulated data at all CHOL concentrations, and observed that these data could be fitted with two Gaussian functions. These Gaussians were centered at 0° , and with the different standard deviations they corresponding to the more and less ordered acyl chains. The obtained standard deviations for the more and less ordered chains only showed minor differences between the force fields (rightmost panel in Fig. 4). We then fitted the C–H angle distributions for all CHOL concentrations with the two Gaussians with the obtained standard deviations and with their means set to 0. These Gaussians provided excellent fits to the data with $R^2 = 0.991 \pm 0.005$ (see the leftmost panel of Fig. 4 for the fits to the Slipids data). The prefactors of the Gaussian functions were considered to represent the fractions of the more and less ordered components in the system. In pure POPC, the ordered component was absent, yet its prevalence increased linearly as a function of CHOL, after saturating at high CHOL concentrations (rightmost panel in Fig. 4). The success of the two-Gaussian fits in describing the data indicates that with all force fields the bilayer is composed of lipids ordered by CHOL and lipids corresponding to their state in pure POPC membrane, *i.e.* corresponding to the “two state model”, and CHOL simply changes the fraction of these two states present. Indeed, single Gaussian fits with the standard deviation as a free parameter fail when CHOL is present, indicating that the ordering does not occur *via* a “collective transition”.




Figure 4: **Distributions of C-H bond orientations.** Leftmost panel: The probability distributions for the angle of C-H vectors of the last 6 carbons in the palmitate chain of POPC with respect to the xy -plane. All distributions are centered around 0, but have different shapes. The markers show data extracted from Slipids simulations at various CHOL concentrations, and the solid lines show two-Gaussian fits with fixed standard deviations to the data. These Gaussians correspond to the more and less ordered palmitate chains. The fixed standard deviations were extracted from a fit of two Gaussians to the data accumulated across all CHOL concentrations. The fit with the fixed standard deviations to all CHOL concentrations are excellent with $R^2 = 0.991 \pm 0.005$ (mean \pm standard deviation). Middle panel: The standard deviations of the two Gaussians fitted to the C-H distribution (such as the one in the leftmost panel) for the four force fields. Rightmost panel: The fraction of the more ordered component as a function of CHOL concentration extracted simply by dividing the prefactor in the Gaussian distribution for the more ordered component by the sum of the two prefactors. Notably, in the CHOL-free system, the ordered component is not present.

3.5 Quality of Force Fields at Various Cholesterol Concentrations

The previous sections highlight that the agreement of the studied force fields with experiment depends on both the parameter in question or the CHOL concentration. To capture these features in an easily digestible form, we have calculated the relative deviations of the form factor minima, the order parameters of the two acyl chains, and the diffusion coefficients between the experiments and all the force fields. The interpolated data were used, and the relative deviation from experiments was obtained by dividing the difference of simulated and experimental data by the latter. These data are shown in the two leftmost columns in Fig. 5 as stacked plots.

The comparison of force fields in the two leftmost columns of Fig. 5 reveals that CHARMM36 and MacRog never provide the best overall agreement with experiment. In fact, for CHARMM36




Figure 5: **Total deviation of the force fields from experimental data.** Two leftmost columns: The relative deviations from the first two minima in the form factor, palmitate and oleate chain order parameters, and diffusion coefficients are shown in a cumulative manner to highlight the overall deviation of the force fields from experimental data. The order parameter deviations are obtained by averaging over the columns in Fig. 2 and normalizing against experimental data. The differences in the form factor minima in Fig. ?? between experiment and simulation were calculated, normalized against experimental data, and summed together. The diffusion coefficient deviation is the difference of values from simulation and experiment in Fig. 3, taken after interpolation to the same CHOL values as shown in Figs. 2 and 1, and normalized against experimental values. Top right panel: The zoomed-in view to the deviation in the location of the first minimum in the form factor. Bottom right panel: Deviation of the density profiles calculated from the form factor using the SDS model. The density profile deviations are calculated as the difference between the simulation and experimental maps in Fig. 1 followed by averaging over the columns and normalization against experiment.

all contributions to the deviation metric are always significant, and it shows the poorest agreement with the form factor minima. MacRog performs relatively well at low CHOL concentrations, but its quality deteriorates significantly upon the addition of CHOL. At low CHOL concentration, Slipids provides the best agreement with experiment, yet its quality decreases slightly at higher CHOL concentrations. Still, Slipids is the second best force field even in the high CHOL regime. Lipid17 provides the worst agreement with experiment at

low CHOL concentrations, but improves significantly towards larger concentrations so that at above ~ 30 mol-% of CHOL, Lipid17 provides the best overall agreement with experiment. Since the contributions from the form factor minima deviations are barely visible for some force fields in the leftmost columns of Fig. 5, we have highlighted the error in the 1st minimum location in the top right panel of Fig. 5. This panel demonstrates that apart from CHARMM36, all force fields capture the location of the 1st minimum relatively well at low CHOL concentrations, but deviate significantly at larger CHOL concentrations. Finally, we also analyzed the deviations in the density profiles in the bottom right panel of Fig. 5. This analysis demonstrates that even the significant deviations as well as the trend as a function of CHOL concentration in the form factor minima are lost as form factors are converted to density profiles. This indicates that the form factor minima are a better target value for force field development and refinement.

3.6 Molecular Level Structure of PC-CHOL Mixtures

Molecular dynamics simulations provide a description of the molecular organization of POPC and CHOL in the bilayer, yet the reliability of this description depends on the quality of the used force field. In the previous section we established that Slipids and Lipid17 provide the best agreement with experimental properties at low and high CHOL concentrations, respectively. Thus, it is reasonable to assume that the molecular pictures provided by these two force fields at the corresponding CHOL concentration regimes are the most realistic.

Earlier studies have analyzed the organization of CHOL and DSPC chains and observed five and three density peaks for DSPC chains and other CHOL molecules around CHOL molecules.⁷⁴

2D density maps RDFs to SI?

Apart from the lateral organization of the molecules, their orientations might also differ between the force fields. However, we observed only small differences in the tilt of CHOL molecules, defined by the angle between a vector spanning the rigid ring structure as well as

the z axis (membrane lies in the x, y plane). The mean angles and their standard deviations are shown in Fig. S13. Lipid17 and Slipids show very consistent values. CHARMM36 has slightly smaller tilt angles than them across all CHOL concentrations. MacRog has the most tilted CHOL molecules at low CHOL concentrations, but their tilt angle is the smallest at large CHOL concentrations. Still, the differences are overall minor.

4 Conclusions

To conclude, the recent all-atom force fields all capture the overall trends of CHOL-PC interaction reasonably well; The addition of CHOL leads to acyl chain ordering, which goes hand in hand with the increase in bilayer thickness. Area per lipid increases less than what would be expected if the areas of POPC and CHOL were additive, manifesting a condensation effect of CHOL. However, beyond these qualitative features, the force fields behave very differently. At the physiological CHOL concentrations in the range from 0 to 30 mol-%, CHOL has a positive partial area in Slipids and Lipid17, a zero partial area in CHARMM36, and a negative partial area in MacRog. In addition to these trends, the absolute values of area per lipid and bilayer thickness vary substantially. At larger CHOL concentrations all force fields predict a too thick membrane, indicating an excessive ordering effect of CHOL. This is indeed verified by the order parameter maps, which suggest that the palmitate chain is overly ordered in all force fields to a varying degree. The best agreement with experiment is achieved with Slipids at low CHOL concentration and with Lipid17 at high CHOL concentration. In terms of dynamics, the lateral diffusion in pure POPC is too fast in Lipid17 and CHARMM36, and the addition of CHOL slows them both down more than in experiments. MacRog has overall too slow dynamics, whereas Slipids agrees quantitatively with the experiment, indicating that it provides the best description of lipid diffusion.

None of the force fields unambiguously provides the best agreement with experiment.

Slipids shows great description of lateral dynamics, and reasonable behavior in terms of membrane thickness and acyl chain ordering, especially at lower and intermediate CHOL concentrations. On the other hand, Lipid17 provides a good description for the palmitate chain ordering, but is otherwise in a relatively poor agreement with experiment except for higher CHOL concentrations. Thus, the choice of force field depends not only on the choice of membrane properties (structural vs. dynamic) relevant to the problem, but also on the CHOL concentration range of interest. At low concentrations, Slipids is a safe choice, but at higher concentrations Lipid17 might provide the most realistic description. Since many research questions will involve other lipids, proteins, sugars, or drugs, so the quality of the force fields describing these additional molecules might be the decisive factor, together with force field compatibility. Finally, it must be noted that while all simulations were performed with their suggested simulation parameters, the different simulation engines might provide slightly different behavior, yet we believe that evaluating the magnitude of these effects is well beyond the scope of this work.

Supporting Information Available

Simulation parameters used for all the studied force fields. Form factor plots from simulations and experiments with the traced minima at various cholesterol concentrations. Interpolated 1st and 2nd minima of the form factors as a function of cholesterol concentration from simulation and experiment. Electron density profiles from simulation and experiment at various cholesterol concentrations. Deuterium order parameters of both acyl chains of POPC from simulation and experiment at various cholesterol concentrations. Deviation of POPC head group order parameters from experiment as a function of cholesterol concentration. Lateral diffusion coefficients with varying cholesterol concentrations as a function of simulation box size from simulations together with the extrapolation of these values to an infinite box size. Diffusion coefficients as a function of cholesterol concentration from simulations with varying

box sizes together with comparison to experiment. Effect of simulation box size on the area per lipid across the studied cholesterol concentrations. Effect of simulation box size on the density profiles across the studied cholesterol concentrations. Cholesterol tilt as a function of cholesterol concentration from simulations.

Acknowledgement

We acknowledge CSC – IT Center for Science for computational resources. JJM thanks the Amber community (Benjamin Madej from the Walker lab and Callum Dickson from the Gould lab) for technical assistance related to the lipid14 force field parameters. MJ thanks the Academy of Finland (Postdoctoral researcher grant no. 338160) and the Emil Aaltonen foundation for funding.

References

- (1) Lorent, J.; Levental, K.; Ganesan, L.; Rivera-Longsworth, G.; Sezgin, E.; Doktorova, M.; Lyman, E.; Levental, I. Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nature chemical biology* **2020**, *16*, 644–652.
- (2) Van Meer, G.; Voelker, D. R.; Feigenson, G. W. Membrane lipids: where they are and how they behave. *Nature reviews Molecular cell biology* **2008**, *9*, 112–124.
- (3) Wang, H.-Y.; Bharti, D.; Levental, I. Membrane heterogeneity beyond the plasma membrane. *Frontiers in Cell and Developmental Biology* **2020**, *8*, 580814.
- (4) Kinnun, J. J.; Bolmatov, D.; Lavrentovich, M. O.; Katsaras, J. Lateral heterogeneity and domain formation in cellular membranes. *Chemistry and Physics of Lipids* **2020**, *232*, 104976.

- (5) Mouritsen, O. G.; Zuckermann, M. J. What's so special about cholesterol? *Lipids* **2004**, *39*, 1101–1113.
- (6) Ipsen, J. H.; Karlström, G.; Mouritsen, O.; Wennerström, H.; Zuckermann, M. Phase equilibria in the phosphatidylcholine-cholesterol system. *Biochim. Biophys. Acta* **1987**, *905*, 162 – 172.
- (7) Kinnunen, P. K. On the principles of functional ordering in biological membranes. *Chemistry and Physics of Lipids* **1991**, *57*, 375 – 399.
- (8) Róg, T.; Pasenkiewicz-Gierula, M.; Vattulainen, I.; Karttunen, M. Ordering effects of cholesterol and its analogues. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **2009**, *1788*, 97–121.
- (9) Simons, K.; Ikonen, E. Functional rafts in cell membranes. *Nature* **1997**, *387*, 569–572.
- (10) Cebecauer, M.; Amaro, M.; Jurkiewicz, P.; Sarmiento, M. J.; Sachl, R.; Cwiklik, L.; Hof, M. Membrane lipid nanodomains. *Chemical reviews* **2018**, *118*, 11259–11297.
- (11) Milovanovic, D.; Honigmann, A.; Koike, S.; Göttfert, F.; Pähler, G.; Junius, M.; Müller, S.; Diederichsen, U.; Janshoff, A.; Grubmüller, H.; Eggeling, C.; Hell, S. W.; van den Bogaart, G.; Jahn, R. Hydrophobic mismatch sorts SNARE proteins into distinct membrane domains. *Nature Communications* **2015**, *6*, 1–10.
- (12) Kelkar, D. A.; Chattopadhyay, A. Modulation of gramicidin channel conformation and organization by hydrophobic mismatch in saturated phosphatidylcholine bilayers. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **2007**, *1768*, 1103–1113.
- (13) Killian, J. A. Hydrophobic mismatch between proteins and lipids in membranes. *Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes* **1998**, *1376*, 401–416.

- (14) Gimpl, G. Interaction of G protein coupled receptors and cholesterol. *Chemistry and physics of lipids* **2016**, *199*, 61–73.
- (15) Guixà-González, R.; Albasanz, J. L.; Rodriguez-Espigares, I.; Pastor, M.; Sanz, F.; Martí-Solano, M.; Manna, M.; Martinez-Seara, H.; Hildebrand, P. W.; Martín, M.; Selent, J. Membrane cholesterol access into a G-protein-coupled receptor. *Nature Communications* **2017**, *8*, 1–12.
- (16) Róg, T.; Vattulainen, I. Cholesterol, sphingolipids, and glycolipids: What do we know about their role in raft-like membranes? *Chem. Phys. Lipids* **2014**, *184*, 82 – 104.
- (17) Berkowitz, M. L. Detailed molecular dynamics simulations of model biological membranes containing cholesterol. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **2009**, *1788*, 86–96.
- (18) Leonard, A. N.; Wang, E.; Monje-Galvan, V.; Klauda, J. B. Developing and testing of lipid force fields with applications to modeling cellular membranes. *Chemical reviews* **2019**, *119*, 6227–6269.
- (19) Marrink, S. J.; Corradi, V.; Souza, P. C.; Ingolfsson, H. I.; Tieleman, D. P.; Sansom, M. S. Computational Modeling of Realistic Cell Membranes. *Chemical Reviews* **2019**, *119*, 6184–6226.
- (20) Brooks, B. R.; Bruccoleri, R. E.; Olafson, B. D.; States, D. J.; Swaminathan, S. a.; Karplus, M. CHARMM: a program for macromolecular energy, minimization, and dynamics calculations. *Journal of computational chemistry* **1983**, *4*, 187–217.
- (21) Cornell, W. D.; Cieplak, P.; Bayly, C. I.; Gould, I. R.; Merz, K. M.; Ferguson, D. M.; Spellmeyer, D. C.; Fox, T.; Caldwell, J. W.; Kollman, P. A. A second generation force field for the simulation of proteins, nucleic acids, and organic molecules. *Journal of the American Chemical Society* **1995**, *117*, 5179–5197.

- (22) Jorgensen, W. L.; Tirado-Rives, J. The OPLS force field for proteins. Energy minimizations for crystals of cyclic peptides and crambin. *J. Am. Chem. Soc.* **1988**, *110*, 1657–1666.
- (23) Harder, E.; Damm, W.; Maple, J.; Wu, C.; Reboul, M.; Xiang, J. Y.; Wang, L.; Lupyan, D.; Dahlgren, M. K.; Knight, J. L., et al. OPLS3: a force field providing broad coverage of drug-like small molecules and proteins. *Journal of chemical theory and computation* **2016**, *12*, 281–296.
- (24) Klauda, J. B.; Kučerka, N.; Brooks, B. R.; Pastor, R. W.; Nagle, J. F. Simulation-Based Methods for Interpreting X-Ray Data from Lipid Bilayers. *Biophys. J.* **2006**, *90*, 2796 – 2807.
- (25) Lim, J. B.; Rogaski, B.; Klauda, J. B. Update of the Cholesterol Force Field Parameters in CHARMM. *J. Phys. Chem. B* **2012**, *116*, 203–210.
- (26) Dickson, C. J.; Madej, B. D.; Skjevik, A. A.; Betz, R. M.; Teigen, K.; Gould, I. R.; Walker, R. C. Lipid14: The Amber Lipid Force Field. *J. Chem. Theory Comput.* **2014**, *10*, 865–879.
- (27) Madej, B. D.; Gould, I. R.; Walker, R. C. A Parameterization of Cholesterol for Mixed Lipid Bilayer Simulation within the Amber Lipid14 Force Field. *J. Phys. Chem. B* **2015**, *119*, 12424–12435.
- (28) Jämbeck, J. P. M.; Lyubartsev, A. P. Derivation and Systematic Validation of a Refined All-Atom Force Field for Phosphatidylcholine Lipids. *J. Phys. Chem. B* **2012**, *116*, 3164–3179.
- (29) Jämbeck, J. P. M.; Lyubartsev, A. P. An Extension and Further Validation of an All-Atomistic Force Field for Biological Membranes. *J. Chem. Theory Comput.* **2012**, *8*, 2938–2948.

- (30) Jämbeck, J. P. M.; Lyubartsev, A. P. Another Piece of the Membrane Puzzle: Extending Slipids Further. *Journal of Chemical Theory and Computation* **2013**, *9*, 774–784.
- (31) Grote, F.; Lyubartsev, A. P. Optimization of slipids force field parameters describing headgroups of phospholipids. *The Journal of Physical Chemistry B* **2020**, *124*, 8784–8793.
- (32) Maciejewski, A.; Pasenkiewicz-Gierula, M.; Cramariuc, O.; Vattulainen, I.; Róg, T. Refined OPLS All-Atom Force Field for Saturated Phosphatidylcholine Bilayers at Full Hydration. *J. Phys. Chem. B* **2014**, *118*, 4571–4581.
- (33) Kulig, W.; Tynkkynen, J.; Javanainen, M.; Manna, M.; Róg, T.; Vattulainen, I.; Jungwirth, P. How well does cholesteryl hemisuccinate mimic cholesterol in saturated phospholipid bilayers? *Journal of Molecular Modeling* **2014**, *20*.
- (34) Kulig, W.; Jurkiewicz, P.; Olżyńska, A.; Tynkkynen, J.; Javanainen, M.; Manna, M.; Róg, T.; Hof, M.; Vattulainen, I.; Jungwirth, P. Experimental determination and computational interpretation of biophysical properties of lipid bilayers enriched by cholesteryl hemisuccinate. *Biochim. Biophys. Acta* **2015**, *1848*, 422 – 432.
- (35) Kulig, W.; Pasenkiewicz-Gierula, M.; Róg, T. Cis and trans unsaturated phosphatidylcholine bilayers: A molecular dynamics simulation study. *Chem. Phys. Lipids* **2016**, *195*, 12 – 20.
- (36) Lee, J.; Cheng, X.; Swails, J. M.; Yeom, M. S.; Eastman, P. K.; Lemkul, J. A.; Wei, S.; Buckner, J.; Jeong, J. C.; Qi, Y.; Jo, S.; Pande, V. S.; Case, D. A.; Brooks, C. L.; MacKerell, A. D.; Klauda, J. B.; Im, W. CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *Journal of Chemical Theory and Computation* **2016**, *12*, 405–413.

- (37) Lee, J.; Hitzenberger, M.; Rieger, M.; Kern, N. R.; Zacharias, M.; Im, W. CHARMM-GUI supports the Amber force fields. *The Journal of chemical physics* **2020**, *153*, 035103.
- (38) Botan, A.; Favela-Rosales, F.; Fuchs, P. F. J.; Javanainen, M.; Kanduč, M.; Kulig, W.; Lamberg, A.; Loison, C.; Lyubartsev, A.; Miettinen, M. S.; Monticelli, L.; Määttä, J.; Ollila, O. H. S.; Retegan, M.; Róg, T.; Santuz, H.; Tynkkynen, J. Toward Atomistic Resolution Structure of Phosphatidylcholine Headgroup and Glycerol Backbone at Different Ambient Conditions. *J. Phys. Chem. B* **2015**, *119*, 15075–15088.
- (39) Catte, A.; Giryh, M.; Javanainen, M.; Loison, C.; Melcr, J.; Miettinen, M. S.; Monticelli, L.; Määttä, J.; Oganessian, V. S.; Ollila, O. S., et al. Molecular electrometer and binding of cations to phospholipid bilayers. *Physical Chemistry Chemical Physics* **2016**, *18*, 32560–32569.
- (40) Antila, H.; Buslaev, P.; Favela-Rosales, F.; Ferreira, T. M.; Gushchin, I.; Javanainen, M.; Kav, B.; Madsen, J. J.; Melcr, J.; Miettinen, M. S., et al. Headgroup structure and cation binding in phosphatidylserine lipid bilayers. *The Journal of Physical Chemistry B* **2019**, *123*, 9066–9079.
- (41) Bacle, A.; Buslaev, P.; Garcia-Fandino, R.; Favela-Rosales, F.; Mendes Ferreira, T.; Fuchs, P. F.; Gushchin, I.; Javanainen, M.; Kiirikki, A. M.; Madsen, J. J., et al. Inverse conformational selection in lipid–protein binding. *Journal of the American Chemical Society* **2021**, *143*, 13701–13709.
- (42) Ferreira, T. M.; Coreta-Gomes, F.; Ollila, O. H. S.; Moreno, M. J.; Vaz, W. L. C.; Topgaard, D. Cholesterol and POPC segmental order parameters in lipid membranes: solid state ^1H - ^{13}C NMR and MD simulation studies. *Phys. Chem. Chem. Phys.* **2013**, *15*, 1976–1989.
- (43) Ollila, O. S.; Pabst, G. Atomistic resolution structure and dynamics of lipid bilayers in

- simulations and experiments. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2016**, *1858*, 2512 – 2528, Biosimulations of lipid membranes coupled to experiments.
- (44) Filippov, A.; Orädd, G.; Lindblom, G. The effect of cholesterol on the lateral diffusion of phospholipids in oriented bilayers. *Biophysical journal* **2003**, *84*, 3079–3086.
- (45) Filippov, A.; Orädd, G.; Lindblom, G. Influence of cholesterol and water content on phospholipid lateral diffusion in bilayers. *Langmuir* **2003**, *19*, 6397–6400.
- (46) Vögele, M.; Hummer, G. Divergent diffusion coefficients in simulations of fluids and lipid membranes. *The Journal of Physical Chemistry B* **2016**, *120*, 8722–8732.
- (47) Vögele, M.; Köfinger, J.; Hummer, G. Hydrodynamics of diffusion in lipid membrane simulations. *Physical Review Letters* **2018**, *120*, 268104.
- (48) Heftberger, P.; Kollmitzer, B.; Heberle, F. A.; Pan, J.; Rappolt, M.; Amenitsch, H.; Kučerka, N.; Katsaras, J.; Pabst, G. Global small-angle X-ray scattering data analysis for multilamellar vesicles: the evolution of the scattering density profile model. *J. Appl. Crystallogr.* **2014**, *47*, 173–180.
- (49) Heftberger, P.; Kollmitzer, B.; Rieder, A. A.; Amenitsch, H.; Pabst, G. In Situ Determination of Structure and Fluctuations of Coexisting Fluid Membrane Domains. *Biophys. J.* **2015**, *108*, 854 – 862.
- (50) Rieder, A.; Koller, D.; Lohner, K.; Pabst, G. Optimizing rapid solvent exchange preparation of multilamellar vesicles. *Chemistry and physics of lipids* **2015**, *186*, 39–44.
- (51) Belička, M.; Weitzer, A.; Pabst, G. High-resolution structure of coexisting nanoscopic and microscopic lipid domains. *Soft Matter* **2017**, *13*, 1823–1833.
- (52) Buboltz, J. T.; Feigenson, G. W. A novel strategy for the preparation of liposomes: rapid solvent exchange. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1999**, *1417*, 232–245.

- (53) Heberle, F.; Pan, J.; Standaert, R.; Drazba, P.; Kučerka, N.; Katsaras, J. Model-based approaches for the determination of lipid bilayer structure from small-angle neutron and X-ray scattering data. *Eur. Biophys. J.* **2012**, *41*, 875–890.
- (54) Kučerka, N.; Nagle, J. F.; Sachs, J. N.; Feller, S. E.; Pencer, J.; Jackson, A.; Katsaras, J. Lipid Bilayer Structure Determined by the Simultaneous Analysis of Neutron and X-Ray Scattering Data. *Biophys. J.* **2008**, *95*, 2356–2367.
- (55) Kučerka, N.; Holland, B. W.; Gray, C. G.; Tomberli, B.; Katsaras, J. Scattering Density Profile Model of POPG Bilayers As Determined by Molecular Dynamics Simulations and Small-Angle Neutron and X-ray Scattering Experiments. *J. Phys. Chem. B* **2012**, *116*, 232–239.
- (56) Kučerka, N.; Perlmutter, J. D.; Pan, J.; Tristram-Nagle, S.; Katsaras, J.; Sachs, J. N. The Effect of Cholesterol on Short- and Long-Chain Monounsaturated Lipid Bilayers as Determined by Molecular Dynamics Simulations and X-Ray Scattering. *Biophys. J.* **2008**, *95*, 2792 – 2805.
- (57) Pan, J.; Cheng, X.; Heberle, F. A.; Mostofian, B.; Kučerka, N.; Drazba, P.; Katsaras, J. Interactions between Ether Phospholipids and Cholesterol As Determined by Scattering and Molecular Dynamics Simulations. *J. Phys. Chem. B* **2012**, *116*, 14829–14838.
- (58) Heftberger, P. Structure and elasticity of fluid membrane domains. Ph.D. thesis, Graz University of Technology, 2015.
- (59) Páll, S.; Zhmurov, A.; Bauer, P.; Abraham, M.; Lundborg, M.; Gray, A.; Hess, B.; Lindahl, E. Heterogeneous parallelization and acceleration of molecular dynamics simulations in GROMACS. *The Journal of Chemical Physics* **2020**, *153*, 134110.
- (60) Ong, E. E.; Liow, J.-L. The temperature-dependent structure, hydrogen bonding and other related dynamic properties of the standard TIP3P and CHARMM-modified TIP3P water models. *Fluid Phase Equilibria* **2019**, *481*, 55–65.

- (61) Marquardt, D.; Heberle, F. A.; Nickels, J. D.; Pabst, G.; Katsaras, J. On scattered waves and lipid domains: detecting membrane rafts with X-rays and neutrons. *Soft Matter* **2015**, *11*, 9055–9072.
- (62) Doktorova, M.; Kučerka, N.; Kinnun, J. J.; Pan, J.; Marquardt, D.; Scott, H. L.; Venable, R. M.; Pastor, R. W.; Wassall, S. R.; Katsaras, J.; Heberle, F. A. Molecular structure of sphingomyelin in fluid phase bilayers determined by the joint analysis of small-angle neutron and X-ray scattering data. *The Journal of Physical Chemistry B* **2020**, *124*, 5186–5200.
- (63) Pabst, G.; Rappolt, M.; Amenitsch, H.; Laggner, P. Structural information from multilamellar liposomes at full hydration: full q-range fitting with high quality x-ray data. *Physical Review E* **2000**, *62*, 4000.
- (64) Kučerka, N.; Tristram-Nagle, S.; Nagle, J. F. Structure of fully hydrated fluid phase lipid bilayers with monounsaturated chains. *The Journal of membrane biology* **2006**, *208*, 193–202.
- (65) Kučerka, N.; Nieh, M. P.; Katsaras, J. Fluid phase lipid areas and bilayer thicknesses of commonly used phosphatidylcholines as a function of temperature. *Biochim. Biophys. Acta* **2011**, *1808*, 2761–2771.
- (66) Leftin, A.; Molugu, T. R.; Job, C.; Beyer, K.; Brown, M. F. Area per Lipid and Cholesterol Interactions in Membranes from Separated Local-Field ^{13}C NMR Spectroscopy. *Biophys. J.* **2014**, *107*, 2274 – 2286.
- (67) Edholm, O.; Nagle, J. F. Areas of molecules in membranes consisting of mixtures. *Biophysical journal* **2005**, *89*, 1827–1832.
- (68) Javanainen, M.; Melcrová, A.; Magarkar, A.; Jurkiewicz, P.; Hof, M.; Jungwirth, P.; Martinez-Seara, H. Two cations, two mechanisms: interactions of sodium and calcium with zwitterionic lipid membranes. *Chemical Communications* **2017**, *53*, 5380–5383.

- (69) Dickson, C. J.; Walker, R. C.; Gould, I. R. Lipid21: Complex Lipid Membrane Simulations with AMBER. *Journal of chemical theory and computation* **2022**, *18*, 1726–1736.
- (70) Yu, Y.; Kramer, A.; Venable, R. M.; Brooks, B. R.; Klauda, J. B.; Pastor, R. W. CHARMM36 lipid force field with explicit treatment of long-range dispersion: parametrization and validation for phosphatidylethanolamine, phosphatidylglycerol, and ether lipids. *Journal of chemical theory and computation* **2021**, *17*, 1581–1595.
- (71) Yu, Y.; Kramer, A.; Venable, R. M.; Simmonett, A. C.; MacKerell Jr, A. D.; Klauda, J. B.; Pastor, R. W.; Brooks, B. R. Semi-automated optimization of the CHARMM36 lipid force field to include explicit treatment of long-range dispersion. *Journal of chemical theory and computation* **2021**, *17*, 1562–1580.
- (72) Camley, B. A.; Lerner, M. G.; Pastor, R. W.; Brown, F. L. Strong influence of periodic boundary conditions on lateral diffusion in lipid bilayer membranes. *The Journal of chemical physics* **2015**, *143*, 12B604_1.
- (73) Leonard, A. N.; Simmonett, A. C.; Pickard IV, F. C.; Huang, J.; Venable, R. M.; Klauda, J. B.; Brooks, B. R.; Pastor, R. W. Comparison of additive and polarizable models with explicit treatment of long-range Lennard-Jones interactions using alkane simulations. *Journal of Chemical Theory and Computation* **2018**, *14*, 948–958.
- (74) Martinez-Seara, H.; Róg, T.; Karttunen, M.; Vattulainen, I.; Reigada, R. Cholesterol induces specific spatial and orientational order in cholesterol/phospholipid membranes. *PloS one* **2010**, *5*, e11162.

ToDo

P.

1. Update Author list & Affiliations	1
--	---

2. Check that the uses of “model” and “force field” are consistent	2
3. Run Lipid21 for revision?	6
4. Results: volmaps of chains around cholesterol?	9
5. Results: SI: RDFs of POPC chains and CHOL around CHOL	9
6. Unify colours in figures	10
7. Cholesterol order parameters? Into the deviation metric?	16