

NMRlipids III: Lipid-cholesterol interactions in atomistic resolution molecular dynamics simulations

Fernando Favela-Rosales,¹ Peter Heftberger,² Matti Javanainen,^{3,4} Josef Melcr,⁵ O. H. Samuli Ollila,^{5,6,*} Georg Pabst,² and Thomas Piggot⁷

¹*Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14-740, 07000 México D.F., México*

²*University of Graz*

³*Department of Physics, Tampere University of Technology, Tampere, Finland*

⁴*University of Helsinki*

⁵*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic*

⁶*Institute of Biotechnology, University of Helsinki*

⁷*University of Southampton*

(Dated: April 12, 2017)

The quantitative quality of lipid-cholesterol interactions in atomistic resolution models will be determined against NMR and scattering data.

INTRODUCTION

Details of intermolecular interactions determine the phase behaviour and details lateral organization of lipid and bilayers containing cholesterol [1]. **1. Fix Ole's name among the authors of this paper.** Formation of lateral heterogeneities [2], lipid rafts [3] and superlattices [4] in cellular membranes have been suggested to be driven by interactions between cholesterol and lipids. While detailed experimental information of these interactions is relatively sparse, atomistic resolution molecular dynamics simulations have been widely applied to give detailed explanations of lipid bilayer lateral organization and lipid-cholesterol interactions [5, 6]. However, the simulations must reproduce the measured experimental details, such as NMR order parameters and scattering form factors, to be useful in interpretation of molecular details in lipid bilayer mixtures.

Simulations qualitatively reproduce the cholesterol condensation effect, but quantitative comparison to robust experimental data has not been typically done. This is partly due to the lack of availability of systematic experimental data sets for the comparison and partly due to the lack of protocols for validating intermolecular interactions in lipid bilayer simulations against experiments. Scattering experiments are more difficult to interpret for mixed lipid bilayers than for single component bilayers [7–9], thus area per molecule values, one of the main quantity used to compare simulations to experiments, have not been available for lipid cholesterol mixtures. Systematic experimental data set for lipid C-H bond order parameters with different cholesterol concentrations in POPC bilayer has been published only relatively recently [10].

In this work we present the experimental scattering form factor data for POPC-cholesterol mixtures by systematically increasing the cholesterol concentration, scanning the same concentrations that were previously studied with ssNMR [10]. Our goal is to show that the combination of systematically measured C-H bond order parameter and scattering form factor data can be used to validate the quality of lipid-cholesterol intermolecular interactions in MD simulations. MD simulations can be also potentially used to give structural interpreta-

tion for form factors measured from mixed systems, which is a major challenge for current methods [7–9]. The approach should be also applicable for mixed lipid bilayers with other than lipid-cholesterol mixtures.

RESULTS AND DISCUSSION

Structural sampling of molecules in bilayers with cholesterol

Order parameters for C-H bond vectors in lipid bilayer systems, measured with ¹³C or ²H NMR techniques, give indirect information about the structural sampling of individual molecules [11]. The observed increase of acyl chain order parameters with added cholesterol in simulations and experiments can be explained by increased trans conformations in acyl chains [10], which is suggested to play critical role in the phase behaviour of PC-lipid-cholesterol mixtures [1]. Consequently, the correct cholesterol ordering effect is expected to be a necessary condition for a model used to understand lipid-cholesterol phase behaviour.

Acyl chain order parameters for pure 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) bilayer and mixture with 50 mol% of cholesterol from different simulations and experiments are shown in Fig. 1. Phase separation is not observed for this system [10, 12], thus the intermolecular interactions are expected to have maximum effect in equimolar mixture. Experimental acyl chain order parameters are typically well reproduced in the state of the art lipid models for pure lipid bilayers (for review see [11]). This is also mainly observed in the data presented here, except that CHARMM36 slightly overestimates the order parameters and MACROG model underestimates order parameters in the beginning of sn-1 chain. In equimolar mixtures with cholesterol all models overestimate the order parameters. For Berger/Höltje and MACROG models the overestimation is significant, while less severe for CHARMM36 and Slipid models.

2. Is overestimation in CHARMM36 and Slipid significant or not?

3. Lipid14 results from recent article should be added in this figure




FIG. 1: Order parameters from simulations and experiments for acyl chains of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC).

The order parameter changes as a function of cholesterol for each segment are shown in Fig. 2 (currently only sn-1).

4.This figure should be finished and similar for sn-2 should be done as well

5.Also experimental cholesterol order parameters are available. Maybe these should be calculated from simulations as well.

Lipid bilayer dimensions and density profiles as a function of cholesterol

Cholesterol induced changes in lipid packing, thickness and density profiles are suggested to be relevant, for example, for membrane protein interactions and cellular membrane lateral organization [?]. Experimental and simulation studies show that the cholesterol induces the so called "condensing effect", i.e. acyl chain ordering increases the membrane thickness and reduces the area per molecule in lipid bilayers [?]. This is

demonstrated from simulations in Fig. 3, showing the area per PC headgroups and per total number of molecules as a function of cholesterol concentration from different models. All models show reduction of area per total amount of molecules, partly due to the smaller area covered by the cholesterol than lipids, but also because lipids ordered by cholesterol require less space. Due to the latter effect, the area per PC headgroup does not essentially increase up to the addition of ~15 mol% of cholesterol.

6.Membrane thickness to be plotted as well.

Bilayer dimensions are indirectly experimentally accessible by scattering methods and have been widely used to validate lipid bilayer simulations [11?]. While sophisticated models are applied to extract area per molecule and thickness for single component lipid bilayers, the results for multicomponent systems are more difficult to interpret [7? -9]. In such case the validation can be done by comparing directly measurable




FIG. 2: Order parameter changes from simulations and experiments for each segment in sn-1 chain of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) as a function of cholesterol concentration.




FIG. 3: Area per molecules calculated from different simulation models as a function of cholesterol concentration. The solid symbols are area per total amount of molecules (chol+PC) and the empty symbols area per PC headgroups. Top figure shows absolute values and bottom figure shows changes respect to pure lipid system.

form factors between simulations and experiments. If simulation reproduces the form factor, it can be also used to interpret the experiments. The form factors from different simulation models with different cholesterol concentrations are compared to experimental data in Fig. 4.

7. Discussion to be finished when the data is solid.

CONCLUSIONS

Cholesterol ordering effect is overestimated in Berger/Holtje and MacRog models. Slight overestimation is observed also in CHARMM36 and Slipid models, but more careful analysis is required to conclude if this is significant or not.




FIG. 4: Form factors from simulations and experiments.

8. Form factor calculation method should be double checked.

9. Experimental form factor amplitudes are not scaled to match with simulations,
as done usually

10. Not all experimental and simulation data is here.

METHODS

X-ray scattering experiments

SAXS data on POPC multilamellar vesicles (MLVs) at various cholesterol concentrations has been measured. Data have been obtained at the EMBL BioSAXS beamline (Hamburg) using 20 keV photons, $T = 27^\circ\text{C}$. Data were analyzed in terms of the SDP-GAP model described in Heftberger et al., J. Appl. Cryst. 2013 and Heftberger et al. Biophys. J. 2015. Data from MLVs are a convolute of structure factor (the crystalline lattice) and form factor. By fitting the scattered intensity data we obtain both contributions. Here we posted only form factors (ASCII format). For information on the quality of the fit we also give plots of the fitted intensity data. The electron den-

sity profile has been modelled in terms of the SDP model (see papers by Kucerka and coworkers), that is volume distribution functions are modelled by individual Gaussians or error functions. Cholesterol is also accounted for by two Gaussians. This model has been proposed by Jianjun Pan (USF, Tampa, FL), but is to the best of our knowledge not published (see also PhD Thesis by Peter Heftberger). Additional figures show the volume distribution functions and the resulting electron density profiles.

Authors to consult and potentially include in publications using this data: Peter Heftberger (peter.heftberger@gmx.at), Georg Pabst (georg.pabst@uni-graz.at)

Molecular dynamics simulations

Simulated systems are listed in Table I and the full simulation details are given in references or in supplementary material.

TABLE I: Simulated lipid bilayers containing cholesterol. The simulation file data sets marked with * include also part of the trajectory. ^a The number of lipid molecules ^b The number of cholesterol molecules ^c Cholesterol concentration (mol%) ^d The number of water molecules ^e Simulation temperature ^f The total simulation time ^g Time frames used in the analysis ^h Reference link for the downloadable simulation files ⁱ Reference for the full simulation details

| Force field | lipid | ^a N _l | ^b N _{chol} | ^c C _{CHOL} | ^d N _w | ^e T (K) | ^f t _{sim} (ns) | ^g t _{anal} (ns) | ^h Files | ⁱ Details |
|-------------------------|-------|-----------------------------|--------------------------------|--------------------------------|-----------------------------|--------------------|------------------------------------|-------------------------------------|--------------------|----------------------|
| Berger-POPC-07 [13] | POPC | 128 | 0 | 0% | 7290 | 298 | 270 | 240 | [14]* | [15] |
| Höltje-CHOL-13 [10, 16] | POPC | 120 | 8 | 6% | 7290 | 298 | 100 | 80 | [17]* | [10] |
| | POPC | 110 | 18 | 14% | 8481 | 298 | 100 | 80 | [18]* | [10] |
| | POPC | 84 | 44 | 34% | 6794 | 298 | 100 | 80 | [19]* | [10] |
| | POPC | 64 | 64 | 50% | 10314 | 298 | 100 | 80 | [20]* | [10] |
| | POPC | 50 | 78 | 61% | 5782 | 298 | 100 | 80 | [21]* | [10] |
| CHARMM36[22, 23] | POPC | 200 | 0 | 0% | 9000 | 310 | ? | 100 | [24]* | SI |
| | POPC | 200 | 22 | 10% | 9000 | 310 | ? | 100 | [24]* | SI |
| | POPC | 200 | 50 | 20% | 9000 | 310 | ? | 100 | [24]* | SI |
| | POPC | 200 | 86 | 30% | 9000 | 310 | ? | 100 | [24]* | SI |
| | POPC | 200 | 134 | 40% | 15030 | 310 | 109 | 100 | [25]* | SI |
| | POPC | 200 | 200 | 50% | 18000 | 310 | 109 | 100 | [25]* | SI |
| Slipids[26–28] | POPC | 200 | 0 | 0% | ? | 310 | ? | 100 | [29]* | SI |
| | POPC | 512 | 0 | 0% | 23943 | 310 | 170 | 100 | [30]* | SI |
| | POPC | 200 | 22 | 10% | ? | 310 | ? | 100 | [29]* | SI |
| | POPC | 200 | 50 | 20% | ? | 310 | ? | 100 | [29]* | SI |
| | POPC | 200 | 86 | 30% | ? | 310 | ? | 100 | [29]* | SI |
| | POPC | 358 | 154 | 30% | 21183 | 298 | 170 | 100 | [31]* | SI |
| | POPC | 200 | 134 | 40% | ? | 310 | 109 | 100 | [32]* | SI |
| | POPC | 200 | 200 | 50% | ? | 310 | 109 | 100 | [32]* | SI |
| | POPC | 256 | 256 | 50% | 20334 | 298 | 170 | 100 | [33]* | SI |
| MacRog[34] | POPC | 128 | 0 | 0% | 6400 | 310 | 400 | 200 | [35]* | [36] |
| | POPC | 114 | 14 | 11% | 6400 | 310 | 400 | 200 | [35]* | [36] |
| | POPC | 72 | 56 | 44% | 6400 | 310 | 400 | 200 | [35]* | [36] |
| | POPC | 64 | 64 | 50% | 6400 | 310 | 400 | 200 | [35]* | [36] |
| | POPC | 56 | 72 | 56% | 6400 | 310 | 400 | 200 | [35]* | [36] |

SUPPLEMENTARY INFORMATION

CHARMM36 results from different simulation packages

The results from CHARMM36 model for lipid bilayers from different simulation packages have been reported to give different results in the literature [37, 38]. The results are mainly dependent on different Lennard-Jones cut-off settings, but all the details are not quite understood. In this work we use the results from Gromacs 5 with settings suggested to be optimal by Gromacs webpage. **11.List these settings here or in SI.** We also compared the results from Gromacs 5 with these settings to the results simulated with NAMD, OpenMM and literature values. Based on comparison shown in Fig. 5, we conclude that Gromacs 5 with settings suggested in webpage gives consistent results with the literature and other simulation packages. Thus these results are used in the main body of the

paper. However, order parameters are slightly overestimated respected to the experiments also with these settings.

12.Should be extend this discussion based on discussion at <https://github.com/NMRLipids/NmrLipidsCholXray/issues/4> ?

* samuli.ollila@helsinki.fi

- [1] J. H. Ipsen, G. Karlström, O. Mourtsen, H. Wennerström, and M. Zuckermann, *Biochim. Biophys. Acta* **905**, 162 (1987).
- [2] P. K. Kinnunen, *Chemistry and Physics of Lipids* **57**, 375 (1991).
- [3] K. Simons and E. Ikonen, *Nature* **387**, 569 (1997).
- [4] P. Somerharju, J. A. Virtanen, K. H. Cheng, and M. Hermanson, *Biochim. Biophys. Acta - Biomembranes* **1788**, 12 (2009).
- [5] T. Róg and I. Vattulainen, *Chem. Phys. Lipids* **184**, 82 (2014).
- [6] A. J. Sodt, M. L. Sandar, K. Gawrisch, R. W. Pastor, and E. Lyman, *J. Am. Chem. Soc.* **136**, 725 (2014).
- [7] J. Pan, X. Cheng, F. A. Heberle, B. Mostofian, N. Kučerka,




FIG. 5: Results for CHARMM36 model [22] from different simulation packages. Discussion going on at <https://github.com/NMRLipids/NmrLipidsCholXray/issues/4>.

P. Drazba, and J. Katsaras, *J. Phys. Chem. B* **116**, 14829 (2012).
 [8] P. Heftberger, B. Kollmitzer, A. Rieder, H. Amenitsch, and G. Pabst, *Biophys. J.* **108**, 854 (2015), ISSN 0006-3495.
 [9] D. Marquardt, F. A. Heberle, J. D. Nickels, G. Pabst, and J. Katsaras, *Soft Matter* **11**, 9055 (2015).
 [10] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
 [11] O. S. Ollila and G. Pabst, *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1858**, 2512 (2016), biosimulations of lipid membranes coupled to experiments.
 [12] I. Ionova, V. Livshits, and D. Marsh, *Biophysical Journal* **102**, 1856 (2012).
 [13] S. Ollila, M. T. Hyvönen, and I. Vattulainen, *J. Phys. Chem. B* **111**, 3139 (2007).
 [14] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC bilayer (Berger model delivered by Tieleman, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13279>.
 [15] T. M. Ferreira, O. H. S. Ollila, R. Pigliapochi, A. P. Dabkowska,

and D. Topgaard, *J. Chem. Phys.* **142**, 044905 (2015).
 [16] M. Höltje, T. Förster, B. Brandt, T. Engels, W. von Rybinski, and H.-D. Höltje, *Biochim. Biophys. Acta* **1511**, 156 (2001).
 [17] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (7 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13282>.
 [18] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (15 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13281>.
 [19] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (34 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13283>.
 [20] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (50 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13285>.
 [21] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (60 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13286>.
 [22] J. B. Klauda, R. M. Venable, J. A. Freites, J. W. O'Connor, D. J. Tobias, C. Mondragon-Ramirez, I. Vorobyov, A. D. MacKerell Jr, and R. W. Pastor, *J. Phys. Chem. B* **114**, 7830 (2010).
 [23] J. B. Lim, B. Rogaski, and J. B. Klauda, *J. Phys. Chem. B* **116**, 203 (2012).
 [24] M. Javanainen, *POPC with 0, 10, 20, and 30 mol-310 K. Charmm36 force field.* (2016), URL <https://doi.org/10.5281/zenodo.159759>.
 [25] M. Javanainen, *POPC with 40 and 50 mol-Slipids force field.* (2016), URL <https://doi.org/10.5281/zenodo.154346>.
 [26] J. P. M. Jämbeck and A. P. Lyubartsev, *J. Chem. Theory Comput.* **8**, 2938 (2012).
 [27] J. P. M. Jämbeck and A. P. Lyubartsev, *J. Phys. Chem. B* **116**, 3164 (2012).
 [28] J. P. M. Jämbeck and A. P. Lyubartsev, *Journal of Chemical Theory and Computation* **9**, 774 (2013).
 [29] M. Javanainen, *POPC with 0, 10, 20, and 30 mol-310 K. Slipids force field.* (2016), URL <https://doi.org/10.5281/zenodo.60607>.
 [30] F. Favela-Rosales, *MD simulation trajectory of a lipid bilayer: Pure POPC in water. SLIPIDS, Gromacs 4.6.3. 2016.* (2016), URL <https://doi.org/10.5281/zenodo.166034>.
 [31] F. Favela-Rosales, *MD simulation trajectory of a lipid bilayer: 70/30 mol% POPC/Cholesterol. SLIPIDS, Gromacs 4.6.3. 2016.* (2016), URL <https://doi.org/10.5281/zenodo.62026>.
 [32] M. Javanainen, *POPC with 40 and 50 mol-Slipids force field.* (2016), URL <https://doi.org/10.5281/zenodo.154346>.
 [33] F. Favela-Rosales, *MD simulation trajectory of a lipid bilayer: 50/50 mol% POPC/Cholesterol. SLIPIDS, Gromacs 4.6.3. 2016.* (2016), URL <https://doi.org/10.5281/zenodo.159434>.
 [34] W. Kulig, M. Pasenkiewicz-Gierula, and T. Róg, *Chem. Phys. Lipids* **195**, 12 (2016).

- [35] M. Javanainen, *POPC/Cholesterol @ 310K. 0, 10, 40, 50 and 60 mol-cholesterol. Model by Maciejewski and Róg* (2015), URL {<http://dx.doi.org/10.5281/zenodo.13877>}.
- [36] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduč, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., *J. Phys. Chem. B* **119**, 15075 (2015).
- [37] T. J. Piggot, Á. Piñeiro, and S. Khalid, *J. Chem. Theory Comput.* **8**, 4593 (2012).
- [38] J. Lee, X. Cheng, J. M. Swails, M. S. Yeom, P. K. Eastman, J. A. Lemkul, S. Wei, J. Buckner, J. C. Jeong, Y. Qi, et al., *Journal of Chemical Theory and Computation* **12**, 405 (2016).

ToDo

- | | | |
|---|----|---|
| 1. Fix Ole's name among the authors of this paper. . . | P. | 1 |
| 2. Is overestimation in CHARMM36 and Slipid significant or not? | 1 | |
| 3. Lipid14 results from recent article should be added in this figure | 1 | |
| 4. This figure should be finished and similar for sn-2 should be done as well | 2 | |
| 5. Also experimental cholesterol order parameters are available. Maybe these should be calculated from simulations as well. | 2 | |
| 6. Membrane thickness to be plotted as well. | 2 | |
| 7. Discussion to be finished when the data is solid. . . | 3 | |
| 8. Form factor calculation method should be double checked. | 4 | |
| 9. Experimental form factor amplitudes are not scaled to match with simulations, as done usually | 4 | |
| 10. Not all experimental and simulation data is here. . . | 4 | |
| 11. List these settings here or in SI. | 5 | |
| 12. Should be extend this discussion based on discussion at https://github.com/NMRLipids/NmrLipidsCholXray/issues/4 ? | 5 | |