

Quantitative quality of lipid-cholesterol interactions in atomistic resolution molecular dynamics simulations

Peter Heftberger,¹ O. H. Samuli Ollila,^{2,*} and Georg Pabst¹

¹*University of Graz*

²*Aalto University*

(Dated: August 26, 2016)

The quantitative quality of lipid-cholesterol interactions in atomistic resolution models will be determined against NMR and scattering data.

INTRODUCTION

The quantitative quality of lipid-cholesterol interactions in atomistic resolution models will be determined against NMR and scattering data.

The order parameter changes as a function of cholesterol for each segment are shown in Fig. 2 (currently only sn-1).

COMPARISON OF FORM FACTORS BETWEEN EXPERIMENTS AND SIMULATIONS

The form factors calculated from different simulations with different cholesterol content are shown in Fig. 3.

METHODS

X-ray scattering experiments

This folder contains SAXS data on POPC multilamellar vesicles (MLVs) at various cholesterol concentrations. Data have been obtained at the EMBL BioSAXS beamline (Hamburg) using 20 keV photons, $T = 27^\circ\text{C}$. Data were analyzed in terms of the SDP-GAP model described in Heftberger et al., J. Appl. Cryst. 2013 and Heftberger et al. Biophys. J. 2015. Data from MLVs are a convolute of structure factor (the crystalline lattice) and form factor. By fitting the scattered intensity data we obtain both contributions. Here we posted only form factors (ASCII format). For information on the quality of the fit we also give plots of the fitted intensity data. The electron density profile has been modelled in terms of the SDP model (see papers by Kucerka and coworkers), that is volume distribution functions are modelled by individual Gaussians or error functions. Cholesterol is also accounted for by two Gaussians. This model has been proposed by Jianjun Pan (USF, Tampa, FL), but is to the best of our knowledge not published (see also PhD Thesis by Peter Heftberger). Additional figures show the volume distribution functions and the resulting electron density profiles.

Authors to consult and potentially include in publications using this data: Peter Heftberger (peter.heftberger@gmx.at), Georg Pabst (georg.pabst@uni-graz.at)

CONCLUSIONS

Molecular dynamics simulations

COMPARISON OF ACYL CHAIN ORDER PARAMETERS BETWEEN EXPERIMENTS AND SIMULATIONS

Order parameters from simulations and experiments for acyl chains of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) are shown in Fig. 1.

TABLE I: Simulated lipid bilayers containing cholesterol. The simulation file data sets marked with * include also part of the trajectory. ^a The number of lipid molecules ^b The number of cholesterol molecules ^c Cholesterol concentration (mol%) ^d The number of water molecules ^e Simulation temperature ^f The total simulation time ^g Time frames used in the analysis ^h Reference link for the downloadable simulation files ⁱ Reference for the full simulation details

Force field	lipid	^a N _l	^b N _{chol}	^c C _{CHOL}	^d N _w	^e T (K)	^f t _{sim} (ns)	^g t _{anal} (ns)	^h Files	ⁱ Details
Berger-POPC-07 [1]	POPC	128	0	0%	7290	298	270	240	[2]*	[3]
/Höltje-CHOL-13 [4, 5]										
	POPC	120	8	6%	7290	298	100	80	[6]*	[5]
	POPC	110	18	14%	8481	298	100	80	[7]*	[5]
	POPC	84	44	34%	6794	298	100	80	[8]*	[5]
	POPC	64	64	50%	10314	298	100	80	[9]*	[5]
	POPC	50	78	61%	5782	298	100	80	[10]*	[5]
CHARMM36[11, 12]	POPC	128	0	0%	5120	303	150	100	[13]*	[14]
	POPC	100	24	19%	4960	303	200	100	[15]*	[14]
	POPC	80	80	50%	4496	303	200	100	[16]*	[14]
MacRog[17]	POPC	128	0	0%	6400	310	400	200	[18]*	[14]
	POPC	114	14	11%	6400	310	400	200	[18]*	[14]
	POPC	72	56	44%	6400	310	400	200	[18]*	[14]
	POPC	64	64	50%	6400	310	400	200	[18]*	[14]
	POPC	56	72	56%	6400	310	400	200	[18]*	[14]

* samuli.ollila@aalto.fi

- [1] S. Ollila, M. T. Hyvönen, and I. Vattulainen, *J. Phys. Chem. B* **111**, 3139 (2007).
- [2] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC bilayer (Berger model delivered by Tieleman, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13279>.
- [3] T. M. Ferreira, O. H. S. Ollila, R. Pigliapochi, A. P. Dabkowska, and D. Topgaard, *J. Chem. Phys.* **142**, 044905 (2015).
- [4] M. Höltje, T. Förster, B. Brandt, T. Engels, W. von Rybinski, and H.-D. Höltje, *Biochim. Biophys. Acta* **1511**, 156 (2001).
- [5] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
- [6] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (7 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13282>.
- [7] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (15 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13281>.
- [8] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (34 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13283>.
- [9] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (50 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13285>.
- [10] O. H. S. Ollila, T. Ferreira, and D. Topgaard, *MD simulation trajectory and related files for POPC/cholesterol (60 mol%) bilayer (Berger model delivered by Tieleman, modified Höltje, Gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.13286>.
- [11] J. B. Klauda, R. M. Venable, J. A. Freites, J. W. O'Connor, D. J. Tobias, C. Mondragon-Ramirez, I. Vorobyov, A. D. MacKerell Jr, and R. W. Pastor, *J. Phys. Chem. B* **114**, 7830 (2010).
- [12] J. B. Lim, B. Rogaski, and J. B. Klauda, *J. Phys. Chem. B* **116**, 203 (2012).
- [13] H. Santuz, *MD simulation trajectory and related files for POPC bilayer (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.14066>.
- [14] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduć, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., *J. Phys. Chem. B* **119**, 15075 (2015).
- [15] H. Santuz, *MD simulation trajectory for POPC/20% Chol bilayer (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.14067>.
- [16] H. Santuz, *MD simulation trajectory for POPC/50% Chol bilayer (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.14068>.
- [17] W. Kulig, M. Pasenkiewicz-Gierula, and T. Róg, *Chem. Phys. Lipids* **195**, 12 (2016).
- [18] M. Javanainen, *POPC/Cholesterol @ 310K. 0, 10, 40, 50 and 60 mol-cholesterol. Model by Maciejewski and Róg* (2015), URL <http://dx.doi.org/10.5281/zenodo.13877>.

ToDo

P.




FIG. 1: Order parameters from simulations and experiments for acyl chains of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC).

1. Why are the order parameters of CHARMM36 too large compared to simulations even without cholesterol? Discussion in

<https://github.com/NMRLipids/NmrLipidsCholXray/issues/4>

2. Why there is decrease in order parameters towards beginning of acyl chain in MacRog model without cholesterol?

3. Do the results suggest that condensation effect is too strong in MacRog and Berger models? Discussion in <https://github.com/NMRLipids/NmrLipidsCholXray/issues/5>

- | | | |
|---|--|---|
| 1. Why are the order parameters of CHARMM36 too large compared to simulations even without cholesterol? Discussion in https://github.com/NMRLipids/NmrLipidsCholXray/issues/4 | 4. Form factor calculation method should be double checked. | 5 |
| 2. Why there is decrease in order parameters towards beginning of acyl chain in MacRog model without cholesterol? | 5. Experimental form factor amplitudes are not scaled to match with simulations, as done usually | 5 |
| 3. Do the results suggest that condensation effect is too strong in MacRog and Berger models? Discussion in https://github.com/NMRLipids/NmrLipidsCholXray/issues/5 | 6. Not all experimental and simulation data is here. | 5 |




FIG. 2: Order parameter changes from simulations and experiments for each segment in sn-1 chain of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) as a function of cholesterol concentration.




FIG. 3: Form factors from simulations and experiments.

4. Form factor calculation method should be double checked.

5. Experimental form factor amplitudes are not scaled to match with simulations,
as done usually

6. Not all experimental and simulation data is here.