

Cite this: DOI: 10.1039/xxxxxxxxxx

Molecular electrometer and binding of cations to phospholipid bilayers[†]

Andrea Catte,^{a,‡} Mykhailo Giryh,^b Matti Javanainen,^{c,d} Claire Loison,^e Josef Melcr,^f Markus S. Miettinen,^{g,h} Luca Monticelli,ⁱ Jukka Määttä,^j Vasily S. Oganessian,^a O. H. Samuli Ollila,^{*b} Joona Tynkkynen,^c and Sergey Vilov,^e

Received Date
Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

Despite the vast amount of experimental and theoretical studies on the binding affinity of cations — especially the biologically relevant Na^+ and Ca^{2+} — for phospholipid bilayers, there is no consensus in the literature. Here we show that by interpreting changes in the choline headgroup order parameters according to the 'molecular electrometer' concept [Seelig *et al.*, *Biochemistry*, 1987, **26**, 7535], one can directly compare the ion binding affinities between simulations and experiments. Our findings strongly support the view that in contrast to Ca^{2+} and other multivalent ions, Na^+ and other monovalent ions (except Li^+) do not specifically bind to phosphatidylcholine lipid bilayers at sub-molar concentrations. However, the Na^+ binding affinity was overestimated by several molecular dynamics simulation models, resulting in artificially positively charged bilayers and exaggerated structural effects in the lipid headgroups. While qualitatively correct headgroup order parameter response was observed with Ca^{2+} binding in all the tested models, no model had sufficient quantitative accuracy to interpret the Ca^{2+} :lipid stoichiometry or the induced atomistic resolution structural changes. All scientific contributions to this open collaboration work were made publicly, using `nmrlipids.blogspot.fi` as the main communication platform.

1 Introduction

Due to its high physiological importance — nerve cell signalling being the prime example — interaction of cations with phospholipid membranes has been widely studied via theory, simulations, and experiments. The relative ion binding affinities are generally agreed to follow the Hofmeister series^{1–9}, however, consen-

sus on the quantitative affinities is currently lacking. Until 1990, the consensus (documented in two extensive reviews^{2,3}) was that while multivalent cations interact significantly with phospholipid bilayers, for monovalent cations (with the exception of Li^+) the interactions are weak. This conclusion has since been strengthened by further studies showing that bilayer properties remain unaltered upon the addition of sub-molar concentrations of monovalent salt^{4,10,11}. Since 2000, however, another view has emerged, suggesting much stronger interactions between phospholipids and monovalent cations, and strong Na^+ binding in particular^{6–9,12–18}.

The pre-2000 view has the experimental support that (in contrast to the significant effects caused by any multivalent cations) sub-molar concentrations of NaCl have a negligible effect on phospholipid infrared spectra⁴, area per molecule¹⁰, dipole potential²⁰, lateral diffusion¹¹, and choline head group order parameters¹⁹; in addition, the water sorption isotherm of a NaCl–phospholipid system is highly similar to that of a pure NaCl solution — indicating that the ion–lipid interaction is very weak⁴.

The post-2000 'strong binding' view rests on experimental and above all simulational findings. At sub-molar NaCl concentrations, the rotational and translational dynamics of membrane-embedded fluorescent probes decreased^{7,9,12}, and atomic force microscopy (AFM) experiments showed changes in bilayer hard-

^a School of Chemistry, University of East Anglia, Norwich, NR4 7TJ, United Kingdom
^b Department of Neuroscience and Biomedical Engineering, Aalto University, Espoo, Finland
^c Tampere University of Technology, Tampere, Finland
^d University of Helsinki, Helsinki, Finland
^e Univ Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622, LYON, France
^f Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, Flemingovo nám. 2, 16610 Prague 6, Czech Republic, Charles University in Prague, Faculty of Mathematics and Physics, Ke Karlovu 3, 121 16 Prague 2, Czech Republic
^g Fachbereich Physik, Freie Universität Berlin, Berlin, Germany
^h Max Planck Institute of Colloids and Interfaces, Department of Theory and Bio-Systems, Potsdam, Germany
ⁱ Institut de Biologie et Chimie des Protéines (IBCP), CNRS UMR 5086, Lyon, France
^j Aalto University, Espoo, Finland

* Author to whom correspondence may be addressed. E-mail: samuli.ollila@aalto.fi.

[†] Electronic Supplementary Information (ESI) available: 5 figures, detailed technical discussion and simulation details. See DOI: 10.1039/b000000x/

[‡] The authors are listed in alphabetical order.

ness^{14–18}; in atomistic molecular dynamics (MD) simulations, phospholipid bilayers consistently bound Na⁺, although the binding strength depended on the model used^{12,13,22,24?–27}.

Some observables have been interpreted in favour of both views. For example, as the effect of monovalent ions (except Li⁺) on the phase transition temperature is tiny (compared to the effect of multivalent ions), it was initially interpreted as an indication that only multivalent ions and Li⁺ specifically bind to phospholipid bilayers²; however, such a small effect in calorimetric measurements was later interpreted to indicate that also Na⁺ binds^{8,12}. Similarly, the lack of significant positive electrophoretic mobility of phosphatidylcholine (PC) vesicles in the presence of NaCl (again in contrast to multivalent ions and Li⁺) suggested weak binding of Na⁺^{1,8,14,15,21}; however, these data were also explained by a countering effect of the Cl[−] ions^{22,23}. Furthermore, to reduce the area per lipid in scattering experiments, molar concentrations of NaCl were required¹⁰, indicating weak ion–lipid interaction; in MD simulations, however, already orders of magnitude lower concentrations resulted in Na⁺ binding and a clear reduction of area per lipid^{12?}. Finally, lipid lateral diffusion was unaltered by NaCl in noninvasive NMR experiments¹¹; however, as it was reduced upon Na⁺ binding in simulations, the reduced lateral diffusion of fluorescent probes^{7,9,12} has been interpreted to support the post-2000 ‘strong binding’ view.

In this paper, we set out to solve the apparent contradictions between the pre-2000 and post-2000 views. To this end, we employ the ‘molecular electrometer’ concept, according to which the changes in the C–H order parameters of the α and β carbons in the phospholipid head group (see Fig. 1) can be used to measure the ion affinity for a PC lipid bilayer^{19,28?–30}. As the order parameters can be accurately measured in experiments and directly compared to simulations[?], applying the molecular electrometer as a function of cation concentration allows the comparison of binding affinity between simulations and experiments. In addition to demonstrating the usefulness of this general concept, we show that the response of the α and β order parameters to penetrating cations is qualitatively correct in MD simulations, but that in several models the affinity of Na⁺ for PC bilayers is grossly overestimated. Moreover, we show that the accuracy of lipid–Ca²⁺ interactions in current models is not enough for atomistic resolution interpretation of NMR experiments.

This work was done as an Open Collaboration at nmrlipids.blogspot.fi; all the related files¹⁰⁵ and almost all the simulation data (<https://zenodo.org/collection/user-nmrlipids>) are openly available.

2 Results and Discussion

2.1 Background: Molecular electrometer in experiments

The basis for the molecular electrometer is the experimental observation that binding of any charged objects (ions, peptides, anesthetics, amphiphiles) on a PC bilayer interface induced systematic changes in the choline α and β segment C–H order parameters^{19,28?–30}. Being systematic, these changes could be employed for determining the binding affinities of the charged objects in question. Originally the molecular electrometer was



Fig. 1 Chemical structure of 1-palmitoyl-2-oleoylphosphatidylcholine (POPC), and the definition of γ , β , α , g_1 , g_2 and g_3 segments.

devised for cations^{19,28?}, but further experimental quantification with various positively and negatively charged molecules showed that the choline order parameters S_{CH}^{α} and S_{CH}^{β} in general vary linearly with small amount of bound charge per lipid^{28?–30}. Let now $S_{CH}^i(0)$, where i refers to either α or β , denote the order parameter in the absence of bound charge; the empirically observed linear relation can then be written as[?]

$$\Delta S_{CH}^i = S_{CH}^i(X^{\pm}) - S_{CH}^i(0) = \frac{4m_i}{3\chi} X^{\pm}. \quad (1)$$

Here X^{\pm} is the amount of bound charge per lipid, m_i an empirical constant depending on the valency and position of bound charge, and the value of the quadrupole coupling constant $\chi \approx 167$ kHz.

With bound positive charge, the absolute value of the β segment order parameter increases and the α segment order parameter decreases (and *vice versa* for negative charge)^{19,28?–30}. However, as $S_{CH}^{\beta}(0) < 0$ while $S_{CH}^{\alpha}(0) > 0$ ^{32–34}, both ΔS_{CH}^{β} and ΔS_{CH}^{α} in fact decrease with bound positive charge (and increase with bound negative charge). Consequently, values of m_i are negative for bound positive charges; for Ca²⁺ binding to POPC bilayer (in the presence of 100 mM NaCl), combination of atomic absorption spectra and ²H NMR experiments gave $m_{\alpha} = -20.5$ and $m_{\beta} = -10.0$ ²⁸. This decrease can be rationalised by electrostatically induced tilting of the choline P–N dipole^{29?,30} — also seen in simulations^{25?–27} — and is in line with the order parameter increase related to the P–N vector tilting more parallel to the membrane plane seen with decreasing hydration levels³⁵.

Quantification of ΔS_{CH}^{α} and ΔS_{CH}^{β} for a wide range of different cations (aqueous cations, cationic peptides, cationic anesthetics) has revealed that $\Delta S_{CH}^{\beta}/\Delta S_{CH}^{\alpha} \approx 0.5$ ^{??}. More specifically, the relation $\Delta S_{CH}^{\beta} = 0.43\Delta S_{CH}^{\alpha}$ was found to hold for DPPC bilayers at various CaCl₂ concentrations¹⁹.

2.2 Molecular electrometer in MD simulations

The black curves in Fig. 2 show how the headgroup order parameters for DPPC and POPC bilayers change in H² NMR experiments as a function of salt solution concentration^{19,28}: Only mi-




Fig. 2 Changes in the PC lipid headgroup β (top row) and α (bottom) segment order parameters in response to NaCl (left column) or CaCl_2 (right) salt solution concentration increase. Comparison between simulations (Table 1) and experiments (DPPCs from Ref. 19, POPC from Ref. 28). The signs of the experimental values, from experiments without ions^{32–34}, can be assumed unchanged at these salt concentrations²⁸⁷. We stress that none of the models reproduces the order parameters without salt within experimental error, indicating structural inaccuracies of varying severity in all of them³⁵. Note that the relatively large drop in CHARMM36 at 450 mM CaCl_2 arose from more equilibrated binding due to a very long simulation time, see ESI[†].

nor changes are seen as a function of $[\text{NaCl}]$, but the effect of $[\text{CaCl}_2]$ is an order of magnitude larger. Thus, according to the molecular electrometer, the monovalent Na^+ ions have negligible affinity for PC lipid bilayers at concentrations up to 1 M, while binding of Ca^{2+} ions at the same concentration is significant^{19,28}.

Figure 2 also reports order parameter changes calculated from MD simulations of DPPC and POPC lipid bilayers as a function of NaCl or CaCl_2 initial concentrations in solution (for details of the simulated systems see Table 1 and ESI[†]). Note that although none of these MD models reproduces within experimental uncertainty the order parameters for a pure PC bilayer without ions (Fig. 2 in Ref. 35), which indicates structural inaccuracies of varying severity in all models³⁵, all the models qualitatively reproduce the experimentally observed headgroup order parameter increase with dehydration³⁵. Similarly here (Fig. 2) the presence of cations led to the decrease of S_{CH}^{α} and S_{CH}^{β} , in qualitative agreement with experiments. The changes were, however, overestimated by most models, which according to the molecular electrometer indicates overbinding of cations in most MD simulations.

While the molecular electrometer is well established in experi-

ments (see Sec. 2.1 above), it is not *a priori* clear that it works in simulations. The overestimated order parameter decrease could, in principle, arise from an exaggerated response of the choline headgroups to the binding cations, instead of overbinding. Therefore, to evaluate the usability of the molecular electrometer in MD simulations, we analysed the relation between cation binding and choline order parameter decrease in simulations.

According to the molecular electrometer, the order parameter changes are linearly proportional to the amount of bound cations (Eq. (1)). Figure 3 shows this proportionality in MD simulations (see ESI[†] for the definition of bound ions); in keeping with the molecular electrometer, a roughly linear correlation between bound cation charge and order parameter change was found in all the eight models. Note that quantitative comparison of the proportionality constants (i.e. slopes in Fig. 3) between different models and experimental slopes ($m_{\alpha} = -20.5$ and $m_{\beta} = -10.0$ for Ca^{2+} binding in DPPC bilayer in the presence of 100mM NaCl²⁸) is not straightforward since the simulation slopes depend on the definition used for bound ions (see ESI[†]).

We note that the quantitative comparison of order parameter




Fig. 3 Change of order parameters (from salt-free solution) of the β and α segments, ΔS_{CH}^{β} and ΔS_{CH}^{α} , as a function of bound cation charge. Eight MD simulation models compared; the two lines per model denote to the two hydrogens per carbon. The order parameters as well as the bound charge calculated separately for each leaflet; cations residing between the bilayer centre and the density maximum of Phosphorus considered bound; error bars (shaded) show standard error of mean over lipids.

changes in response to bound charge should be more straightforward for systems with charged amphiphiles fully associated in the bilayer, as the amount of bound charge is then explicitly known in both simulations and experiments. In such a comparison between experiments^{30?} and previously published Berger-model-based simulations[?], we could not rule out overestimation of order parameter response to bound cations (slopes m_{α} and m_{β}), see ESI[†]. This might, in principle, explain the overestimated order parameter response of the Berger model to CaCl_2 , but not to NaCl (see discussion in ESI[†]). Since simulation data with charged amphiphiles are not available for other models, an extended comparison with different models is left for further studies.

Figure 3 shows that the decrease of order parameters clearly correlated with the amount of bound cations in simulations. This is also evident from Fig. 4, which shows the Na^+ density profiles of the MD models ordered according to the order parameter change (in Fig. 2) from the smallest (top) to the largest (bottom). The general trend in the figure is that the Na^+ density peaks are larger for models with larger changes in order parameters, in line with the observed correlation between cation binding and order parameter decrease in Fig. 3.




Fig. 4 Na^+ (solid line) and Cl^- (dashed) distributions along the lipid bilayer normal from MD simulations at several NaCl concentrations. The eight MD models are ordered according to their strength of order parameter change in response to NaCl (Fig. 2) from the weakest (top panel) to the strongest (bottom). The light green vertical lines indicate the locations of the Phosphorus maxima, used to define bound cations in Fig. 3.




Fig. 5 Relation between $\Delta S_{\text{CH}}^{\beta}$ and $\Delta S_{\text{CH}}^{\alpha}$ from experiments¹⁹ and different simulation models. Solid line is $\Delta S_{\text{CH}}^{\beta} = 0.43\Delta S_{\text{CH}}^{\alpha}$ determined for DPPC bilayer from ^2H NMR experiment with various CaCl_2 concentrations¹⁹.

Figure 5 compares the relation between $\Delta S_{\text{CH}}^{\beta}$ and $\Delta S_{\text{CH}}^{\alpha}$ in experiments¹⁹ and in MD models. Only Lipid14 gave $\Delta S_{\text{CH}}^{\beta}/\Delta S_{\text{CH}}^{\alpha}$ ratio in agreement with the experimental ratio; all other models underestimated the α segment order parameter decrease with bound cations with respect to the β segment decrease.

In conclusion, a clear correlation between bound cations and order parameter decrease was observed for all simulation models. Consequently, the molecular electrometer can be used to compare the cation binding affinity between experiments and simulations. However, we found that quantitatively the response of α and β segment order parameters to bound cations in simulations did not generally agree with the experiments; e.g., the $\Delta S_{\text{CH}}^{\beta}/\Delta S_{\text{CH}}^{\alpha}$ ratio agreed with experiments only in the Lipid14 model (Fig. 5). Thus, the observed overestimation of the order parameter changes with salt concentrations could, in principle, arise from overbinding of cations or from an oversensitive lipid headgroup response to the bound cations (see also discussion in ESI[†]). A careful analysis with current lipid models is performed in the next section.

2.3 Cation binding in different simulation models

The order parameter changes (Fig. 2) and density distributions (Fig. 4) demonstrated significantly different Na^+ binding affinities in different simulation models. The best agreement with experiments (lowest $\Delta S_{\text{CH}}^{\alpha}$ and $\Delta S_{\text{CH}}^{\beta}$) was observed for the three models (Orange, Lipid14, CHARMM36; see Fig. 2) that predicted the lowest Na^+ densities near the bilayer (Fig. 4). All the other models clearly overestimated the choline order parameter responses to NaCl (Fig. 2) — and notably the strength of the overestimation was clearly linked to the strength of the Na^+ binding affinity (compare Figs. 2 and 4), which leads us to conclude that Na^+ binding affinity was overestimated in all these models.

As in the best three models the order parameter changes with NaCl were small (< 0.02), the achieved statistical accuracy did not allow us to conclude which of the three had the most realistic Na^+

binding affinity, especially at physiological NaCl concentrations ($\sim 150 \text{ mM}$) relevant for most applications. The overestimated binding in the other models raises questions concerning the quality of predictions from these models when NaCl is present. Especially interactions between charged molecules and the bilayer might be significantly affected by the strong Na^+ binding, which gives the otherwise neutral bilayer an effective positive charge.

Significant Ca^{2+} binding affinity for phosphatidylcholine bilayers at sub-molar concentrations is agreed on in the literature^{2,3,19,28}, however, several details remain under discussion. Simulations suggest that Ca^{2+} binds to lipid carbonyl oxygens with a coordination number of 4.2¹³, while interpretation of NMR and scattering experiments suggest that one Ca^{2+} interacts mainly with the choline groups^{92–94} of two phospholipid molecules²⁸. A simulation model correctly reproducing the order parameter changes would resolve the discussion by giving atomistic resolution interpretation for the experiments.

As a function of CaCl_2 concentration, all models but one (CHARMM36 with the recent heptahydrated Ca^{2+} by Yoo et al.⁷) overestimated the order parameter decrease (Fig. 2), which according to the molecular electrometer indicates too strong Ca^{2+} binding. (We note that while this is the most likely scenario for the models that overestimated changes in both order parameters, for CaCl_2 it is possible also that the headgroup response is oversensitive to bound cations, see ESI[†].) In CHARMM36 with the heptahydrated Ca^{2+} by Yoo et al.⁷, $\Delta S_{\text{CH}}^{\beta}$ was overestimated but $\Delta S_{\text{CH}}^{\alpha}$ underestimated (Fig. 2), in line with the $\Delta S_{\text{CH}}^{\beta}/\Delta S_{\text{CH}}^{\alpha}$ ratio in CHARMM36 being larger than in experiments (Fig. 5). As we do not know whether $\Delta S_{\text{CH}}^{\beta}$ or $\Delta S_{\text{CH}}^{\alpha}$ was more realistic, we cannot conclude whether Ca^{2+} binding was too strong or too weak in CHARMM36. This could be resolved by comparing against experimental data with a known amount of bound charge (e.g., amphiphilic cations³⁰²), however, such simulation data are not currently available.

The density distributions with CaCl_2 showed significant Ca^{2+} binding in all models (Fig. 6), however, some differences occurred in details. The Berger model predicted deeper penetration (density maximum at $\sim 1.8 \text{ nm}$) compared to other models ($\sim 2 \text{ nm}$); the latter value is probably more realistic as ^1H NMR and neutron scattering data indicate that Ca^{2+} interacts mainly with the choline group^{2,92–94}. In CHARMM36 (but not in Slipids) practically all Ca^{2+} ions present in the simulation bound the bilayer within $2 \mu\text{s}$ (Fig. 6 and ESI[†]), which hints that the Ca^{2+} binding affinity of CHARMM36 is among the strongest of these models.

The origin of inaccuracies in lipid-ion interactions and binding affinities is far from clear. Potential candidates are, e.g., discrepancies in the ion models^{88–90}, incomplete treatment of electronic polarizability⁸⁷, and inaccuracies in the lipid headgroup description³⁵.

Considering the ion models, Cordomi et al.²⁵ showed the Na^+ binding affinity to decrease when ion radius is increased; however, in their DPPC bilayer simulations (with the OPLS-AA force field²) even the largest Na^+ radii still resulted in significant binding. In our results, the Slipids force field gave essentially similar binding affinity with ion parameters from Refs. 73 and 69,70

Table 1 List of MD simulations. The salt concentrations calculated as $[\text{salt}] = N_c \times [\text{water}] / N_w$, where $[\text{water}] = 55.5 \text{ M}$; these correspond the concentrations reported in the experiments by Akutsu et al.¹⁹. The lipid force fields named as in our previous work³⁵.

force field for lipids / ions	lipid	salt	[salt] (mM)	^a N _l	^b N _w	^c N _c	^d T (K)	^e t _{sim} (ns)	^f t _{anal} (ns)	^g files
Berger-POPC-07 ³⁶ / –	POPC	no	0	128	7290	0	298	270	50	37
Berger-POPC-07 ³⁶ / ffgm ^x ³⁸	"	NaCl	340	"	7202	44	"	110	"	39
Berger-POPC-07 ³⁶ / ffgm ^x ³⁸	"	CaCl ₂	340	"	7157	"	"	108	58	40
Berger-DPPC-97 ⁴¹ / –	DPPC	no	0	72	2880	0	323	60	50	42
Berger-DPPC-97 ⁴¹ / ffgm ^x ³⁸	"	NaCl	150	"	"	8	"	120	60	43
Berger-DPPC-97 ⁴¹ / ffgm ^x ³⁸	"	"	1000	"	2778	51	"	"	"	44
BergerOPLS-DPPC-06 ⁴⁵ / –	DPPC	no	0	72	2880	0	323	120	60	46
BergerOPLS-DPPC-06 ⁴⁵ / OPLS ⁴⁷	"	NaCl	150	"	"	8	"	"	"	48
BergerOPLS-DPPC-06 ⁴⁵ / OPLS ⁴⁷	"	"	1000	"	2778	51	"	"	"	49
CHARMM36 ⁵⁰ / –	POPC	no	0	128	5210	0	303	200	150	?
CHARMM36 ⁵⁰ / –	"	"	0	72	2242	"	"	30	20	51
CHARMM36 ⁵⁰ / CHARMM36 ⁵²	"	NaCl	350	"	2085	13	"	80	60	53
CHARMM36 ⁵⁰ / CHARMM36 ⁵²	"	"	690	"	"	26	"	73	"	54
CHARMM36 ⁵⁰ / CHARMM36 ⁵²	"	"	950	"	2168	37	"	80	"	55
CHARMM36 ⁵⁰ / CHARMM36	"	CaCl ₂	350	128	6400	35	"	200	100	56
CHARMM36 ⁵⁰ / CHARMM36	"	"	450	200	9000	73	310	2000	"	?
CHARMM36 ⁵⁰ / CHARMM36	"	"	670	128	6400	67	303	200	120	57
CHARMM36 ⁵⁰ / CHARMM36	"	"	1000	"	"	100	"	"	100	58
CHARMM36 ⁵⁰ / –	DPPC	no	0	128	8000	0	323	170	150	–
CHARMM36 ⁵⁰ / Yoo [?]	"	CaCl ₂	430	"	7760	60	"	200	170	–
CHARMM36 ⁵⁰ / Yoo [?]	"	"	890	"	7520	120	"	"	"	–
MacRog ⁵⁹ / –	POPC	no	0	128	6400	0	310	400	200	?
MacRog ⁵⁹ / –	"	"	0	288	14400	"	"	90	40	60
MacRog ⁵⁹ / OPLS ⁴⁷	"	NaCl	100	"	14554	27	"	"	50	61
MacRog ⁵⁹ / OPLS ⁴⁷	"	"	210	"	14500	54	"	"	"	"
MacRog ⁵⁹ / OPLS ⁴⁷	"	"	310	"	14446	81	"	"	"	"
MacRog ⁵⁹ / OPLS ⁴⁷	"	"	420	"	14392	108	"	"	"	"
Orange / –	POPC	no	0	72	2880	0	298	60	50	62
Orange / OPLS ⁴⁷	"	NaCl	140	"	2866	7	"	120	60	63
Orange / OPLS ⁴⁷	"	"	510	"	2802	26	"	"	100	64
Orange / OPLS ⁴⁷	"	"	1000	"	2780	50	"	"	80	65
Orange / OPLS	"	CaCl ₂	510	"	2802	26	"	"	60	66
Slipids ⁷¹ / –	POPC	no	0	128	5120	0	310	200	150	72
Slipids ⁷¹ / AMBER ⁷³	"	NaCl	130	200	9000	21	"	105	100	74
Slipids ⁷¹ / AMBER ⁴⁷	"	CaCl ₂	450	"	"	73	"	2000	"	?
Slipids ⁶⁷ / –	DPPC	no	0	128	3840	0	323	150	100	68
Slipids ⁶⁷ / AMBER ^{69,70}	"	NaCl	150	600	18000	49	"	100	40	–
Slipids ⁶⁷ / AMBER ^{69,70}	"	"	850	128	3726	57	"	205	200	?
Slipids ⁶⁷ / AMBER ^{69,70}	"	"	1750	"	3612	114	"	105	100	"
Slipids ⁶⁷ / AMBER ^{69,70}	"	"	2570	"	3514	163	"	"	"	"
Lipid14 ⁷⁷ / –	POPC	no	0	128	5120	0	298	205	200	78
Lipid14 ⁷⁷ / AMBER ⁴⁷	"	NaCl	150	"	"	12	"	"	"	79
Lipid14 ⁷⁷ / AMBER ⁴⁷	"	"	1000	"	"	77	"	"	"	80
Lipid14 ⁷⁷ / AMBER ⁴⁷	"	CaCl ₂	350	"	6400	35	"	200	100	81
Lipid14 ⁷⁷ / AMBER ⁴⁷	"	"	1000	"	"	100	"	"	"	82
Ulmschneiders ⁸³ / –	POPC	no	0	128	5120	0	298	2×205	2×200	84
Ulmschneiders ⁸³ / OPLS ⁴⁷	"	NaCl	150	"	"	12	"	205	200	85
Ulmschneiders ⁸³ / OPLS ⁴⁷	"	"	1000	"	"	77	"	"	"	86

^a Number of lipid molecules

^b Number of water molecules

^c Number of cations

^d Simulation temperature

^e Total simulation time

^f Time used for analysis

^g Reference for simulation files




Fig. 6 Ca^{2+} (solid line) and Cl^- (dashed) distributions along the lipid bilayer normal from MD simulations. For clarity, only one CaCl_2 concentration per MD model is shown; see ESI[†] for a plot including all the available concentrations. The light green vertical lines indicate the locations of the Phosphorus maxima, used to define bound cations in Fig. 3.

(Fig. 4). Further, compensation of missing electronic polarizability by scaling the ion charge^{87?} reduced Na^+ binding in Berger, Berger-OPLS and Slipids, but not enough to reach agreement with experiments (ESI[†]). The charge-scaled Ca^{2+} model⁷⁵ slightly reduced binding in CHARMM36, but did not have significant influence in Slipids (ESI[†]). The heptahydrated Ca^{2+} ions by Yoo et al.⁷ significantly reduced Ca^{2+} binding in CHARMM36 (Fig. 6), however, the model must be further analysed to fully interpret the results.

The lipid models may also have a significant influence on ion binding behaviour. For example, the same ion model and non-bonded parameters are used in Orange and Berger-OPLS⁴⁵, but while Na^+ ion binding affinity appeared realistic in Orange, it was significantly overestimated in Berger-OPLS (Fig. 4). However, realistic Na^+ binding does not automatically imply realistic Ca^{2+} binding (see Orange, Lipid14, and CHARMM36 in Fig. 2) or realistic choline order parameter response to bound charge (see Orange and CHARMM36 in Fig. 5). It should also be noted that the low binding affinity of Na^+ in CHARMM36 is due to the additional repulsion (NBFIX⁵²) added between the sodium ions and lipid oxygens (ESI[†]), and that in the Ca^{2+} model by Yoo et al.⁷ the calcium is forced to be solvated solely by water. Altogether, our results indicate that probably both, lipid and ion force field parameters, need improvement to correctly predict the cation binding affinity, and the associated structural changes.

3 Conclusions

In accordance with the molecular electrometer^{19,28?–30}, cation binding to lipid bilayers was accompanied with a decrease in the C–H order parameters of the PC head group α and β carbons in all the simulation models tested (Fig. 3) — despite of the known inaccuracies in the actual atomistic resolution structures³⁵. Hence, the molecular electrometer allowed a direct comparison of Na^+ binding affinity between simulations and noninvasive NMR experiments. The comparison revealed that most models overestimated Na^+ binding; only Orange, Lipid14, and CHARMM36 predicted realistic binding affinities. None of the tested models had the accuracy required to interpret the Ca^{2+} :lipid stoichiometry or the induced structural changes with atomistic resolution.

Taken together, our results corroborate the pre-2000 view that at sub-molar concentrations, in contrast to Ca^{2+} and other multivalent ions^{1–4,10,11,19–21,28}, Na^+ and other monovalent ions (except Li^+) do not specifically bind to phospholipid bilayers. Concerning the interpretation of existing experimental data, our work supports Ceve's view² that the observed small shift in phase transition temperature is not indicative of Na^+ binding. Further, our findings are in line with the noninvasive NMR spectroscopy work of Filippov et al.¹¹ that proved the results of Refs. 7,9,12 to be explainable by direct interactions between Na^+ ions and fluorescent probes. Finally, as spectroscopic methods are in general more sensitive to atomistic details in fluid-like environment than AFM, our work indirectly suggests that the ion binding reported from AFM experiments on fluid-like lipid bilayer systems^{14–18} might be confounded with other physical features of the system. Concerning contradictions in MD simulation results, we reinterpret the strong Na^+ binding as an artefact of several simulation models, e.g., the Berger model used in Refs. 12,13.

The artificial specific Na^+ binding in MD simulations may lead to doubtful results, as it effectively results in a positively charged phosphatidylcholine lipid bilayer even at physiological NaCl concentrations. Such a charged bilayer will have distinctly different interactions with charged objects than what a (more realistic) model without specific Na^+ binding would predict. Furthermore, the overestimation of binding affinity may extend from ions to other positively charged objects, say, membrane protein seg-

ments. This would affect lipid–protein interactions and could explain, for example, certain contradicting results on electrostatic interactions between charged protein segments and lipid bilayers^{96,97}. In conclusion, more careful studies and model development on lipid bilayer–charged object interactions are urgently called for to make molecular dynamics simulations directly usable in a physiologically relevant electrolytic environment.

This work was done as a fully open collaboration, using `nmrlipids.blogspot.fi` as the communication platform. All the scientific contributions were communicated publicly through this blog or the GitHub repository¹⁰⁵. All the related content and data are available at Ref. 105.

Acknowledgements: AC and VSO wish to thank the Research Computing Service at UEA for access to the High Performance Computing Cluster; VSO acknowledges the Engineering and Physical Sciences Research Council in the UK for financial support (EP/L001322/1). MG acknowledges financial support from Finnish Center of International Mobility (Fellowship TM-9363). J. Melcr acknowledges computational resources provided by the CESNET LM2015042 and the CERIT Scientific Cloud LM2015085 projects under the program "Projects of Large Research, Development, and Innovations Infrastructure". MSM acknowledges financial support from the Volkswagen Foundation (86110). LM acknowledges funding from the Institut National de la Sante et de la Recherche Medicale (INSERM). OHSO acknowledges Tiago Ferreira for very useful discussions, the Emil Aaltonen foundation for financial support, Aalto Science-IT project and CSC-IT Center for Science for computational resources.

References

- 1 M. Eisenberg, T. Gresalfi, T. Riccio, and S. McLaughlin, *Biochemistry* **18**, 5213 (1979).
- 2 G. Cevc, *Biochim. Biophys. Acta - Rev. Biomemb.* **1031**, 311 (1990).
- 3 J.-F. Tocanne and J. Teissie, *Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes* **1031**, 111 (1990).
- 4 H. Binder and O. Zschörnig, *Chem. Phys. Lipids* **115**, 39 (2002).
- 5 J. J. Garcia-Celma, L. Hatahet, W. Kunz, and K. Fendler, *Langmuir* **23**, 10074 (2007).
- 6 E. Leontidis and A. Aroti, *The Journal of Physical Chemistry B* **113**, 1460 (2009).
- 7 R. Vacha, S. W. I. Siu, M. Petrov, R. A. Böckmann, J. Barucha-Kraszewska, P. Jurkiewicz, M. Hof, M. L. Berkowitz, and P. Jungwirth, *J. Phys. Chem. A* **113**, 7235 (2009).
- 8 B. Klasczyk, V. Knecht, R. Lipowsky, and R. Dimova, *Langmuir* **26**, 18951 (2010).
- 9 F. F. Harb and B. Tinland, *Langmuir* **29**, 5540 (2013).
- 10 G. Pabst, A. Hodzic, J. Strancar, S. Danner, M. Rappolt, and P. Laggner, *Biophys. J.* **93**, 2688 (2007).
- 11 A. Filippov, G. Orädd, and G. Lindblom, *Chemistry and Physics of Lipids* **159**, 81 (2009).
- 12 R. A. Böckmann, A. Hac, T. Heimburg, and H. Grubmüller, *Biophys. J.* **85**, 1647 (2003).
- 13 R. A. Böckmann and H. Grubmüller, *Ang. Chem. Int. Ed.* **43**, 1021 (2004).
- 14 S. Garcia-Manyes, G. Oncins, and F. Sanz, *Biophys. J.* **89**, 1812 (2005).
- 15 S. Garcia-Manyes, G. Oncins, and F. Sanz, *Electrochimica Acta* **51**, 5029 (2006), ISSN 0013-4686, bioelectrochemistry 2005 Bioelectrochemistry 2005, URL <http://www.sciencedirect.com/science/article/pii/S0013468606002775>.
- 16 T. Fukuma, M. J. Higgins, and S. P. Jarvis, *Phys. Rev. Lett.* **98**, 106101 (2007).
- 17 U. Ferber, G. Kaggwa, and S. Jarvis, *European Biophysics Journal* **40**, 329 (2011), ISSN 0175-7571, URL <http://dx.doi.org/10.1007/s00249-010-0650-7>.
- 18 L. Redondo-Morata, G. Oncins, and F. Sanz, *Biophysical Journal* **102**, 66 (2012).
- 19 H. Akutsu and J. Seelig, *Biochemistry* **20**, 7366 (1981).
- 20 R. J. Clarke and C. L. Aijpfert, *Biophysical Journal* **76**, 2614 (1999).
- 21 S. A. TATULIAN, *European Journal of Biochemistry* **170**, 413 (1987), ISSN 1432-1033, URL <http://dx.doi.org/10.1111/j.1432-1033.1987.tb13715.x>.
- 22 M. L. Berkowitz, D. L. Bostick, and S. Pandit, *Chem. Rev.* **106**, 1527 (2006).
- 23 V. Knecht and B. Klasczyk, *Biophys. J.* **104**, 818 (2013).
- 24 J. N. Sachs, H. Nanda, H. I. Petrache, and T. B. Woolf, *Biophys. J.* **86**, 3772 (2004).
- 25 A. Cordomila, O. Edholm, and J. J. Perez, *J. Chem. Theo. Comput.* **5**, 2125 (2009).
- 26 C. Valley, J. Perlmutter, A. Braun, and J. Sachs, *J. Membr. Biol.* **244**, 35 (2011).
- 27 M. L. Berkowitz and R. Vacha, *Acc. Chem. Res.* **45**, 74 (2012).
- 28 C. Altenbach and J. Seelig, *Biochemistry* **23**, 3913 (1984).
- 29 J. Seelig, P. M. MacDonald, and P. G. Scherer, *Biochemistry* **26**, 7535 (1987).
- 30 P. G. Scherer and J. Seelig, *Biochemistry* **28**, 7720 (1989).
- 31 O. H. S. Ollila and G. Pabst, *Atomistic resolution structure and dynamics of lipid bilayers in simulations and experiments* (2015), submitted to BBA, URL https://github.com/NMRLipids/NMRLipids_V-Review/blob/master/Manuscript/review_manu.pdf.
- 32 M. Hong, K. Schmidt-Rohr, and A. Pines, *Journal of the American Chemical Society* **117**, 3310 (1995).
- 33 M. Hong, K. Schmidt-Rohr, and D. Nanz, *Biophysical Journal* **69**, 1939 (1995).
- 34 J. D. Gross, D. E. Warschawski, and R. G. Griffin, *Journal of the American Chemical Society* **119**, 796 (1997).
- 35 A. Botan, A. Catte, F. Favela, P. Fuchs, M. Javanainen, W. Kulig, A. Lamberg, M. S. Miettinen, L. Monticelli, J. Määtä, et al., *Towards atomistic resolution structure of phosphatidylcholine glycerol backbone and choline headgroup at different ambient conditions*, <http://arxiv.org/abs/1309.2131v2> (2015), nMRLipids project, `nmrlipids.blogspot.fi`, 1309.2131.

- 36 S. Ollila, M. T. Hyvönen, and I. Vattulainen, *J. Phys. Chem. B* **111**, 3139 (2007).
- 37 O. H. S. Ollila, T. Ferreira, and D. Topgaard (2014), URL {<http://dx.doi.org/10.5281/zenodo.13279>}.
- 38 T. P. Straatsma and H. J. C. Berendsen, *The Journal of Chemical Physics* **89** (1988).
- 39 O. O. H. Samuli, *MD simulation trajectory and related files for POPC bilayer with 340mM NaCl (Berger model delivered by Tieleman, ffgmx ions, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.32144>.
- 40 O. O. H. Samuli, *MD simulation trajectory and related files for POPC bilayer with 340mM CaCl₂ (Berger model delivered by Tieleman, ffgmx ions, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.32173>.
- 41 S.-J. Marrink, O. Berger, P. Tieleman, and F. Jähnig, *Biophysical Journal* **74**, 931 (1998).
- 42 J. Määttä (2015), URL {<http://dx.doi.org/10.5281/zenodo.13934>}.
- 43 J. Määttä, *Dppc_berger_nacl* (2015), URL <http://dx.doi.org/10.5281/zenodo.16319>.
- 44 J. Määttä, *Dppc_berger_nacl_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.17210>.
- 45 D. P. Tieleman, J. L. MacCallum, W. L. Ash, C. Kandt, Z. Xu, and L. Monticelli, *J. Phys. Condens. Matter* **18**, S1221 (2006).
- 46 J. Määttä, *Dppc_berger_opls06* (2015), URL <http://dx.doi.org/10.5281/zenodo.17237>.
- 47 J. Åqvist, *The Journal of Physical Chemistry* **94**, 8021 (1990).
- 48 J. Määttä, *Dppc_berger_opls06_nacl* (2015), URL <http://dx.doi.org/10.5281/zenodo.16484>.
- 49 J. Määttä, *Dppc_berger_opls06_nacl_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.17208>.
- 50 J. B. Klauda, R. M. Venable, J. A. Freites, J. W. O'Connor, D. J. Tobias, C. Mondragon-Ramirez, I. Vorobyov, A. D. M. Jr, and R. W. Pastor, *J. Phys. Chem. B* **114**, 7830 (2010).
- 51 O. O. H. Samuli and M. Miettinen (2015), URL {<http://dx.doi.org/10.5281/zenodo.13944>}.
- 52 R. M. Venable, Y. Luo, K. Gawrisch, B. Roux, and R. W. Pastor, *The Journal of Physical Chemistry B* **117**, 10183 (2013).
- 53 S. Ollila, *MD simulation trajectory and related files for POPC bilayer with 350mM NaCl (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.32496>.
- 54 S. Ollila, *MD simulation trajectory and related files for POPC bilayer with 690mM NaCl (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.32497>.
- 55 S. Ollila, *MD simulation trajectory and related files for POPC bilayer with 950mM NaCl (CHARMM36, Gromacs 4.5)* (2015), URL <http://dx.doi.org/10.5281/zenodo.32498>.
- 56 G. Mykhailo and O. O. H. Samuli, *Popc_charmm36_cacl2_035mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.35159>.
- 57 G. Mykhailo and O. O. H. Samuli, *Popc_charmm36_cacl2_067mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.35160>.
- 58 G. Mykhailo and O. O. H. Samuli, *Popc_charmm36_cacl2_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.35156>.
- 59 A. Maciejewski, M. Pasenkiewicz-Gierula, O. Cramariuc, I. Vattulainen, and T. Rog, *The Journal of Physical Chemistry B* **118**, 4571 (2014).
- 60 M. Javanainen (2014), URL {<http://dx.doi.org/10.5281/zenodo.13498>}.
- 61 M. Javanainen, *POPC @ 310K, varying amounts of NaCl. Model by Maciejewski and Rog* (2015), URL <http://dx.doi.org/10.5281/zenodo.14976>.
- 62 O. H. S. Ollila, J. Määttä, and L. Monticelli, *MD simulation trajectory for POPC bilayer (Orange, Gromacs 4.5.)* (2015), URL <http://dx.doi.org/10.5281/zenodo.34488>.
- 63 O. H. S. Ollila, J. Määttä, and L. Monticelli, *MD simulation trajectory for POPC bilayer with 140mM NaCl (Orange, Gromacs 4.5.)* (2015), URL <http://dx.doi.org/10.5281/zenodo.34491>.
- 64 O. H. S. Ollila, J. Määttä, and L. Monticelli, *MD simulation trajectory for POPC bilayer with 510mM NaCl (Orange, Gromacs 4.5.)* (2015), URL <http://dx.doi.org/10.5281/zenodo.34490>.
- 65 S. Ollila, J. Määttä, and L. Monticelli, *MD simulation trajectory for POPC bilayer with 1000mM NaCl (Orange, Gromacs 4.5.)* (2015), URL <http://dx.doi.org/10.5281/zenodo.34497>.
- 66 O. H. S. Ollila, J. Määttä, and L. Monticelli, *MD simulation trajectory for POPC bilayer with 510mM CaCl₂ (Orange, Gromacs 4.5.)* (2015), URL <http://dx.doi.org/10.5281/zenodo.34498>.
- 67 J. P. M. Jämbeck and A. P. Lyubartsev, *The Journal of Physical Chemistry B* **116**, 3164 (2012).
- 68 J. Määttä (2014), URL {<http://dx.doi.org/10.5281/zenodo.13287>}.
- 69 D. Beglov and B. Roux, *The Journal of Chemical Physics* **100** (1994).
- 70 B. Roux, *Biophysical Journal* **71**, 3177 (1996), ISSN 0006-3495, URL <http://www.sciencedirect.com/science/article/pii/S0006349596795115>.
- 71 J. P. M. Jämbeck and A. P. Lyubartsev, *Journal of Chemical Theory and Computation* **8**, 2938 (2012).
- 72 M. Javanainen, *Popc @ 310K, slipids force field.* (2015), DOI: 10.5281/zenodo.13887.
- 73 D. E. Smith and L. X. Dang, *The Journal of Chemical Physics* **100** (1994).
- 74 M. Javanainen, *POPC @ 310K, 130 mM of NaCl. Slipids with ions by Smith & Dang* (2015), URL <http://dx.doi.org/10.5281/zenodo.35275>.
- 75 M. Kohagen, P. E. Mason, and P. Jungwirth, *The Journal of*

- Physical Chemistry B **118**, 7902 (2014).
- 76 M. Javanainen, *POPC @ 310K, 450 mM of CaCl₂. Slipids with ECC-scaled ions* (2016), URL <http://dx.doi.org/10.5281/zenodo.45007>.
 - 77 C. J. Dickson, B. D. Madej, Å. A. Skjevik, R. M. Betz, K. Teigen, I. R. Gould, and R. C. Walker, *Journal of Chemical Theory and Computation* **10**, 865 (2014).
 - 78 M. Girych and O. H. S. Ollila, *Popc_amber_lipid14_verlet* (2015), URL <http://dx.doi.org/10.5281/zenodo.30898>.
 - 79 M. Girych and O. H. S. Ollila, *Popc_amber_lipid14_nacl_015mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.30891>.
 - 80 M. Girych and O. H. S. Ollila, *Popc_amber_lipid14_nacl_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.30865>.
 - 81 G. Mykhailo and O. O. H. Samuli, *Popc_amber_lipid14_cacl2_035mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.34415>.
 - 82 G. Mykhailo and O. O. H. Samuli, *Popc_amber_lipid14_cacl2_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.35074>.
 - 83 J. P. Ulmschneider and M. B. Ulmschneider, *Journal of Chemical Theory and Computation* **5**, 1803 (2009).
 - 84 M. Girych and O. H. S. Ollila, *Popc_ulmschneider_opls_verlet_group* (2015), URL <http://dx.doi.org/10.5281/zenodo.30904>.
 - 85 M. Girych and O. H. S. Ollila, *Popc_ulmschneider_opls_nacl_015mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.30892>.
 - 86 M. Girych and O. H. S. Ollila, *Popc_ulmschneider_opls_nacl_1mol* (2015), URL <http://dx.doi.org/10.5281/zenodo.30894>.
 - 87 I. Leontyev and A. Stuchebrukhov, *Phys. Chem. Chem. Phys.* **13**, 2613 (2011).
 - 88 B. Hess, C. Holm, and N. van der Vegt, *The Journal of Chemical Physics* **124** (2006).
 - 89 A. A. Chen, , and R. V. Pappu, *The Journal of Physical Chemistry B* **111**, 11884 (2007).
 - 90 M. M. Reif, M. Winger, and C. Oostenbrink, *Journal of Chemical Theory and Computation* **9**, 1247 (2013), PMID: 23418406, <http://dx.doi.org/10.1021/ct300874c>, URL <http://dx.doi.org/10.1021/ct300874c>.
 - 91 M. Kohagen, P. E. Mason, and P. Jungwirth, *The Journal of Physical Chemistry B* **0**, null (0), PMID: 26172524, <http://dx.doi.org/10.1021/acs.jpccb.5b05221>, URL <http://dx.doi.org/10.1021/acs.jpccb.5b05221>.
 - 92 H. Hauser, M. C. Phillips, B. Levine, and R. Williams, *Nature* **261**, 390 (1976).
 - 93 H. Hauser, W. Guyer, B. Levine, P. Skrabal, and R. Williams, *Biochimica et Biophysica Acta (BBA) - Biomembranes* **508**, 450 (1978), ISSN 0005-2736, URL <http://www.sciencedirect.com/science/article/pii/0005273678900913>.
 - 94 L. Herbette, C. Napolitano, and R. McDaniel, *Biophysical Journal* **46**, 677 (1984).
 - 95 H. Hauser, W. Guyer, and F. Paltauf, *Chemistry and Physics of Lipids* **29**, 103 (1981).
 - 96 A. Arkhipov, Y. Shan, R. Das, N. Endres, M. Eastwood, D. Wemmer, J. Kuriyan, and D. Shaw, *Cell* **152**, 557 (2013).
 - 97 K. Kaszuba, M. Grzybek, A. Orłowski, R. Danne, T. Råsg, K. Simons, Å. Coskun, and I. Vattulainen, *Proceedings of the National Academy of Sciences* **112**, 4334 (2015).
 - 98 J. Määttä (2015), URL <http://dx.doi.org/10.5281/zenodo.16320>.
 - 99 J. Määttä, *Dppc_berger_nacl_1mol_scaled* (2015), URL <http://dx.doi.org/10.5281/zenodo.17228>.
 - 100 J. Määttä (2015), URL <http://dx.doi.org/10.5281/zenodo.16485>.
 - 101 J. Määttä, *Dppc_berger_opls06_nacl_1mol_scaled* (2015), URL <http://dx.doi.org/10.5281/zenodo.17209>.
 - 102 M. Javanainen, *POPC @ 310K, varying amounts of NaCl. Slipids with ECC-scaled ions* (2015), URL <http://dx.doi.org/10.5281/zenodo.35193>.
 - 103 M. Javanainen, *POPC @ 310K, 450 mM of CaCl₂. Charmm36 with ECC-scaled ions* (2016), URL <http://dx.doi.org/10.5281/zenodo.45008>.
 - 104 D. van der Spoel, E. Lindahl, B. Hess, A. R. van Buuren, E. Apol, P. J. Meulenhoff, D. P. Tieleman, A. L. T. M. Sijbers, K. A. Feenstra, R. van Drunen, et al., *GROMACS user manual version 4.0* (2005), URL www.gromacs.org.
 - 105 O. H. S. Ollila and et al. (2015), URL <https://github.com/NMRLipids/lipid-ionINTERACTION>.
 - 106 T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, *Phys. Chem. Chem. Phys.* **15**, 1976 (2013).
 - 107 O. Berger, O. Edholm, and F. Jähnig, *Biophys. J.* **72**, 2002 (1997).
 - 108 M. Bachar, P. Brunelle, D. P. Tieleman, and A. Rauk, *J. Phys. Chem. B* **108**, 7170 (2004).
 - 109 B. Hess, H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije, *J. Comput. Chem.* **18**, 1463 (1997).
 - 110 B. Hess, *Journal of Chemical Theory and Computation* **4**, 116 (2008).
 - 111 T. Darden, D. York, and L. Pedersen, *The Journal of Chemical Physics* **98** (1993).
 - 112 U. L. Essman, M. L. Perera, M. L. Berkowitz, T. Larden, H. Lee, and L. G. Pedersen, *J. Chem. Phys.* **103**, 8577 (1995).
 - 113 G. Bussi, D. Donadio, and M. Parrinello, *The Journal of Chemical Physics* **126** (2007).
 - 114 M. Parrinello and A. Rahman, *J. Appl. Phys.* **52**, 7182 (1981).
 - 115 S. Pronk, S. Páll, R. Schulz, P. Larsson, P. Bjelkmar, R. Apostolov, M. R. Shirts, J. C. Smith, P. M. Kasson, D. van der Spoel, et al., *Bioinformatics* **29**, 845 (2013).

- 116 H. J. C. Berendsen, J. P. M. Postma, W. F. van Gunsteren, A. DiNola, and J. R. Haak, *J. Chem. Phys.* **81**, 3684 (1984).
- 117 J. Lee, X. Cheng, J. M. Swails, M. S. Yeom, P. K. Eastman, J. A. Lemkul, S. Wei, J. Buckner, J. C. Jeong, Y. Qi, et al., *Journal of Chemical Theory and Computation* **0**, null (0).
- 118 M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl, *SoftwareX* **1**, 19 (2015), ISSN 2352-7110, URL <http://www.sciencedirect.com/science/article/pii/S2352711015000059>.
- 119 S. Nose, *Mol. Phys.* **52**, 255 (1984).
- 120 W. G. Hoover, *Phys. Rev. A* **31**, 1695 (1985).
- 121 S. Páll and B. Hess, *Computer Physics Communications* **184**, 2641 (2013), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/article/pii/S0010465513001975>.
- 122 M. Abraham, D. van der Spoel, E. Lindahl, B. Hess, and the GROMACS development team, *GROMACS user manual version 5.0.7* (2015), URL www.gromacs.org.
- 123 W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, and M. L. Klein, *The Journal of Chemical Physics* **79** (1983).
- 124 O. H. S. Ollila and M. Retegan, *Md simulation trajectory and related files for popc bilayer (lipid14, gromacs 4.5)* (2014), URL <http://dx.doi.org/10.5281/zenodo.12767>.
- 125 J. Domański, P. Stansfeld, M. Sansom, and O. Beckstein, *The Journal of Membrane Biology* **236**, 255 (2010), ISSN 0022-2631.