

Deep Generative Models

Lecture 3

Roman Isachenko

Moscow Institute of Physics and Technology

Autumn, 2021

Recap of previous lecture

MLE problem for autoregressive model

$$\theta^* = \arg \max_{\theta} p(\mathbf{X}|\theta) = \arg \max_{\theta} \sum_{i=1}^n \sum_{j=1}^m \log p(x_{ij}|\mathbf{x}_{i,1:j-1}\theta).$$

Sampling

$$\hat{x}_1 \sim p(x_1|\theta), \quad \hat{x}_2 \sim p(x_2|\hat{x}_1, \theta), \dots, \quad \hat{x}_m \sim p(x_m|\hat{\mathbf{x}}_{1:m-1}, \theta)$$

New generated object is $\hat{\mathbf{x}} = (\hat{x}_1, \hat{x}_2, \dots, \hat{x}_m)$.

Masking helps to make neural network autoregressive.

- ▶ **MADE** - masked autoencoder (MLP).
- ▶ **WaveNet** - masked 1D convolutions.
- ▶ **PixelCNN** - masked 2D convolutions.

PixelCNN++ uses discretized mixture of logistic distribution to make the output distribution more natural.

Recap of previous lecture

Posterior distribution

$$p(\boldsymbol{\theta}|\mathbf{X}) = \frac{p(\mathbf{X}|\boldsymbol{\theta})p(\boldsymbol{\theta})}{p(\mathbf{X})} = \frac{p(\mathbf{X}|\boldsymbol{\theta})p(\boldsymbol{\theta})}{\int p(\mathbf{X}|\boldsymbol{\theta})p(\boldsymbol{\theta})d\boldsymbol{\theta}}$$

Bayesian inference

$$p(\mathbf{x}|\mathbf{X}) = \int p(\mathbf{x}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\mathbf{X})d\boldsymbol{\theta}$$

Maximum a posteriori (MAP) estimation

$$\boldsymbol{\theta}^* = \arg \max_{\boldsymbol{\theta}} p(\boldsymbol{\theta}|\mathbf{X}) = \arg \max_{\boldsymbol{\theta}} (\log p(\mathbf{X}|\boldsymbol{\theta}) + \log p(\boldsymbol{\theta}))$$

MAP inference

$$p(\mathbf{x}|\mathbf{X}) = \int p(\mathbf{x}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\mathbf{X})d\boldsymbol{\theta} \approx p(\mathbf{x}|\boldsymbol{\theta}^*).$$

Latent variable models (LVM)

MLE problem

$$\theta^* = \arg \max_{\theta} p(\mathbf{X}|\theta) = \arg \max_{\theta} \prod_{i=1}^n p(\mathbf{x}_i|\theta) = \arg \max_{\theta} \sum_{i=1}^n \log p(\mathbf{x}_i|\theta).$$

The distribution $p(\mathbf{x}|\theta)$ could be very complex and intractable (as well as real distribution $\pi(\mathbf{x})$).

Extended probabilistic model

Introduce latent variable \mathbf{z} for each sample \mathbf{x}

$$p(\mathbf{x}, \mathbf{z}|\theta) = p(\mathbf{x}|\mathbf{z}, \theta)p(\mathbf{z}); \quad \log p(\mathbf{x}, \mathbf{z}|\theta) = \log p(\mathbf{x}|\mathbf{z}, \theta) + \log p(\mathbf{z}).$$

$$p(\mathbf{x}|\theta) = \int p(\mathbf{x}, \mathbf{z}|\theta) d\mathbf{z} = \int p(\mathbf{x}|\mathbf{z}, \theta)p(\mathbf{z}) d\mathbf{z}.$$

Motivation


The distributions $p(\mathbf{x}|\mathbf{z}, \theta)$ and $p(\mathbf{z})$ could be quite simple.

Latent variable models (LVM)


$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \log \int p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})p(\mathbf{z})d\mathbf{z} \rightarrow \max_{\boldsymbol{\theta}}$$

Examples

Mixture of gaussians



PCA model




- ▶ $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_z, \boldsymbol{\Sigma}_z)$
- ▶ $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \sigma^2\mathbf{I})$
- ▶ $p(\mathbf{z}) = \text{Categorical}(\boldsymbol{\pi})$
- ▶ $p(\mathbf{z}) = \mathcal{N}(\mathbf{z}|\mathbf{0}, \mathbf{I})$

Latent variable models (LVM)

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \log \int p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) p(\mathbf{z}) d\mathbf{z} \rightarrow \max_{\boldsymbol{\theta}}$$

PCA projects original data \mathbf{X} onto a low dimensional latent space while maximizing the variance of the projected data.



- ▶ $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \sigma^2\mathbf{I})$
- ▶ $p(\mathbf{z}) = \mathcal{N}(\mathbf{z}|0, \mathbf{I})$
- ▶ $p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{W}\mathbf{W}^T + \sigma^2\mathbf{I})$
- ▶ $p(\mathbf{z}|\mathbf{x}) = \mathcal{N}(\mathbf{M}^{-1}\mathbf{W}^T(\mathbf{x} - \boldsymbol{\mu}), \sigma^2\mathbf{M})$, where $\mathbf{M} = \mathbf{W}\mathbf{W}^T + \sigma^2\mathbf{I}$

Maximum likelihood estimation for LVM

MLE for extended problem


$$\begin{aligned}\theta^* &= \arg \max_{\theta} p(\mathbf{X}, \mathbf{Z} | \theta) = \arg \max_{\theta} \prod_{i=1}^n p(\mathbf{x}_i, \mathbf{z}_i | \theta) = \\ &= \arg \max_{\theta} \sum_{i=1}^n \log p(\mathbf{x}_i, \mathbf{z}_i | \theta).\end{aligned}$$

However, \mathbf{Z} is unknown.

MLE for original problem

$$\begin{aligned}\theta^* &= \arg \max_{\theta} \log p(\mathbf{X} | \theta) = \arg \max_{\theta} \sum_{i=1}^n \log p(\mathbf{x}_i | \theta) = \\ &= \arg \max_{\theta} \sum_{i=1}^n \log \int p(\mathbf{x}_i, \mathbf{z}_i | \theta) d\mathbf{z}_i = \\ &= \arg \max_{\theta} \log \sum_{i=1}^n \int p(\mathbf{x}_i | \mathbf{z}_i, \theta) p(\mathbf{z}_i) d\mathbf{z}_i.\end{aligned}$$

Naive approach



Monte-Carlo estimation

$$p(\mathbf{x}|\theta) = \int p(\mathbf{x}|\mathbf{z}, \theta) p(\mathbf{z}) d\mathbf{z} = \mathbb{E}_{p(\mathbf{z})} p(\mathbf{x}|\mathbf{z}, \theta) \approx \frac{1}{K} \sum_{k=1}^K p(\mathbf{x}|\mathbf{z}_k, \theta)$$

Challenge: to cover the space properly, the number of samples grows exponentially with respect to dimensionality of \mathbf{z} .

Variational lower bound

Derivation 1

$$\begin{aligned}\log p(\mathbf{x}|\boldsymbol{\theta}) &= \log \int p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) d\mathbf{z} = \\ &= \log \int \frac{q(\mathbf{z})}{q(\mathbf{z})} p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) d\mathbf{z} = \log \mathbb{E}_q \left[\frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} \right] \geq \\ &\geq \mathbb{E}_q \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} = \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} d\mathbf{z} = \mathcal{L}(q, \boldsymbol{\theta})\end{aligned}$$

Derivation 2

$$\begin{aligned}\log p(\mathbf{x}|\boldsymbol{\theta}) &= \int q(\mathbf{z}) \log p(\mathbf{x}|\boldsymbol{\theta}) d\mathbf{z} = \\ &= \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})} d\mathbf{z} = \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) q(\mathbf{z})}{p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}) q(\mathbf{z})} d\mathbf{z} = \\ &= \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} d\mathbf{z} + \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})} d\mathbf{z} = \\ &= \mathcal{L}(q, \boldsymbol{\theta}) + KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(q, \boldsymbol{\theta}).\end{aligned}$$

Variational lower bound

Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}(q, \theta) &= \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z} | \theta)}{q(\mathbf{z})} d\mathbf{z} = \\ &= \int q(\mathbf{z}) \log p(\mathbf{x} | \mathbf{z}, \theta) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p(\mathbf{x} | \mathbf{z}, \theta) - KL(q(\mathbf{z}) || p(\mathbf{z}))\end{aligned}$$

Log-likelihood decomposition

$$\log p(\mathbf{x} | \theta) = \mathbb{E}_q \log p(\mathbf{x} | \mathbf{z}, \theta) - KL(q(\mathbf{z}) || p(\mathbf{z})) + KL(q(\mathbf{z}) || p(\mathbf{z} | \mathbf{x}, \theta)).$$

Instead of maximizing incomplete likelihood, maximize ELBO (equivalently minimize KL)

$$\max_{\theta} p(\mathbf{x} | \theta) \quad \rightarrow \quad \max_{q, \theta} \mathcal{L}(q, \theta) \equiv \min_{q, \theta} KL(q(\mathbf{z}) || p(\mathbf{z} | \mathbf{x}, \theta)).$$

EM-algorithm

$$\mathcal{L}(q, \theta) = \int q(\mathbf{z}) \log p(\mathbf{x}|\mathbf{z}, \theta) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z}.$$

Block-coordinate optimization

- ▶ Initialize θ^* ;
- ▶ E-step


$$\begin{aligned} q^*(\mathbf{z}) &= \arg \max_q \mathcal{L}(q, \theta^*) = \\ &= \arg \min_q KL(q(\mathbf{z}) || p(\mathbf{z}|\mathbf{x}, \theta^*)) = p(\mathbf{z}|\mathbf{x}, \theta^*); \end{aligned}$$

- ▶ M-step

$$\theta^* = \arg \max_{\theta} \mathcal{L}(q^*, \theta);$$

- ▶ Repeat E-step and M-step until convergence.

EM illustration



Amortized variational inference

E-step

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}(q, \boldsymbol{\theta}^*) = \arg \min_q KL(q||p) = p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}^*).$$

- ▶ $p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}^*)$ could be **intractable**;
- ▶ $q(\mathbf{z})$ is different for each object \mathbf{x} .

Idea

Restrict a family of all possible distributions $q(\mathbf{z})$ to a parametric class $q(\mathbf{z}|\mathbf{x}, \phi)$ conditioned on samples \mathbf{x} with parameters ϕ .

Variational Bayes

- ▶ E-step

$$\phi_k = \phi_{k-1} + \eta \nabla_{\phi} \mathcal{L}(\phi, \boldsymbol{\theta}_{k-1})|_{\phi=\phi_{k-1}}$$

- ▶ M-step

$$\boldsymbol{\theta}_k = \boldsymbol{\theta}_{k-1} + \eta \nabla_{\boldsymbol{\theta}} \mathcal{L}(\phi_k, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\boldsymbol{\theta}_{k-1}}$$

Variational EM-algorithm

ELBO

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta}).$$

► E-step

$$\boldsymbol{\phi}_k = \boldsymbol{\phi}_{k-1} + \eta \nabla_{\boldsymbol{\phi}} \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta}_{k-1})|_{\boldsymbol{\phi}=\boldsymbol{\phi}_{k-1}},$$

where $\boldsymbol{\phi}$ – parameters of variational distribution $q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})$.

► M-step

$$\boldsymbol{\theta}_k = \boldsymbol{\theta}_{k-1} + \eta \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\phi}_k, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\boldsymbol{\theta}_{k-1}},$$

where $\boldsymbol{\theta}$ – parameters of the generative distribution $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})$.

Now all we have to do is to obtain two gradients $\nabla_{\boldsymbol{\phi}} \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta})$, $\nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta})$.

Challenge: Number of samples n could be huge (we need to derive unbiased stochastic gradients).

ELBO gradients

$$\mathcal{L}(\phi, \theta) = \mathbb{E}_q \left[\log p(\mathbf{x}|\mathbf{z}, \theta) + \log \frac{p(\mathbf{z})}{q(\mathbf{z}|\mathbf{x}, \phi)} \right] \rightarrow \max_{\phi, \theta}.$$

M-step: $\nabla_{\theta} \mathcal{L}(\phi, \theta)$

$$\begin{aligned} \nabla_{\theta} \mathcal{L}(\phi, \theta) &= \int q(\mathbf{z}|\mathbf{x}, \phi) \nabla_{\theta} \log p(\mathbf{x}|\mathbf{z}, \theta) d\mathbf{z} \approx \\ &\approx \nabla_{\theta} \log p(\mathbf{x}|\mathbf{z}^*, \theta), \quad \mathbf{z}^* \sim q(\mathbf{z}|\mathbf{x}, \phi). \end{aligned}$$

E-step: $\nabla_{\phi} \mathcal{L}(\phi, \theta)$

Difference from M-step: density function $q(\mathbf{z}|\mathbf{x}, \phi)$ depends on the parameters ϕ , it is impossible to use the Monte-Carlo estimation:

$$\begin{aligned} \nabla_{\phi} \mathcal{L}(\phi, \theta) &= \nabla_{\phi} \int q(\mathbf{z}|\mathbf{x}, \phi) \left[\log p(\mathbf{x}|\mathbf{z}, \theta) + \log \frac{p(\mathbf{z})}{q(\mathbf{z}|\mathbf{x}, \phi)} \right] d\mathbf{z} \\ &\neq \int q(\mathbf{z}|\mathbf{x}, \phi) \nabla_{\phi} \left[\log p(\mathbf{x}|\mathbf{z}, \theta) + \log \frac{p(\mathbf{z})}{q(\mathbf{z}|\mathbf{x}, \phi)} \right] d\mathbf{z} \end{aligned}$$

Reparametrization trick

Law of the unconscious statistician (LOTUS)

Let X be a random variable and let $Y = g(X)$. Then

$$\mathbb{E}_{p_Y} Y = \mathbb{E}_{p_X} g(X) = \int g(\mathbf{x}) p(\mathbf{x}) d\mathbf{x}.$$

Examples

- ▶ $r(x) = \mathcal{N}(x|0, 1)$, $y = \sigma \cdot x + \mu$, $p_Y(y|\theta) = \mathcal{N}(y|\mu, \sigma^2)$, $\theta = [\mu, \sigma]$.
- ▶ $\epsilon^* \sim r(\epsilon)$, $\mathbf{z} = g(\mathbf{x}, \epsilon, \phi)$, $\mathbf{z} \sim q(\mathbf{z}|\mathbf{x}, \phi)$

$$\begin{aligned} \nabla_{\phi} \int q(\mathbf{z}|\mathbf{x}, \phi) f(\mathbf{z}) d\mathbf{z} &= \nabla_{\phi} \int r(\epsilon) f(\mathbf{z}) d\epsilon \\ &= \int r(\epsilon) \nabla_{\phi} f(g(\mathbf{x}, \epsilon, \phi)) d\epsilon \approx \nabla_{\phi} f(g(\mathbf{x}, \epsilon^*, \phi)) \end{aligned}$$

ELBO gradient (E-step, $\nabla_{\phi} \mathcal{L}(\phi, \theta)$)

$$\begin{aligned}\nabla_{\phi} \mathcal{L}(\phi, \theta) &= \nabla_{\phi} \int q(\mathbf{z}|\mathbf{x}, \phi) \log p(\mathbf{x}|\mathbf{z}, \theta) d\mathbf{z} - \nabla_{\phi} \text{KL}(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z})) \\ &= \int r(\epsilon) \nabla_{\phi} \log p(\mathbf{x}|g(\mathbf{x}, \epsilon, \phi), \theta) d\epsilon - \nabla_{\phi} \text{KL}(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z})) \\ &\approx \nabla_{\phi} \log p(\mathbf{x}|g(\mathbf{x}, \epsilon^*, \phi), \theta) - \nabla_{\phi} \text{KL}(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z}))\end{aligned}$$

Variational assumption

$$r(\epsilon) = \mathcal{N}(0, \mathbf{I}); \quad q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mu_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x})).$$

$$\mathbf{z} = g(\mathbf{x}, \epsilon, \phi) = \sigma_{\phi}(\mathbf{x}) \cdot \epsilon + \mu_{\phi}(\mathbf{x}).$$

Here $\mu_{\phi}(\cdot), \sigma_{\phi}(\cdot)$ are parameterized functions (outputs of neural network).

- ▶ $p(\mathbf{z})$ – prior distribution on latent variables \mathbf{z} . We could specify any distribution that we want. Let say $p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I})$.
- ▶ $p(\mathbf{x}|\mathbf{z}, \theta)$ – generative distribution. Since it is a parameterized function let it be neural network with parameters θ .

Variational autoencoder (VAE)

Final algorithm

- ▶ pick $i \sim U[1, n]$;
- ▶ compute a stochastic gradient w.r.t. ϕ

$$\begin{aligned}\nabla_{\phi} \mathcal{L}(\phi, \theta) \approx & \nabla_{\phi} \log p(\mathbf{x} | g(\mathbf{x}, \epsilon^*, \phi), \theta) \\ & - \nabla_{\phi} \text{KL}(q(\mathbf{z} | \mathbf{x}, \phi) || p(\mathbf{z})), \quad \epsilon^* \sim r(\epsilon);\end{aligned}$$

- ▶ compute a stochastic gradient w.r.t. θ


$$\nabla_{\theta} \mathcal{L}(\phi, \theta) \approx \nabla_{\theta} \log p(\mathbf{x} | \mathbf{z}^*, \theta), \quad \mathbf{z}^* \sim q(\mathbf{z} | \mathbf{x}, \phi);$$

- ▶ update θ, ϕ according to the selected optimization method (SGD, Adam, RMSProp):


$$\begin{aligned}\phi &:= \phi + \eta \nabla_{\phi} \mathcal{L}(\phi, \theta), \\ \theta &:= \theta + \eta \nabla_{\theta} \mathcal{L}(\phi, \theta).\end{aligned}$$

Variational autoencoder (VAE)

- ▶ VAE learns stochastic mapping between \mathbf{x} -space, from complicated distribution $\pi(\mathbf{x})$, and a latent \mathbf{z} -space, with simple distribution.
- ▶ The generative model learns a joint distribution $p(\mathbf{x}, \mathbf{z}|\theta) = p(\mathbf{z})p(\mathbf{x}|\mathbf{z}, \theta)$, with a prior distribution $p(\mathbf{z})$, and a stochastic decoder $p(\mathbf{x}|\mathbf{z}, \theta)$.
- ▶ The stochastic encoder $q(\mathbf{z}|\mathbf{x}, \phi)$ (inference model), approximates the true but intractable posterior $p(\mathbf{z}|\mathbf{x}, \theta)$ of the generative model.



Variational Autoencoder



Variational autoencoder (VAE)

- ▶ Encoder $q(\mathbf{z}|\mathbf{x}, \phi) = \text{NN}_e(\mathbf{x}, \phi)$ outputs $\mu_\phi(\mathbf{x})$ and $\sigma_\phi(\mathbf{x})$.
- ▶ Decoder $p(\mathbf{x}|\mathbf{z}, \theta) = \text{NN}_d(\mathbf{z}, \theta)$ outputs parameters of the sample distribution.




image credit:

<https://lilianweng.github.io/lil-log/2018/08/12/from-autoencoder-to-beta-vae.html>

Bayesian framework

Posterior distribution

$$p(\theta|\mathbf{X}) = \frac{p(\mathbf{X}|\theta)p(\theta)}{p(\mathbf{X})} = \frac{p(\mathbf{X}|\theta)p(\theta)}{\int p(\mathbf{X}|\theta)p(\theta)d\theta}$$

Bayesian inference

$$p(\mathbf{x}|\mathbf{X}) = \int p(\mathbf{x}|\theta)p(\theta|\mathbf{X})d\theta$$

Maximum a posteriori (MAP) estimation

$$\theta^* = \arg \max_{\theta} \log p(\theta|\mathbf{X}) = \arg \max_{\theta} (\log p(\mathbf{X}|\theta) + \log p(\theta))$$

MAP inference

$$p(\mathbf{x}|\mathbf{X}) = \int p(\mathbf{x}|\theta)p(\theta|\mathbf{X})d\theta = \int p(\mathbf{x}|\theta)\delta(\theta - \theta^*)d\theta \approx p(\mathbf{x}|\theta^*).$$

VAE as Bayesian model

Posterior distribution

$$p(\boldsymbol{\theta}|\mathbf{X}) = \frac{p(\mathbf{X}|\boldsymbol{\theta})p(\boldsymbol{\theta})}{p(\mathbf{X})}$$

ELBO

$$\begin{aligned}\log p(\boldsymbol{\theta}|\mathbf{X}) &= \log p(\mathbf{X}|\boldsymbol{\theta}) + \log p(\boldsymbol{\theta}) - \log p(\mathbf{X}) \\ &= \mathcal{L}(q, \boldsymbol{\theta}) + KL(q||p) + \log p(\boldsymbol{\theta}) - \log p(\mathbf{X}) \\ &\geq [\mathcal{L}(q, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta})] - \log p(\mathbf{X}).\end{aligned}$$

EM-algorithm

► E-step

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}(q, \boldsymbol{\theta}^*) = \arg \min_q KL(q||p) = p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}^*);$$

► M-step

$$\boldsymbol{\theta}^* = \arg \max_{\boldsymbol{\theta}} [\mathcal{L}(q, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta})].$$

Summary

- ▶ LVM introduces latent representation of observed samples to make model more interpretable.
- ▶ LVM maximizes variational evidence lower bound (ELBO) to find MLE of model parameters.
- ▶ The general variational EM algorithm maximizes ELBO objective.
- ▶ Amortized inference allows to efficiently compute stochastic gradients for ELBO using Monte-Carlo estimation.
- ▶ The reparametrization trick gets unbiased gradients w.r.t to a variational posterior distribution.
- ▶ The VAE model is an LVM with two neural network: for stochastic encoder $q(\mathbf{z}|\mathbf{x}, \phi)$ and for stochastic decoder $p(\mathbf{x}|\mathbf{z}, \theta)$.
- ▶ VAE is not a "true" bayesian model since parameters θ do not have a prior distribution.