

Deep Generative Models

Lecture 7

Roman Isachenko

Moscow Institute of Physics and Technology

Autumn, 2021

Variational posterior

ELBO

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$

- ▶ In E-step of EM-algorithm we wish $KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) = 0$.
(In this case the lower bound is tight $\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta})$).
- ▶ Normal variational distribution $q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_\phi(\mathbf{x}), \boldsymbol{\sigma}_\phi^2(\mathbf{x}))$ is poor (e.g. has only one mode).
- ▶ Flows models convert a simple base distribution to a complex one using invertible transformation with simple Jacobian. How to use flows in VAE?

Flows in VAE posterior

Apply a sequence of transformations to the random variable

$$\mathbf{z}_0 \sim q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_\phi(\mathbf{x}), \boldsymbol{\sigma}_\phi^2(\mathbf{x})).$$

Here, $q(\mathbf{z}|\mathbf{x}, \phi)$ (which is a VAE encoder) plays a role of a base distribution.

$$\mathbf{z}_0 \xrightarrow{g_1} \mathbf{z}_1 \xrightarrow{g_2} \dots \xrightarrow{g_K} \mathbf{z}_K, \quad \mathbf{z}_K = g(\mathbf{z}_0), \quad g = g_K \circ \dots \circ g_1.$$

Each g_k is a flow transformation (e.g. planar, coupling layer) parameterized by ϕ_k .

$$\begin{aligned} \log q_K(\mathbf{z}_K|\mathbf{x}, \phi, \{\phi_k\}_{k=1}^K) &= \log q(\mathbf{z}_0|\mathbf{x}, \phi) \\ &\quad - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right|. \end{aligned}$$

Flows in VAE posterior

ELBO

$$p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta}) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})} \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})} \rightarrow \max_{\boldsymbol{\phi}, \boldsymbol{\theta}}.$$

Flow model in latent space

$$\log q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*) = \log q(\mathbf{z}_0|\mathbf{x}, \boldsymbol{\phi}) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \boldsymbol{\phi}_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

Let use $q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*)$, $\boldsymbol{\phi}_* = \{\boldsymbol{\phi}, \boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_K\}$ as a variational distribution. Here $\boldsymbol{\phi}$ – encoder parameters, $\{\boldsymbol{\phi}_k\}_{k=1}^K$ – flow parameters.

- ▶ Encoder outputs base distribution $q(\mathbf{z}_0|\mathbf{x}, \boldsymbol{\phi})$.
- ▶ Flow model $\mathbf{z}_K = g(\mathbf{z}_0, \{\boldsymbol{\phi}_k\}_{k=1}^K)$ transforms the base distribution $q(\mathbf{z}_0|\mathbf{x}, \boldsymbol{\phi})$ to the distribution $q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*)$.
- ▶ Distribution $q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*)$ is used as a variational distribution for ELBO maximization.

Flows in VAE posterior

Flow model in latent space

$$\log q_K(\mathbf{z}_K|\mathbf{x}, \phi_*) = \log q(\mathbf{z}_0|\mathbf{x}, \phi) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

ELBO objective

$$\begin{aligned}\mathcal{L}(\phi, \theta) &= \mathbb{E}_{q_K(\mathbf{z}_K|\mathbf{x}, \phi_*)} \log \frac{p(\mathbf{x}, \mathbf{z}_K|\theta)}{q_K(\mathbf{z}_K|\mathbf{x}, \phi_*)} \\ &= \mathbb{E}_{q_K(\mathbf{z}_K|\mathbf{x}, \phi_*)} [\log p(\mathbf{x}, \mathbf{z}_K|\theta) - \log q_K(\mathbf{z}_K|\mathbf{x}, \phi_*)] \\ &= \mathbb{E}_{q_K(\mathbf{z}_K|\mathbf{x}, \phi_*)} \log p(\mathbf{x}|\mathbf{z}_K, \theta) - KL(q_K(\mathbf{z}_K|\mathbf{x}, \phi_*) || p(\mathbf{z}_K)).\end{aligned}$$

The second term in ELBO is reverse KL divergence. Planar flows was originally proposed for variational inference in VAE.

Flows in VAE posterior

Variational distribution

$$\log q_K(\mathbf{z}_K | \mathbf{x}, \phi_*) = \log q(\mathbf{z}_0 | \mathbf{x}, \phi) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

ELBO objective

$$\begin{aligned} \mathcal{L}(\phi, \theta) &= \mathbb{E}_{q_K(\mathbf{z}_K | \mathbf{x}, \phi_*)} [\log p(\mathbf{x}, \mathbf{z}_K | \theta) - \log q_K(\mathbf{z}_K | \mathbf{x}, \phi_*)] \\ &= \mathbb{E}_{q(\mathbf{z}_0 | \mathbf{x}, \phi)} [\log p(\mathbf{x}, \mathbf{z}_K | \theta) - \log q_K(\mathbf{z}_K | \mathbf{x}, \phi_*)] \Big|_{\mathbf{z}_K = g(\mathbf{z}_0, \{\phi_k\}_{k=1}^K)} \\ &= \mathbb{E}_{q(\mathbf{z}_0 | \mathbf{x}, \phi)} \left[\log p(\mathbf{x}, \mathbf{z}_K | \theta) - \log q(\mathbf{z}_0 | \mathbf{x}, \phi) + \right. \\ &\quad \left. + \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right| \right]. \end{aligned}$$

Flows in VAE posterior

Variational distribution

$$\log q_K(\mathbf{z}_K | \mathbf{x}, \phi_*) = \log q(\mathbf{z}_0 | \mathbf{x}, \phi) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

ELBO objective

$$\begin{aligned} \mathcal{L}(\phi, \theta) = \mathbb{E}_{q(\mathbf{z}_0 | \mathbf{x}, \phi)} & \left[\log p(\mathbf{x}, \mathbf{z}_K | \theta) - \log q(\mathbf{z}_0 | \mathbf{x}, \phi) + \right. \\ & \left. + \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right| \right]. \end{aligned}$$

- ▶ Obtain samples \mathbf{z}_0 from the encoder.
- ▶ Apply flow model $\mathbf{z}_K = g(\mathbf{z}_0, \{\phi_k\}_{k=1}^K)$.
- ▶ Compute likelihood for \mathbf{z}_K using the decoder, base distribution for \mathbf{z}_0 and the Jacobian.
- ▶ We do not need inverse flow function, if we use flows in variational inference.

Recap of previous lecture

ELBO

$$p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}(\boldsymbol{\phi}, \boldsymbol{\theta}) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})} \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi})} \rightarrow \max_{\boldsymbol{\phi}, \boldsymbol{\theta}}.$$

- ▶ Normal variational distribution
 $q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\boldsymbol{\phi}}(\mathbf{x}), \boldsymbol{\sigma}_{\boldsymbol{\phi}}^2(\mathbf{x}))$ is poor (e.g. has only one mode).
- ▶ Flows models convert a simple base distribution to a complex one using an invertible transformation with simple Jacobian.

Flow model in latent space

$$\log q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*) = \log q(\mathbf{z}_0|\mathbf{x}, \boldsymbol{\phi}) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \boldsymbol{\phi}_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

Let's use $q_K(\mathbf{z}_K|\mathbf{x}, \boldsymbol{\phi}_*)$, $\boldsymbol{\phi}_* = \{\boldsymbol{\phi}, \boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_K\}$ as a variational distribution. Here, $\boldsymbol{\phi}$ – encoder parameters, $\{\boldsymbol{\phi}_k\}_{k=1}^K$ – flow parameters.

Recap of previous lecture

Variational distribution

$$\log q_K(\mathbf{z}_K | \mathbf{x}, \phi_*) = \log q(\mathbf{z}_0 | \mathbf{x}, \phi) - \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right|.$$

ELBO objective

$$\begin{aligned} \mathcal{L}(\phi, \theta) = \mathbb{E}_{q(\mathbf{z}_0 | \mathbf{x}, \phi)} & \left[\log p(\mathbf{x}, \mathbf{z}_K | \theta) - \log q(\mathbf{z}_0 | \mathbf{x}, \phi) + \right. \\ & \left. + \sum_{k=1}^K \log \left| \det \left(\frac{\partial g_k(\mathbf{z}_{k-1}, \phi_k)}{\partial \mathbf{z}_{k-1}} \right) \right| \right]. \end{aligned}$$

- ▶ Obtain samples \mathbf{z}_0 from the encoder.
- ▶ Apply flow model $\mathbf{z}_K = g(\mathbf{z}_0, \{\phi_k\}_{k=1}^K)$.
- ▶ Compute likelihood for \mathbf{z}_K using the decoder, base distribution for \mathbf{z}_0 and the Jacobian.
- ▶ We do not need an inverse flow function if we use flows in variational inference.

Inverse autoregressive flow (IAF)

$$\begin{aligned}\mathbf{x} = g(\mathbf{z}, \boldsymbol{\theta}) &\Rightarrow x_i = \tilde{\sigma}_i(\mathbf{z}_{1:i-1}) \cdot z_i + \tilde{\mu}_i(\mathbf{z}_{1:i-1}). \\ \mathbf{z} = f(\mathbf{x}, \boldsymbol{\theta}) &\Rightarrow z_i = (x_i - \tilde{\mu}_i(\mathbf{z}_{1:i-1})) \cdot \frac{1}{\tilde{\sigma}_i(\mathbf{z}_{1:i-1})}.\end{aligned}$$

Reverse KL for flow model



$$KL(p||\pi) = \mathbb{E}_{p(\mathbf{z})} \left[\log p(\mathbf{z}) - \log \left| \det \left(\frac{\partial g(\mathbf{z}, \boldsymbol{\theta})}{\partial \mathbf{z}} \right) \right| - \log \pi(g(\mathbf{z}, \boldsymbol{\theta})) \right]$$

- ▶ We don't need to think about computing the function $f(\mathbf{x}, \boldsymbol{\theta})$.
- ▶ Inverse autoregressive flow is a natural choice for using flows in VAE:


$$\mathbf{z}_0 = \boldsymbol{\sigma}(\mathbf{x}) \odot \boldsymbol{\epsilon} + \boldsymbol{\mu}(\mathbf{x}), \quad \boldsymbol{\epsilon} \sim \mathcal{N}(0, 1); \quad \sim q(\mathbf{z}_0|\mathbf{x}, \phi).$$

$$\mathbf{z}_k = \tilde{\boldsymbol{\sigma}}_k(\mathbf{z}_{k-1}) \odot \mathbf{z}_{k-1} + \tilde{\boldsymbol{\mu}}_k(\mathbf{z}_{k-1}), \quad k \geq 1; \quad \sim q_k(\mathbf{z}_k|\mathbf{x}, \phi, \{\phi_j\}_{j=1}^k).$$


Inverse autoregressive flow (IAF)



(a) Prior distribution



(b) Posteriors in standard VAE



(c) Posteriors in VAE with IAF

Dequantization

- ▶ Images are discrete data, pixels lie in the $[0, 255]$ integer domain (the model is $P(\mathbf{x}|\boldsymbol{\theta}) = \text{Categorical}(\boldsymbol{\pi}(\boldsymbol{\theta}))$).
- ▶ Flow is a continuous model (it works with continuous data \mathbf{x}).

By fitting a continuous density model to discrete data, one can produce a degenerate solution with all probability mass on discrete values.


How to convert a discrete data distribution to a continuous one?

Uniform dequantization

$$\mathbf{x} \sim \text{Categorical}(\boldsymbol{\pi})$$

$$\mathbf{u} \sim U[0, 1]$$

$$\mathbf{y} = \mathbf{x} + \mathbf{u} \sim \text{Continuous}$$



Uniform dequantization

Statement

Fitting continuous model $p(\mathbf{y}|\boldsymbol{\theta})$ on uniformly dequantized data $\mathbf{y} = \mathbf{x} + \mathbf{u}$, $\mathbf{u} \sim U[0, 1]$ is equivalent to maximization of a lower bound on log-likelihood for a discrete model:

$$P(\mathbf{x}|\boldsymbol{\theta}) = \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u}$$

Thus, the maximisation of continuous model log-likelihood on \mathbf{y} can't lead to the a collapse onto the discrete data (the objective is bounded above by the discrete model log-likelihood).

Proof

$$\begin{aligned} \log P(\mathbf{x}|\boldsymbol{\theta}) &= \log \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} \geq \\ &\geq \int_{U[0,1]} \log p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} = \log p(\mathbf{y}|\boldsymbol{\theta}). \end{aligned}$$

Dequantization

- ▶ Images are discrete data, pixels lie in the $\{0, 255\}$ integer domain (the model is $P(\mathbf{x}|\theta) = \text{Categorical}(\pi(\theta))$).
- ▶ Flow is a continuous model (it works with continuous data \mathbf{x}).

By fitting a continuous density model to discrete data, one can produce a degenerate solution with all probability mass on discrete values.


How to convert a discrete data distribution to a continuous one?

Uniform dequantization

$$\mathbf{x} \sim \text{Categorical}(\pi)$$

$$\mathbf{u} \sim U[0, 1]$$

$$\mathbf{y} = \mathbf{x} + \mathbf{u} \sim \text{Continuous}$$



Uniform dequantization

Statement


Fitting continuous model $p(\mathbf{y}|\boldsymbol{\theta})$ on uniformly dequantized data $\mathbf{y} = \mathbf{x} + \mathbf{u}$, $\mathbf{u} \sim U[0, 1]$ is equivalent to maximization of a lower bound on log-likelihood for a discrete model:

$$P(\mathbf{x}|\boldsymbol{\theta}) = \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u}$$

Proof

$$\begin{aligned}\mathbb{E}_{\pi} \log p(\mathbf{y}|\boldsymbol{\theta}) &= \int \pi(\mathbf{y}) \log p(\mathbf{y}|\boldsymbol{\theta}) d\mathbf{y} = \\ &= \sum \pi(\mathbf{x}) \int_{U[0,1]} \log p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} \leq \\ &\leq \sum \pi(\mathbf{x}) \log \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} = \\ &= \sum \pi(\mathbf{x}) \log P(\mathbf{x}|\boldsymbol{\theta}) = \mathbb{E}_{\pi} \log P(\mathbf{x}|\boldsymbol{\theta}).\end{aligned}$$

Variational dequantization



- ▶ $p(\mathbf{y}|\boldsymbol{\theta})$ assign uniform density to unit hypercubes $\mathbf{x} + U[0, 1]$ (left fig).
- ▶ Neural network density models are smooth function approximators (right fig).
- ▶ Smooth dequantization is more natural.

How to perform the smooth dequantization?

Flow++

Variational dequantization

Introduce variational dequantization noise distribution $q(\mathbf{u}|\mathbf{x})$ and treat it as an approximate posterior.

Variational lower bound

$$\begin{aligned}\log P(\mathbf{x}|\boldsymbol{\theta}) &= \left[\log \int q(\mathbf{u}|\mathbf{x}) \frac{p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u} \right] \geq \\ &\geq \int q(\mathbf{u}|\mathbf{x}) \log \frac{p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u} = \mathcal{L}(q, \boldsymbol{\theta}).\end{aligned}$$

Uniform dequantization bound

$$\log P(\mathbf{x}|\boldsymbol{\theta}) = \log \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} \geq \int_{U[0,1]} \log p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u}.$$

Variational lower bound

$$\mathcal{L}(q, \theta) = \int q(\mathbf{u}|\mathbf{x}) \log \frac{p(\mathbf{x} + \mathbf{u}|\theta)}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u}.$$

Let $\mathbf{u} = h(\epsilon, \phi)$ is a flow model with base distribution $\epsilon \sim p(\epsilon) = \mathcal{N}(0, \mathbf{I})$:

$$q(\mathbf{u}|\mathbf{x}) = p(h^{-1}(\mathbf{u}, \phi)) \cdot \left| \det \frac{\partial h^{-1}(\mathbf{u}, \phi)}{\partial \mathbf{u}} \right|.$$

Then

$$\log P(\mathbf{x}|\theta) \geq \mathcal{L}(\phi, \theta) = \int p(\epsilon) \log \left(\frac{p(\mathbf{x} + h(\epsilon, \phi)|\theta)}{p(\epsilon) \cdot \left| \det \frac{\partial h(\epsilon, \phi)}{\partial \epsilon} \right|^{-1}} \right) d\epsilon.$$


Variational lower

$$\log P(\mathbf{x}|\boldsymbol{\theta}) \geq \int p(\boldsymbol{\epsilon}) \log \left(\frac{p(\mathbf{x} + h(\boldsymbol{\epsilon}, \phi))}{p(\boldsymbol{\epsilon}) \cdot \left| \det \frac{\partial h(\boldsymbol{\epsilon}, \phi)}{\partial \boldsymbol{\epsilon}} \right|^{-1}} \right) d\boldsymbol{\epsilon}.$$


- ▶ If $p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})$ is also a flow model, it is straightforward to calculate stochastic gradient of this ELBO.
- ▶ Uniform dequantization is a special case of variational dequantization ($q(\mathbf{u}|\mathbf{x}) = U[0, 1]$). The gap between $\log P(\mathbf{x}|\boldsymbol{\theta})$ and the derived ELBO is $KL(q(\mathbf{u}|\mathbf{x})||p(\mathbf{u}|\mathbf{x}))$.
- ▶ In the case of uniform dequantization the model unnaturally places uniform density over each hypercube $\mathbf{x} + U[0, 1]$ due to inexpressive distribution q .

Table 1. Unconditional image modeling results in bits/dim

Model family	Model	CIFAR10	ImageNet 32x32	ImageNet 64x64
Non-autoregressive	RealNVP (Dinh et al., 2016)	3.49	4.28	—
	Glow (Kingma & Dhariwal, 2018)	3.35	4.09	3.81
	IAF-VAE (Kingma et al., 2016)	3.11	—	—
	Flow++ (ours)	3.08	3.86	3.69
Autoregressive	Multiscale PixelCNN (Reed et al., 2017)	—	3.95	3.70
	PixelCNN (van den Oord et al., 2016b)	3.14	—	—
	PixelRNN (van den Oord et al., 2016b)	3.00	3.86	3.63
	Gated PixelCNN (van den Oord et al., 2016c)	3.03	3.83	3.57
	PixelCNN++ (Salimans et al., 2017)	2.92	—	—
	Image Transformer (Parmar et al., 2018)	2.90	3.77	—
	PixelSNAIL (Chen et al., 2017)	2.85	3.80	3.52



(a) PixelCNN



(b) Flow++

VAE limitations

- ▶ Poor variational posterior distribution (encoder)

$$q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\phi}(\mathbf{x}), \boldsymbol{\sigma}_{\phi}^2(\mathbf{x})).$$

- ▶ Poor prior distribution

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ Poor probabilistic model (decoder)

$$p(\mathbf{x}|\mathbf{z}, \theta) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\theta}(\mathbf{z}), \boldsymbol{\sigma}_{\theta}^2(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\theta) - \mathcal{L}(q, \theta) = (?).$$

ELBO interpretations

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(\phi, \boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(\phi, \boldsymbol{\theta}).$$

$$\mathcal{L}(\phi, \boldsymbol{\theta}) = \int q(\mathbf{z}|\mathbf{x}, \phi) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x}, \phi)} d\mathbf{z}.$$

- Evidence minus posterior KL

$$\mathcal{L}(q, \boldsymbol{\theta}) = \log p(\mathbf{x}|\boldsymbol{\theta}) - KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$

- Average negative energy plus entropy

$$\begin{aligned}\mathcal{L}(q, \boldsymbol{\theta}) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) - \log q(\mathbf{z}|\mathbf{x}, \phi)] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) + \mathbb{H}[q(\mathbf{z}|\mathbf{x}, \phi)].\end{aligned}$$

- Average reconstruction minus KL to prior

$$\begin{aligned}\mathcal{L}(q, \boldsymbol{\theta}) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) + \log p(\mathbf{z}) - \log q(\mathbf{z}|\mathbf{x}, \phi)] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) - KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z})).\end{aligned}$$

ELBO surgery

$$\frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(q, \theta) = \frac{1}{n} \sum_{i=1}^n [\mathbb{E}_{q(\mathbf{z}|\mathbf{x}_i)} \log p(\mathbf{x}_i|\mathbf{z}, \theta) - KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z}))].$$

Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}],$$

- ▶ $q(\mathbf{z}) = \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i)$ – **aggregated** posterior distribution.
- ▶ $\mathbb{I}_q[\mathbf{x}, \mathbf{z}]$ – mutual information between \mathbf{x} and \mathbf{z} under empirical data distribution and distribution $q(\mathbf{z}|\mathbf{x})$.
- ▶ First term pushes $q(\mathbf{z})$ towards the prior $p(\mathbf{z})$.
- ▶ Second term reduces the amount of information about \mathbf{x} stored in \mathbf{z} .

ELBO surgery

Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}].$$

Proof

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})} d\mathbf{z} = \\ &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z})q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})q(\mathbf{z})} d\mathbf{z} = \int \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} + \\ &+ \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{q(\mathbf{z})} d\mathbf{z} = KL(q(\mathbf{z})||p(\mathbf{z})) + \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z})) \end{aligned}$$

ELBO surgery

Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}],$$

Proof (continued)

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z}_i)) = KL(q(\mathbf{z})||p(\mathbf{z})) + \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z}))$$

It could be shown (exercise):

$$\mathbb{I}_q[\mathbf{x}, \mathbf{z}] = \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z})) \in [0, \log n].$$

Summary

- ▶ Uniform dequantization is the simplest form of dequantization.
- ▶ Variational dequantization is a more natural type that was proposed in Flow++ model.
- ▶ The IWAE could get the tighter lower bound to the likelihood, but the training of such model becomes more difficult.
- ▶ The ELBO surgery reveals insights about a prior distribution in VAE. The optimal prior is the aggregated posterior.