

# Matching and Basis Rotation for the Intrinsic Unified Evolution Basis

Niccolò Laurenti

In this document we will explain how the matching and the basis rotation are performed in the Intrinsic Unified Evolution Basis.

## 1 Matching

Fist of all, we have to perform the matching between the basis vectors in the two different flavor schemes, i.e. the  $n_f$  and the  $n_f + 1$  flavor schemes. The matching of the gluon, light quark and heavy quark PDFs are the followings:

$$\begin{aligned} g^{(n_f+1)} &= A_{gg}^S g^{(n_f)} + A_{gq}^S \Sigma_{(n_f)}^{(n_f)} + A_{gH}^S h^{(n_f)} \\ l^{(n_f+1)} &= A_{qq}^{ns} l^{(n_f)} + \frac{1}{2n_f} A_{qg}^S g^{(n_f)} \quad \text{and the same for } \bar{l} \\ h^{(n_f+1)} &= \frac{1}{2} A_{Hg}^S g^{(n_f)} + \frac{1}{2} A_{Hq}^{ps} \Sigma_{(n_f)}^{(n_f)} + A_{HH} h^{(n_f)} \quad \text{and the same for } \bar{h} \end{aligned}$$

From the second relation we get that

$$\begin{aligned} l_+^{(n_f+1)} &= A_{qq}^{ns} l_+^{(n_f)} + \frac{1}{n_f} A_{qg}^S g^{(n_f)} \\ l_-^{(n_f+1)} &= A_{qq}^{ns} l_-^{(n_f)} \\ \Sigma_{(n_f)}^{(n_f+1)} &= A_{qg}^S g^{(n_f)} + A_{qq}^{ns} \Sigma_{(n_f)}^{(n_f)} \\ V_{(n_f)}^{(n_f+1)} &= A_{qq}^{ns} V_{(n_f)}^{(n_f)} \end{aligned}$$

while from the third relation we get that

$$\begin{aligned} h_+^{(n_f+1)} &= A_{Hg}^S g^{(n_f)} + A_{Hq}^{ps} \Sigma_{(n_f)}^{(n_f)} + A_{HH} h_+^{(n_f)} \\ h_-^{(n_f+1)} &= A_{HH} h_-^{(n_f)} \end{aligned}$$

The matching of the components  $\Sigma_{\Delta(n_f)}$ ,  $V_{\Delta(n_f)}$ ,  $T_i$ ,  $V_i$  are diagonal. For the  $V$ s this is trivial since they are composed by  $l_-$ . For  $\Sigma_{\Delta(n_f)}$  instead: being it defined as

$$\Sigma_{\Delta(n_f)}^{(n_f+1)} = \frac{n_d(n_f)}{n_u(n_f)} \sum_{i=1}^{n_u} u_{+i}^{(n_f+1)} - \sum_{i=1}^{n_d} d_{+i}^{(n_f+1)}$$

the gluon contribution cancels, giving the relation

$$\Sigma_{\Delta(n_f)}^{(n_f+1)} = A_{qq}^{ns} \Sigma_{\Delta(n_f)}^{(n_f+1)}$$

The same holds for the  $T_i$  components.

Observe that this holds up to NNLO, since that at N<sup>3</sup>LO we have to consider also the pure singlet components of the light quark matching (I think that we have to add  $\frac{1}{2n_f} A_{qq}^{ps} \Sigma_{(n_f)}^{nf}$  to the matching of  $l^{(n_f+1)}$ , but I'm not 100% sure. In this way the matching of  $l_-$  remains diagonal, as it should be, and the same hold for  $\Sigma_{\Delta(n_f)}$  and  $T_i$ ).

Up to second order the perturbative expansion of the matching terms is given by

$$\begin{aligned} A_{gg}^S &= 1 + a_s A_{gg}^{S(1)} + a_s^2 A_{gg}^{S(2)} \\ A_{gq}^S &= 1 + a_s^2 A_{gq}^{S(2)} \\ A_{gH}^S &= 1 + a_s A_{gH}^{S(1)} \\ A_{qq}^S &= 1 \\ A_{qq}^{ns} &= 1 + a_s^2 A_{qq}^{ns(1)} \\ A_{Hg}^S &= 1 + a_s A_{Hg}^{S(1)} + a_s^2 A_{Hg}^{S(2)} \\ A_{Hq}^{ps} &= 1 + a_s^2 A_{Hq}^{ps(2)} \\ A_{HH} &= 1 + a_s A_{HH}^{(2)} \end{aligned}$$

## 2 Basis Rotation

After the matching we have to perform the basis rotation from the basis with  $\Sigma_{(n_f)}^{(n_f+1)}$ ,  $\Sigma_{\Delta(n_f)}^{(n_f+1)}$ ,  $h_+^{(n_f+1)}$  to the basis with  $\Sigma_{(n_f+1)}^{(n_f+1)}$ ,  $\Sigma_{\Delta(n_f+1)}^{(n_f+1)}$ ,  $T_i^{(n_f+1)}$  (all the considerations that we will do for this basis rotation apply identically to the components  $V$ ,  $V_\Delta$ ,  $h_-$ ,  $V_i$ ). Being all the PDFs in the  $n_f + 1$  flavor scheme, from now on we will drop the superscript  $(n_f + 1)$ .

### 2.1 $\Sigma$

For the  $\Sigma$  component the basis rotation is very simple, being

$$\Sigma_{(n_f+1)} = \Sigma_{(n_f)} + h_+$$

### 2.2 $\Sigma_\Delta$

This component requires a bit more work: starting from

$$\begin{cases} \Sigma &= \Sigma_u + \Sigma_d \\ \Sigma_\Delta &= \frac{n_d}{n_u} \Sigma_u - \Sigma_d \end{cases}$$

we obtain that

$$\begin{cases} \Sigma_u &= \frac{n_u}{n_f} (\Sigma + \Sigma_\Delta) \\ \Sigma_d &= \frac{n_d}{n_f} \Sigma - \frac{n_u}{n_f} \Sigma_\Delta \end{cases}$$

Therefore, we find

$$\Sigma_{\Delta(n_f+1)} = \frac{n_u(n_f+1)}{n_d(n_f+1)} \Sigma_{u(n_f)} - \Sigma_{d(n_f)} + k(n_f) h_+$$

where

$$k(n_f) = \begin{cases} \frac{n_u(n_f+1)}{n_d(n_f+1)} & \text{if h=up-like} \\ -1 & \text{if h=down-like} \end{cases}$$

In the end we find that

$$\Sigma_{\Delta(n_f+1)} = \left( \frac{n_d(n_f+1)}{n_u(n_f+1)} n_u(n_f) - n_d(n_f) \right) \Sigma_{(n_f)} + \frac{n_f+1}{n_u(n_f+1)} \frac{n_u(n_f)}{n_f} \Sigma_{\Delta(n_f)} + k(n_f) h_+$$

### 2.3 $T_i$

In the end, we have to find the rotation for the  $T_i$  component: being

$$\begin{aligned} T_3^d &= d^+ - s^+ = \Sigma_{d(n_f)} - h_+ & \text{for } n_f = 3 \\ T_3^u &= u^+ - c^+ = \Sigma_{u(n_f)} - h_+ & \text{for } n_f = 4 \\ T_8^d &= d^+ + s^+ - 2b^+ = \Sigma_{d(n_f)} - 2h_+ & \text{for } n_f = 5 \\ T_8^u &= u^+ + c^+ - 2t^+ = \Sigma_{u(n_f)} - 2h_+ & \text{for } n_f = 6 \end{aligned}$$

Using the expressions of  $\Sigma_u$  and  $\Sigma_d$  as a function of  $\Sigma$  and  $\Sigma_\Delta$ , we can write that

$$T_i = f_1(n_f) \Sigma_{(n_f)} + f_2(n_f) \Sigma_{\Delta(n_f)} + f_3(n_f) h_+$$

with

$$\begin{aligned} f_1(n_f) &= \begin{cases} \frac{n_u(n_f)}{n_f} & \text{if } h \text{ is up-like } (n_f=3,5) \\ \frac{n_d(n_f)}{n_f} & \text{if } h \text{ is down-like } (n_f=2,4) \end{cases} \\ f_2(n_f) &= \begin{cases} \frac{n_u(n_f)}{n_f} & \text{if } h \text{ is up-like} \\ -\frac{n_u(n_f)}{n_f} & \text{if } h \text{ is down-like} \end{cases} \\ f_3(n_f) &= \begin{cases} -1 & \text{if } h \text{ is } s, c \text{ } (n_f=2,3) \\ -2 & \text{if } h \text{ is } b, t \text{ } (n_f=4,5) \end{cases} \end{aligned}$$