Technical Indicator Assembly Document for NOAA NaMES

Willem Klajbor, The NOAA Ecosystem Indicators Working Group

2021-07-08

Contents

O.	vervi	ew 7	7												
	0.1	Definition of Indicators	7												
1	Chl	Chlorophyll-a 9													
	1.1	Data)												
	1.2	Methods)												
2	Zoo	plankton 11	L												
	2.1	Data	L												
	2.2	Methods													
3	Cor	al Reefs 13	3												
	3.1	Data	}												
	3.2	Methods	3												
4	For	Forage Fish 15													
	4.1	Data	ó												
	4.2	Methods	ó												
5	Seabirds 1														
	5.1	Data	7												
	5.2	Methods	7												
6	Ove	erfished Stocks)												
	6.1	Data)												
	6.2	Methods 10)												

7	Marine 1	Aam i	mals																			23
	7.1 ESA																					23
	7.2 MMI	PA																				23
8	Unusual	Unusual Mortality Events 2														25						
	8.1 Data																					25
	8.2 Meth	ods .																		 •		25
9	Sea Surfa	асе Т	lemp	era	tur	re																27
	9.1 Data																					27
	9.2 Meth	ods .																•	•		•	27
10	Sea Leve	1																				29
	10.1 Data																					29
	10.2 Meth	ods .																		 •	•	29
11	Sea Ice																					31
	11.1 Data																					31
	11.2 Meth	ods .																				32
12	Climate	Indic	es																			33
	12.1 ENS	Ο																				33
	12.2 MEI																					33
	12.3 PDO	٠																				33
	12.4 EPN	Р																				33
	12.5 NAO	٠																				33
	12.6 AMC)																		 •	•	34
13	Coastal I	opu ²	latio	n																		35
	13.1 Data																					35
	13.2 Meth	iods .																	•			35
14	Coastal	Гоuri	ism																			37
	14.1 Data																					37
	14.2 Meth	ods .																		 		37

5

15	Coastal Employment	39
	15.1 Data	39
	15.2 Methods	39
16	Commercial Fishing	41
	16.1 Landings	41
	16.2 Revenue	41
17	Recreational Fishing	43
	17.1 Effort	43
	17.2 Harvest	43
18	Fishing Engagement	45
	18.1 Commercial	45
	18.2 Recreational	45
19	Billion Dollar Disasters	47
	19.1 Data	47
	19.2 Methods	47
20	Beach Closures	51
	20.1 Data	51
	20.2 Methods	51
21	Marine Species Distribution	53
	21.1 Data	53
	21.2 Methods	53

6 CONTENTS

Overview

The National Marine Ecosystem Status web portal provides the status of marine ecosystems across the U.S. and access to NOAA ecosystem indicator information and data. This website is designed to document the data sources and methods used to create the indicators displayed on the site.

0.1 Definition of Indicators

Ecosystem indicators are quantitative and/or qualitative measures of key components of the ecosystem. Marine ecosystems provide food, jobs, security, well-being, and other services to millions of people across the U.S. Yet, marine ecosystems and the people that rely on them are facing increasingly complex challenges. Tracking the status and trends of ocean and coastal ecosystems is critically important to understand how these ecosystems are changing and identify potential issues.

8 CONTENTS

Chlorophyll-a

1.1 Data

Under Construction

1.2 Methods

Zooplankton

2.1 Data

Under Construction

2.2 Methods

Coral Reefs

3.1 Data

Under Construction

3.2 Methods

Forage Fish

4.1 Data

Under Construction

4.2 Methods

Seabirds

5.1 Data

Under Construction

5.2 Methods

Overfished Stocks

Fish play an important role in marine ecosystems, supporting the ecological structure of many marine food webs. Caught by recreational and commercial fisheries, fish support significant parts of coastal economies, and can play an important cultural role in many regions. To understand the health of fish populations - as well as their abundance and distribution, we regularly assess fish stocks - stock assessments. Assessments let us know if a stock is experiencing overfishing or if it is overfished i.e. how much catch is sustainable while maintaining a healthy stock. And, if a stock becomes depleted, stock assessments can help determine what steps may be taken to rebuild it to sustainable levels. Understanding stock assessments helps measure how well we're managing and recovering fish stocks over time.

We present the number of overfished stocks by year in all regions.

6.1 Data

Data are obtained from the NOAA Fisheries Fishery Stock Status website https://www.fisheries.noaa.gov/national/population-assessments/fishery-stock-status-updates. Stocks that meet the criteria for overfished status are summed by year for each region. The status of stocks are available in report form and graphically.

6.2 Methods

This indicator is compiled manually.

After traveling to https://www.fisheries.noaa.gov/national/population-assessments/fishery-stock-status-updates, identify the most up to date status

report. The reports should be available in two formats - through a report and through a visual.

The easiest way to compile the data are using the visual, which should be named "Stocks on the Overfished and Overfishing Lists by Region." After opening the visual (example below), overfished stocks will be displayed by squares and sorted spatially by region. We are only counting the "Overfished" stocks - do not count stocks that are only on the "Overfishing" list.

knitr::include_graphics(rep("overfished.png"))

North Pacific corresponds to the Alaska Region, Pacific corresponds to the California Current, Western Pacific corresponds to Hawaii (but be sure not to count the pacific island specific complexes), and New England corresponds to North Atlantic.

For more information, contact Willem Klajbor (willem.klajbor@noaa.gov) or Stephanie Oakes (stephanie.oakes@noaa.gov).

6.2. METHODS 21




Figure 6.1: NOAA Fisheries 2020 Q4 Stock Status Map

Marine Mammals

7.1 ESA

7.1.1 Data

Under Construction

7.1.2 Methods

 ${\bf Under\ construction}$

7.2 MMPA

7.2.1 Data

Under Construction

7.2.2 Methods

Unusual Mortality Events

8.1 Data

Under Construction

8.2 Methods

Sea Surface Temperature

9.1 Data

Under Construction

9.2 Methods

Sea Level

10.1 Data

Under Construction

10.2 Methods

Sea Ice

Unlike icebergs, glaciers, ice sheets, and ice shelves, which originate on land, sea ice forms, expands, and melts in the ocean. Sea ice influences global climate by reflecting sunlight back into space. Because this solar energy is not absorbed into the ocean, temperatures nearer the poles remain cool. When sea ice melts, the surface area reflecting sunlight decreases, allowing more solar energy to be absorbed by the ocean, causing temperatures to rise. This creates a positive feedback loop. Warmer water temperatures delay ice growth in the autumn and winter, and the ice melts faster the following spring, exposing dark ocean waters for longer periods the following summer.

Sea ice affects the movement of ocean waters. When sea ice forms, ocean salts are left behind. As the seawater gets saltier, its density increases, and it sinks. Surface water is pulled in to replace the sinking water, which in turn becomes cold and salty and sinks. This initiates deep-ocean currents driving the global ocean conveyor belt.

Sea ice is an important element of the Arctic system. It provides an important habitat for biological activity, i.e. algae grows on the bottom of sea ice, forming the basis of the Arctic food web, and it plays a critical role in the life cycle of many marine mammals - seals and polar bears. Sea ice also serves a critical role in supporting Indigenous communities culture and survival. We present the annual sea ice extent in millions of Kilometers for the Arctic region.

11.1 Data

Sea ice data was accessed from the NOAA National Centers for Environmental Information, https://www.ncdc.noaa.gov/snow-and-ice/extent/ , with the data pulled from here: https://www.ncdc.noaa.gov/snow-and-ice/extent/sea-ice/N/ $3/{\rm data.csv}$. The data are plotted in units of million square km.

11.2 Methods

To download the current sea ice data, you can either:

1) Copy/paste the following url into your web browser: https://www.ncdc.noaa.gov/snow-and-ice/extent/sea-ice/N/3/data.csv

or

2) Use the following R code to download the data and import it into your RStudio environment

```
url <-"https://www.ncdc.noaa.gov/snow-and-ice/extent/sea-ice/N/3/data.csv"
# Specify destination where file should be saved
destfile <- "C:/Users/ ... Your Path ... /my folder/output.csv"
#Apply download.file function in R
download.file(url, destfile)</pre>
```

Data were restructured and gauge values were calculated manually.

For more information, contact Willem Klajbor (willem.klajbor@noaa.gov) or Scott Cross (scott.cross@noaa.gov).

Climate Indices

12.1 ENSO

Under Construction

12.2 MEI

Under construction

12.3 PDO

Under Construction

12.4 EPNP

Under Construction

12.5 NAO

12.6 AMO

Coastal Population

13.1 Data

Under Construction

13.2 Methods

Coastal Tourism

14.1 Data

Under Construction

14.2 Methods

Coastal Employment

15.1 Data

Under Construction

15.2 Methods

Commercial Fishing

16.1 Landings

16.1.1 Data

Under Construction

16.1.2 Methods

Under construction

16.2 Revenue

16.2.1 Data

Under Construction

16.2.2 Methods

Recreational Fishing

17.1 Effort

17.1.1 Data

Under Construction

17.1.2 Methods

Under construction

17.2 Harvest

17.2.1 Data

Under Construction

17.2.2 Methods

Fishing Engagement

18.1 Commercial

18.1.1 Data

Under Construction

18.1.2 Methods

Under construction

18.2 Recreational

18.2.1 Data

Under Construction

18.2.2 Methods

Billion Dollar Disasters

In the United States the number of weather and climate-related disasters exceeding 1 billion dollars has been increasing since 1980. These events have significant impacts to coastal economies and communities. The Billion Dollar Disaster indicator provides information on the frequency and the total estimated costs of major weather and climate events that occur in the United States. This indicator compiles the annual number of weather and climate-related disasters across seven event types. Events are included if they are estimated to cause more than one billion U.S. dollars in direct losses. The cost estimates of these events are adjusted for inflation using the Consumer Price Index (CPI) and are based on costs documented in several Federal and private-sector databases. We present the total annual number of disaster events for all regions.

19.1 Data

Billion dollar disaster event frequency data are taken from NOAA's National Centers for Environmental Information (https://www.ncdc.noaa.gov/billions/). The number of disasters within each region were summed for every year of available data. Although the number is the count of unique disaster events within a region, the same disaster can impact multiple regions, meaning a sum across regions will overestimate the unique number of disasters.

19.2 Methods

The Billion Dollar Event Frequency Data displayed on the website were compiled using the following code:

```
PKG <- c("foreign", "stringr", "data.table")</pre>
for (p in PKG) {
  if(!require(p,character.only = TRUE)) {
    install.packages(p)
    require(p, character.only = TRUE)}
#states <- c("AK", "AL", "AR", "AZ", "CA", "CO", "CT", "DE", "FL", "GA", "HI",
              "IA", "ID", "IL", "IN", "KS", "KY", "LA", "MA", "MD", "ME", "MI",
              "MN", "MO", "MS", "MT", "NC", "ND", "NE", "NH", "NJ", "NM", "NV",
#
              "NY", "OH", "OK", "OR", "PA", "RI", "SC", "SD", "TN", "TX", "UT",
#
             "VA", "VT", "WA", "WI", "WV", "WY")
#
states <- c("AK","AL","CA","CT","DE","FL","GA","HI",
             "LA", "MA", "MD", "ME",
             "MS", "NC", "NH", "NJ",
             "NY", "OR", "PA", "RI", "SC", "TX",
             "VA", "WA", "PR", "VI")
#Update Year in URL (2021)
Billion_Storm <- NULL</pre>
for (x in states) {
  temp <- tempfile()</pre>
  temp.connect <- url(paste0("https://www.ncdc.noaa.gov/billions/events-",x,"-1980-202
  temp <- data.table(read.delim(temp.connect, header=TRUE,fill=FALSE, stringsAsFactors</pre>
  temp$State <- x
  Billion_Storm <- rbind(Billion_Storm,temp)</pre>
  unlink(temp)
  rm(temp)
}
Billion_Storm$Begin.Date <- as.character(Billion_Storm$Begin.Date)</pre>
Billion_Storm$Begin.Year <- substr(Billion_Storm$Begin.Date,1,4)</pre>
Billion_Storm$Begin.Date <- as.Date(Billion_Storm$Begin.Date,"%Y %m %d")
Billion_Storm$End.Date <- as.character(Billion_Storm$End.Date)</pre>
Billion_Storm$End.Year <- substr(Billion_Storm$End.Date,1,4)</pre>
Billion_Storm$End.Date <- as.Date(Billion_Storm$End.Date,"%Y %m %d")
Gulf.of.Mexico <- c("FL","AL","LA","MS","TX")</pre>
Northeast <- c("NC","VA","MD","DE","PA","NJ","NY","CT","RI",
                "MA","NH","ME")
Southeast <- c("SC", "GA", "FL")
California.Current <- c("CA", "OR", "WA")
Alaska<- c("AK")
```

19.2. METHODS 49

```
Hawaii <- c("HI")</pre>
Caribbean <- c("PR","VI")</pre>
Storm_Freq <- NULL</pre>
for (x in c("Gulf.of.Mexico","Northeast","Southeast","California.Current","Alaska","Hawaii","Carr
  TEMP <- Billion_Storm[which(Billion_Storm$State%in%get(x)),]</pre>
  TEMP$Disaster <- TEMP$Begin.Date <- TEMP$End.Date <- TEMP$Deaths <- TEMP$State <- TEMP$Begin.Ye
  TEMP <- unique(TEMP)</pre>
  colnames(TEMP)<- c('Name', 'Frequency', 'End.Year')</pre>
  TEMP <- aggregate(Frequency~End.Year, data=TEMP, FUN=length)</pre>
  TEMP$Region <- x
  assign(paste0(x,"_Data", sep=""),TEMP)
  Storm_Freq <- rbind(Storm_Freq,TEMP)</pre>
  rm(TEMP)
}
Storm_Freq_F <- spread(Storm_Freq,Region,Frequency)</pre>
write.csv(Storm_Freq_F,file="C:/Users/... your path.../Billion_Dollar_Storms_1980_Present.csv")
rm(list=ls())
```

Gauge values counted manually.

For more information, please contact Willem Klajbor (willem.klajbor@noaa.gov) or Kate Quigley (kate.quigley@noaa.gov).

Beach Closures

20.1 Data

Under Construction

20.2 Methods

Marine Species Distribution

21.1 Data

Under Construction

21.2 Methods

 ${\bf Under\ construction}$