

Enabling Community Contributions in Formulation Selection Decision Support for Continental Hydrologic Modeling

H31L-0764

OWP OFFICE OF WATER PREDICTION

Guy Litt^{1,2}, Lauren Bolotin^{1,2}, Benjamin Choat^{1,3}

¹NOAA-Affiliate, Office of Water Prediction, National Weather Service, NOAA, Tuscaloosa, AL ²Lynker, Boulder, CO ³WEST Consultants, Inc, Folsom, CA

How might we understand where streamflow predictive models work well or poorly for NextGen?

The **Regionalization and Formulations Testing & Selection (RaFTS)** analysis framework trains machine learning models on catchment attributes to make predictions on:

1. Estimated hydrologic formulation performance (e.g. KGE, NNSE) (Frame et al., in-review)
2. Dominant processes (e.g. Sobol' Sensitivity)
3. Hydrologic signatures (e.g. runoff ratio, FDC midsegment slope)


RaFTS relies on publicly available datasets:

1. Catchment attributes (e.g. NHDplus, hydroatlas)
2. Continental-scale reference catchments commonly modeled (e.g. CAMELS - Addor et al 2017)
3. Published hydrologic model results (e.g. Kratzert et al, 2019, Mai et al, 2022)



RaFTS enables hydrologic data-informed decision making across the hydrofabric: >800k catchments across CONUS + AK, HI, & PR

How RaFTS makes predictions




RaFTS: Open-source analytics tool for hydrologic model regionalization & decision making




Data-informed NextGen formulation selection

The importance of catchment attribute data vary by what's being predicted:




The amount of catchments needed to train a machine-learning model vary by what's being predicted:




Not all metrics or signatures are predictable with a given dataset:


Bias of FDC midsegment slope, Entity-aware LSTM test dataset predicted vs observed, multilayer perceptron




NNSE, Entity-aware LSTM test dataset predicted vs observed, random forest




Top 2% peak flow flow bias, Entity-aware LSTM test dataset predicted vs observed, multilayer perceptron (Poor predictability)



Top 2% peak flow bias, VIC basin validation dataset predicted vs observed, multilayer perceptron



The hydrologic formulations with the RaFTS best-predicted normalized Nash Sutcliffe Efficiency:



Current RaFTS Capabilities

- Standardize predictor datasets (e.g. metrics, hydrologic signatures)
- Retrieve and standardize basin attributes
 - USGS NHDplus attributes
 - hydroatlas
- Aggregate/transform standardized attributes
- Train machine learning models on attributes and predictors
 - Random Forest
 - Multi-layer Perceptron
- Grid search hyperparameter optimization
- Evaluate input datasets
 - Principal component analysis
 - Feature importance
- Evaluate machine learning results
 - Learning curve
 - Testing vs. observed regression
 - Prediction maps
- Make predictions in any basin with a USGS comid and retrievable basin attributes

Open source development of RaFTS for community use:

<https://github.com/NOAA-OWP/formulation-selector>

Possible Future Work:

- Uncertainty quantification
- Containerization
- Optimal best model(s) selector
- Regional-scale analyses
- Cloud-hosted data



Summary

Community-contributed outcomes from continental-scale hydrologic modeling results further improves data-informed decision making using RaFTS, an analysis framework for the community.

RaFTS can predict which hydrologic models may perform best based on a variety of metrics, such as NSE, KGE, high-flow metrics, baseflow metrics, etc, but **requires careful assessment of appropriate predictive ML models.**

Future applications to RaFTS might:

- Decide which processes should be represented & where, in process-based hydrologic models
- Constrain parameter sets following calibration via hydrologic signatures
- Automate decision-making on regionalization strategies

ACKNOWLEDGEMENTS:

The authors would like to thank Sorush Sorourian, Keith Jennings, Nels Frazier, Andy Wood, and Jess Garrett for ideas and contributions surrounding RaFTS.

REFERENCES:

- Addor, N., Newman, A. J., Mizukami, N., & Clark, M. P. (2017). The CAMELS data set: catchment attributes and meteorology for large-sample studies. *Hydrology and Earth System Sciences*, 21(10), 5293-5313.
- Blodgett, D., Johnson, J.M., 2022, nhdplusTools: Tools for Accessing and Working with the NHDPlus, <https://doi.org/10.5066/P97AS8JD>
- Frame, J. M., Araki, R., Bhuiyan, S.A., Bindas, T., Rapp, J., Bolotin, L., Deardorff, E., Liu, Q., Haces-Garcia, F., Liao, M., Frazier, N., & Ogden, F. Machine learning for a heterogeneous water modeling framework. In review.
- Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., & Nearing, G. (2019). Towards learning universal, regional, and local hydrological behaviors via machine learning applied to large-sample datasets. *Hydrology and Earth System Sciences*, 23(12), 5089-5110.
- Mai, J., Craig, J. R., Tolson, B. A., & Arsenault, R. (2022). The sensitivity of simulated streamflow to individual hydrologic processes across North America. *Nature communications*, 13(1), 455.

CONTACT

Website: <https://water.noaa.gov>
Email: nws.nwc@noaa.gov

View my poster and other AGU materials



formulation-selector repo:

