Elliptic Problems in Nonsmooth Domains

Pierre Grisvard



Society for Industrial and Applied Mathematics Philadelphia

Elliptic Problems in Nonsmooth Domains

Books in the Classics in Applied Mathematics series are monographs and textbooks declared out of print by their original publishers, though they are of continued importance and interest to the mathematical community. SIAM publishes this series to ensure that the information presented in these texts is not lost to today's students and researchers.

Editor-in-Chief

Robert E. O'Malley, Jr., University of Washington

Editorial Board

John Boyd, University of Michigan
Susanne Brenner, Louisiana State University
Bernard Deconinck, University of Washington
Leah Edelstein-Keshet, University of British Columbia
William G. Faris, University of Arizona
Nicholas J. Higham, University of Manchester
Peter Hoff, University of Washington

Mark Kot, University of Washington
Peter Olver, University of Minnesota
Philip Protter, Cornell University
Matthew Stephens, The University of Chicago
Divakar Viswanath, University of Michigan
Gerhard Wanner, L'Université de Genève

Classics in Applied Mathematics

C. C. Lin and L. A. Segel, Mathematics Applied to Deterministic Problems in the Natural Sciences

Johan G. F. Belinfante and Bernard Kolman, A Survey of Lie Groups and Lie Algebras with Applications and Computational Methods

James M. Ortega, Numerical Analysis: A Second Course

Anthony V. Fiacco and Garth P. McCormick, Nonlinear Programming: Sequential Unconstrained Minimization Techniques

F. H. Clarke, Optimization and Nonsmooth Analysis

George F. Carrier and Carl E. Pearson, Ordinary Differential Equations

Leo Breiman, Probability

R. Bellman and G. M. Wing, An Introduction to Invariant Imbedding

Abraham Berman and Robert J. Plemmons, Nonnegative Matrices in the Mathematical Sciences

Olvi L. Mangasarian, Nonlinear Programming

*Carl Friedrich Gauss, Theory of the Combination of Observations Least Subject to Errors: Part One, Part Two, Supplement. Translated by G. W. Stewart

U. M. Ascher, R. M. M. Mattheij, and R. D. Russell, Numerical Solution of Boundary Value Problems for Ordinary Differential Equations

K. E. Brenan, S. L. Campbell, and L. R. Petzold, Numerical Solution of Initial-Value Problems in Differential-Algebraic Equations

Charles L. Lawson and Richard J. Hanson, Solving Least Squares Problems

J. E. Dennis, Jr. and Robert B. Schnabel, Numerical Methods for Unconstrained Optimization and Nonlinear Equations

Richard E. Barlow and Frank Proschan, Mathematical Theory of Reliability

Cornelius Lanczos, Linear Differential Operators

Richard Bellman, Introduction to Matrix Analysis, Second Edition

Beresford N. Parlett, The Symmetric Eigenvalue Problem

Richard Haberman, Mathematical Models: Mechanical Vibrations, Population Dynamics, and Traffic Flow

Peter W. M. John, Statistical Design and Analysis of Experiments

Tamer Başar and Geert Jan Olsder, Dynamic Noncooperative Game Theory, Second Edition

Emanuel Parzen, Stochastic Processes

Petar Kokotović, Hassan K. Khalil, and John O'Reilly, Singular Perturbation Methods in Control: Analysis and Design

Jean Dickinson Gibbons, Ingram Olkin, and Milton Sobel, Selecting and Ordering Populations: A New Statistical Methodology

James A. Murdock, Perturbations: Theory and Methods

^{*}First time in print.

Classics in Applied Mathematics (continued)

Ivar Ekeland and Roger Témam, Convex Analysis and Variational Problems

Ivar Stakgold, Boundary Value Problems of Mathematical Physics, Volumes I and II

1. M. Ortega and W. C. Rheinboldt, Iterative Solution of Nonlinear Equations in Several Variables

David Kinderlehrer and Guido Stampacchia, An Introduction to Variational Inequalities and Their Applications

F. Natterer, The Mathematics of Computerized Tomography

Avinash C. Kak and Malcolm Slaney, Principles of Computerized Tomographic Imaging

R. Wong, Asymptotic Approximations of Integrals

O. Axelsson and V. A. Barker, Finite Element Solution of Boundary Value Problems: Theory and Computation

David R. Brillinger, Time Series: Data Analysis and Theory

Joel N. Franklin, Methods of Mathematical Economics: Linear and Nonlinear Programming, Fixed-Point Theorems

Philip Hartman, Ordinary Differential Equations, Second Edition

Michael D. Intriligator, Mathematical Optimization and Economic Theory

Philippe G. Ciarlet, The Finite Element Method for Elliptic Problems

Jane K. Cullum and Ralph A. Willoughby, Lanczos Algorithms for Large Symmetric Eigenvalue Computations, Vol. I: Theory

M. Vidyasagar, Nonlinear Systems Analysis, Second Edition

Robert Mattheij and Jaap Molenaar, Ordinary Differential Equations in Theory and Practice

Shanti S. Gupta and S. Panchapakesan, Multiple Decision Procedures: Theory and Methodology of Selecting and Ranking Populations

Eugene L. Allgower and Kurt Georg, Introduction to Numerical Continuation Methods

Leah Edelstein-Keshet, Mathematical Models in Biology

Heinz-Otto Kreiss and Jens Lorenz, Initial-Boundary Value Problems and the Navier-Stokes Equations

J. L. Hodges, Jr. and E. L. Lehmann, Basic Concepts of Probability and Statistics, Second Edition

George F. Carrier, Max Krook, and Carl E. Pearson, Functions of a Complex Variable: Theory and Technique

Friedrich Pukelsheim, Optimal Design of Experiments

Israel Gohberg, Peter Lancaster, and Leiba Rodman, Invariant Subspaces of Matrices with Applications

Lee A. Segel with G. H. Handelman, Mathematics Applied to Continuum Mechanics

Rajendra Bhatia, Perturbation Bounds for Matrix Eigenvalues

Barry C. Arnold, N. Balakrishnan, and H. N. Nagaraja, A First Course in Order Statistics

Charles A. Desoer and M. Vidyasagar, Feedback Systems: Input-Output Properties

Stephen L. Campbell and Carl D. Meyer, Generalized Inverses of Linear Transformations

Alexander Morgan, Solving Polynomial Systems Using Continuation for Engineering and Scientific Problems

I. Gohberg, P. Lancaster, and L. Rodman, Matrix Polynomials

Galen R. Shorack and Jon A. Wellner, Empirical Processes with Applications to Statistics

Richard W. Cottle, Jong-Shi Pang, and Richard E. Stone, The Linear Complementarity Problem

Rabi N. Bhattacharya and Edward C. Waymire, Stochastic Processes with Applications

Robert J. Adler, The Geometry of Random Fields

Mordecai Avriel, Walter E. Diewert, Siegfried Schaible, and Israel Zang, Generalized Concavity

Rabi N. Bhattacharya and R. Ranga Rao, Normal Approximation and Asymptotic Expansions

Françoise Chatelin, Spectral Approximation of Linear Operators

Yousef Saad, Numerical Methods for Large Eigenvalue Problems, Revised Edition

Achi Brandt and Oren E. Livne, Multigrid Techniques: 1984 Guide with Applications to Fluid Dynamics, Revised Edition

Bernd Fischer, Polynomial Based Iteration Methods for Symmetric Linear Systems

Pierre Grisvard, Elliptic Problems in Nonsmooth Domains

Copyright © 2011 by the Society for Industrial and Applied Mathematics

This SIAM edition is a republication of the work first published by Pitman Publishing Inc., Marshfield, Massachusetts, in 1985.

10987654321

All rights reserved. Printed in the United States of America. No part of this book may be reproduced, stored, or transmitted in any manner without the written permission of the publisher. For information, write to the Society for Industrial and Applied Mathematics, 3600 Market Street, 6th Floor, Philadelphia, PA 19104-2688 USA.

Library of Congress Cataloging-in-Publication Data

Grisvard, P. (Pierre)

Elliptic problems in nonsmooth domains / Pierre Grisvard. -- SIAM ed.

p. cm. -- (Classics in applied mathematics; 69)

Originally published: Boston: Pitman Advanced Pub. Program, 1985.


Includes bibliographical references and index.

ISBN 978-1-611972-02-3 (pbk.)

1. Boundary value problems. Numerical solutions. 2. Differential equations, Elliptic. Numerical solutions. I. Title.

QA379.G74 2011 515'.782--dc23

2011028564



1985: To Catherine, Olivier, Béatrice, and Etienne



2011:

To the memory of our beloved father, from Béatrice, Etienne, & Olivier Grisvard

Contents

	Fore	word xiii
	Pref	ace xv
1		olev spaces 1
	1.1	Motivation 1
	1.2	Boundaries 4
		1.2.1 Graphs and manifolds 5
		1.2.2 Segment property and cone property 10
	1.3	Spaces 14
		1.3.1 Euclidean space 14
		1.3.2 Open subsets of the Euclidean space 16
		1.3.3 Manifolds 19
	1.4	—
		1.4.1 Multiplication and differentiation 20
		1.4.2 Density results 24
		1.4.3 Continuation, compactness and convexity
		inequalities 25
		1.4.4 Imbeddings 27
		1.4.5 Spaces defined on polygons 33
	1.5	
		1.5.1 Hyperplanes 36
		1.5.2 Polygons 42
		1.5.3 Maximal domains of elliptic operators 52
	1.6	
		1.6.1 Normal systems 62
	1.7	A model domain with a cut 74
2	Reg	ular second-order elliptic boundary value problems 81
_	_	Foreword 84
		Variational solution of special problems 84
		2.2.1 Existence and uniqueness 84
		2.2.2 Smoothness 87

X	CONTENTS
	2.3 A priori estimates 92
	2.3.1 An inequality based on the duality mapping 92
	2.3.2 An inequality in the half space 97
	2.3.3 A general a priori estimate 105
	2.4 Existence and uniqueness, the general case 111
	2.4.1 The basic result 111
	2.4.2 Applications of the Fredholm theory and
	the maximum principle 119
	2.5 Other kinds of solutions 128
	2.5.1 More on smoothness 128
	2.5.2 Very weak solution 129
3	Second-order elliptic boundary value problems in convex
	domains 132
	3.1 A priori estimates and the curvature of the boundary 133
	3.1.1 An identity based on integration by parts 133
	3.1.2 A priori inequalities for the Laplace operator
	revisited 138
	3.1.3 A priori inequalities for more general operators 1
	3.2 Boundary value problems in convex domains 147
	3.2.1 Linear boundary conditions 147
	3.2.2 Nonlinear boundary conditions (review) 151
	3.2.3 Nonlinear boundary conditions (continued) 156
	3.2.4 Oblique boundary conditions 167
	3.3 Boundary value problems in domains with turning points
	Second-order elliptic boundary value problems in polygons 18.
	4.1 Foreword 182
	4.2 A priori estimates in an infinite strip 184
	4.2.1 Explicit solution by Fourier transform and
	consequences 184
	4.2.2 L_p bounds for the second derivatives of the
	solution 189
•	4.3 Bounds in a polygon 194
	4.3.1 The L_2 case 194
	4.3.2 The L_p case $(p \neq 2)$ 199
4	4.4 The Fredholm alternative 208
	4.4.1 The semi-Fredholm properties 208
	4.4.2 The adjoint problem 217
	4.4.3 The Fredholm alternative for variational
	problems 226
	4.4.4 The Fredholm alternative for nonvariational
	problems 234

		CONTENTS
5	Mor	e singular solutions 249
	5.1	Behaviour of the derivatives of order higher than two 249
		5.1.1 Special data 250
		5.1.2 A trace theorem 256
		5.1.3 More singular solutions 261
	5.2	Operators with variable coefficients 265
6	Resi	ults in spaces of Hölder functions 274
	6.1	Foreword 274
	6.2	A brief review of Hölder spaces 275
	6.3	Regular second-order elliptic boundary value problems revisited 282

consequences A model fourth-order problem

Smoothness

- Introductory results
- 7.2 305 Singular solutions, the L_2 case
 - Kondratiev's method in weighted spaces 7.2.1

The Schauder inequality in an infinite strip

The Schauder inequality in a polygon and its

Getting rid of the weights

The Schauder inequality

287 Second-order elliptic problems in polygons revisited

295

- Singular solutions, the L_p case 328
 - 7.3.1 A priori inequalities
 - 7.3.2 335 Smoothness
 - 7.3.3 340 The related Stokes problem

Miscellaneous 345

6.3.1

6.3.2

6.4.1

6.4.2

- The Dirichlet problem for a strongly nonlinear equation
- 8.2 Some three-dimensional results (an outline)
 - 8.2.1 357 **Edges**
 - Conical points and vertices 361
- 8.3 372 The heat equation
- The numerical solution of elliptic problems with singularities 384
 - 8.4.1 Weighted spaces and mesh refinements
 - 394 Augmenting the space of trial functions
 - 8.4.3 Calculating the stress intensity factor 396

400 Bibliography

Foreword

Since the publication of Pierre Grisvard's monograph in 1985, the theory of elliptic problems in nonsmooth domains has become increasingly important for research in partial differential equations and their numerical solutions. While significant advances have occurred during the last two decades, Grisvard's monograph remains an excellent introduction to the subject and a good source for the basic material.

I had the good fortune to obtain one of the last available copies of this monograph in the early 1990s just before it went out of print, and it has become a regular reference in my work ever since. I am, therefore, delighted to be able to play a role in bringing this monograph back into print in the SIAM Classics in Applied Mathematics series. I hope a new generation of mathematicians will also find it useful.

Susanne C. Brenner Louisiana State University June 2011

Preface

In this book, we focus our attention on elliptic boundary value problems in domains with nonsmooth boundaries and problems with mixed boundary conditions. So far this topic has been mainly ignored. Indeed most of the available mathematical theories about elliptic boundary value problems deal with domains with very smooth boundaries; few of them deal with mixed boundary conditions. However, the majority of the elliptic boundary value problems which arise in practice are naturally posed in domains whose geometry is simple but not smooth. These domains are very often three-dimensional polyhedra. For the purpose of solving them numerically these problems are usually reduced to two-dimensional domains. Thus the domains are plane polygons and the boundary conditions are mixed. Accordingly this book is primarily intended for mathematicians working in the field of elliptic partial differential equations as well as for numerical analysts and users of such elliptic equations.

Perhaps the main feature of elliptic boundary value problems in a domain with smooth boundary is the so-called 'shift theorem'. Let us describe it on the simplest example, the Dirichlet problem for the Laplace equation. This will be our model problem throughout this introduction. Accordingly we consider a function u which is a solution of the equation

$$\Delta u = f \tag{1}$$

in a bounded open subset Ω of the two-dimensional Euclidean space R^2 . Here the function f is given and we assume that u coincides with some smooth given function g on the boundary Γ of Ω . The shift theorem can be phrased in the framework of either the Sobolev spaces or the Hölder spaces. Here, for simplicity, we describe only the Sobolev version.

We denote by $W_p^m(\Omega)$ the space of those functions defined in Ω whose derivatives up to the order m have their pth power integrable in Ω . We assume that p is strictly greater than 1 and is finite. For the time being, we also assume that the boundary of Ω is smooth, i.e. is a C^∞ manifold. Then when f is given in $W_p^m(\Omega)$, the corresponding solution u of the

xvi PREFACE

problem (1) belongs to $W_p^{m+2}(\Omega)$. In other words the order of the Sobolev space is shifted from m to m+2, by the inverse operator of Δ .

The particular case when p=2 has a simpler proof and is usually the only one needed by numerical analysts. However, the general case when p is allowed to differ from 2 (especially p large) is useful when one studies nonlinear boundary value problems by some kind of linearization or fixed point method. Most of the current error estimates for the numerical solution of an elliptic boundary value problem rely on this shift theorem. Therefore it is particularly important to know whether or not the same result holds for boundary value problems in a domain with a nonsmooth boundary.

From now on let us assume that Ω has one corner. For convenience we assume that this corner is at the origin of \mathbb{R}^2 and that, in some neighbourhood of the corner, Ω coincides with the sector

$$G = \{(r \cos \theta, r \sin \theta); r > 0, 0 < \theta < \omega\}$$

in the usual polar coordinates, where ω is the size of the angle at the origin. Otherwise we assume that Γ is smooth. For each positive integer k, we define a function u_k in the following way:

$$u_k = r^{k\pi/\omega} \sin(k\pi\theta/\omega)$$

when $k\pi/\omega$ is not an integer and

$$u_k = r^{k\pi/\omega} \{ \ln r \sin (k\pi\theta/\omega) + \theta \cos (k\pi\theta/\omega) \}$$

when $k\pi/\omega$ is an integer. It is readily seen that u_k is harmonic in Ω (thus $f_k = \Delta u_k = 0$) and that u_k coincides with a smooth function g_k on Γ . Indeed u_k vanishes on Γ near the origin when $k\pi/\omega$ is not an integer, while it vanishes on one side of G (for $\theta = 0$) and coincides with the polynomial $(-1)^k \omega r^{k\pi/\omega}$ on the other side of G (for $\theta = \omega$) when $k\pi/\omega$ is an integer. Consequently if the shift theorem were valid on Ω , u_k ought to belong to the intersection of all the Sobolev spaces on Ω . This would imply that u_k has all its derivatives of all orders continuous in the closure of Ω by the well-known Sobolev imbedding theorem. However, it is easy to check from the explicit formula above for u_k , that u_k is l times continuously differentiable if and only if l is strictly smaller that $k\pi/\omega$. A little extra work shows that u_k belongs to the Sobolev space $W_p^l(\Omega)$ if and only if its Sobolev exponent l-2/p is strictly smaller than $k\pi/\omega$, again.

So much for the shift theorem when Ω has a corner. Surprisingly enough, the functions u_k are all we need to formulate an alternative statement. Indeed, when f is given in $W_p^m(\Omega)$, the corresponding solution u of the problem (1) has the following property: there exist numbers c_k such that

$$u - \sum c_k u_k \in W_p^{m+2}(\Omega)$$

where the k in the summation ranges over all integers such that

$$\pi/\omega \leq k\pi/\omega \leq m+2-2/p$$
,

provided the Sobolev exponent m+2-2/p is not an integer itself. The limitation on k in the summation means that we exclude the u_k which belong to the space $W_p^{m+2}(\Omega)$. This result demonstrates that the solution has the usual regularity far from the corner while it describes accurately the behaviour near the corner of that part of the solution which does not belong to the required space.

The terms in the expansion of u above coincide with the terms in the formal power series derived by Lehman (1959).

The above modified version of the shift theorem does not express a regularity result in the whole of Ω . Thus the following question remains open: under which assumptions of f does the solution u belong to $W_p^{m+2}(\Omega)$? In other words when do the coefficients c_k vanish? These are continuous linear functionals of the data f and g. It turns out that they are local functionals if and only if $k\pi/\omega$ is an integer. This means that they only depend on the restriction of the data f and g to any neighbourhood of the corner. For instance we have

$$c_1 = f(0, 0)/\pi$$

when $\omega = \pi/2$. On the other hand when $k\pi/\omega$ is not an integer the functional c_k is global; this means that c_k may not vanish even when the data f and g are zero near the corner. As a consequence the functional c_k depends on the geometry of Ω far from the corner and it is not possible to make it explicit in such a general case.

Deriving similar modified shift theorems for various boundary value problems is what this book is about. Let us now proceed with a detailed description of the various chapters.

Chapter 1

The properties of the Sobolev spaces have been thoroughly investigated even when they are defined on very rough domains. We review the only properties we need without proofs and rely on the well-known book by Nečas (1967) for the proofs and references. In dealing with boundary value problems, one cannot skip a preliminary study of the boundary values of the functions belonging to Sobolev spaces. Very little is available about this question when the boundary is a polygon, although a complete answer has been given by Nikol'skii (1956, 1958), in the framework of slightly different spaces more suitable in the approximation theory. Accordingly we describe completely the boundary properties of

xviii PREFACE

functions belonging to Sobolev spaces on domains with polygonal boundaries. We include the proofs which turn out to be very similar to Nikol'skii's proofs. Some extensions of the classical Green formula are also worked out in the spirit of Lions and Magenes (1963) in the more general case of nonsmooth domains. This is why Chapter 1 is surprisingly long.

Chapter 2

As a first step toward the generalization of the classical shift theorem, we attempt to find the minimal assumptions under which one of the classical methods of proof can be worked out. Our technique is to look at the problem locally, flatten the boundary by a change of variables, freeze the coefficients and use partial Fourier transforms. Basically this is the method followed in Agmon et al. (1959). It turns out that the minimal assumption under which one obtains solutions in the Sobolev space $W_p^m(\Omega)$ is that the boundary Γ is of class C^m . This means that Γ can be locally represented as the graph of a C^m function. Actually one can allow a boundary of class $C^{m-1,1}$. Consequently a variational solution to a second-order boundary value problem is shown to belong to $W_p^2(\Omega)$ provided the boundary is at least of class $C^{1,1}$ This assumption does not allow a polygonal boundary. We recall that $C^{1,1}$ denotes the class of the functions with Lipschitz first derivatives.

Chapter 3

The classical method outlined above includes the proof of an *a priori* estimate which looks roughly like this:

$$\int_{\Omega} \left| \frac{\partial u}{\partial x_i \, \partial x_i} \right|^p \, \mathrm{d}x \le C \int_{\Omega} |\Delta u|^p \, \mathrm{d}x + \text{lower-order terms.}$$
 (2)

Usually we have very poor control of the constant C involved in this inequality. This is due to the local character of the method of proof. However in the case when p=2, an alternative proof based on integration by parts leads to a very accurate evaluation of the constant C. This is achieved under very general (possibly nonlinear) boundary conditions on u, in any n-dimensional domain. Such a proof (for the Dirichlet boundary condition) goes back to Caccioppoli (1950–51). It turns out that the constant C depends only on the negative part of the curvature of Γ (roughly speaking). This allows one to take limits with respect to the domain Ω and to prove some regularity results in general convex domains as well as in domains with turning points. Such a technique has been used for the first time by Kadlec (1964).

xix

Chapters 4 and 5

These chapters are devoted to the proof of a modified shift theorem similar to the one outlined at the beginning of this introduction for general boundary value problems for the Laplace equation in a plane polygon. On each side of the polygon, the condition is either a Dirichlet or a Neumann or an oblique boundary condition. In Chapter 4 we prove the regularity of the second derivatives of the solution, while Chapter 5 focuses on the higher derivatives. In addition, some boundary value problems involving operators with variable coefficients as well as nonhomogeneous operators are investigated.

Chapter 6

The same boundary value problems as in Chapters 4 and 5 are investigated in the framework of the spaces $C^{m,\sigma}(\bar{\Omega})$, i.e. the space of the functions which are m times continuously differentiable and whose derivatives of order m fulfil a uniform Hölder condition of order σ throughout Ω (0 < σ < 1).

Chapter 7

This chapter is focused on the Dirichlet problem for the biharmonic equation in a plane polygon. We have chosen this particular problem as our model fourth-order problem because of its importance in several physical questions (bending of plates, elasticity, fluid dynamics). Again we prove a suitably modified shift theorem in the Sobolev spaces $W_p^m(\Omega)$. We follow very closely the method of Kondratiev (1967a) when p=2. The shift theorem is also reformulated for the linear Stokes system and for the stationary Navier-Stokes equations in a plane polygon.

Chapter 8

This chapter includes miscellaneous topics all closely related to the content of the previous chapters.

First, the Dirichlet problem for a strongly nonlinear elliptic equation in a convex plane polygon is solved by applying a classical global inversion theorem following a work by Najmi (1978). The method relies strongly on the results of Chapters 4 and 5.

The method of Chapter 3 is adapted to the heat equation for a domain which is not time-like (with only one space variable for simplicity). Here we follow a work by Sadallah (1976, 1977, 1978).

The third section of Chapter 8 describes without complete proofs the few results about the behaviour of the solution of a second-order boundary value problem in a three-dimensional polyhedron.

Finally the fourth section is devoted to the consequences of the results of the previous chapters for the numerical analysis of boundary value problems.

Singular solutions like the u_k above have a strong polluting effect on the classical finite element methods. This difficulty is usually overcome in two main ways which are described in this section. The first consists (in a few words) in augmenting the usual spaces of trial functions by the addition of some of the singular solutions which have been explicitly calculated here.

The second relies on mesh refinements near the corners. Again the way the mesh has to be refined is governed by the behaviour of the singular solutions near the corners. We give here an analysis of the related error estimates.

In conclusion, let me acknowledge that this book has been strongly influenced by the outstanding paper by Kondratiev about general elliptic boundary value problems in domains with conical points.

I wish to express my gratitude to the many mathematical colleagues in the Universities of Algiers, Maryland and Nice, with whom I have had so many fruitful talks.

Finally I wish to express my sincere appreciation to Pitman Publishing for their most efficient handling of the publication of this book.

Nice August. 1984 P.G.