Fitting Cell 186

Idea 1: Initilize based on relative onset or offset like apearence and sharpness

Issues:
1. Some cells are early or late onset cells that look like offset cells but are not.
   If initilized as an offset cell it is very hard to have it fit correctly.

Idea 2: Try to fit cell 186 "semi-automatically" and then use the information that you get from that fit in order to inform future fitting

Issues:
1. Perhaps it is possilbe to run a "pre-fit" on a wide range of options and narrow down whether or not a cell is an onset of offset cell.
   However, this operation is realtively expensive, and could have memory issues.

Idea 3: Initialize all cells in 7 different set regiemes

Initialize Cells to be
* Sharp Onset
* Sharp Offset
* Sharp Both
* Broad Onset
* Broad Offset
* Broad Both
* Low FR relatively tonic

These options should cover all of our bases.

Issues:
1. Sharp cell's alpha STRF looks problematic for Offset responding cells.
2. Old Issue: By "ice-berging" onset cells you invoke a latency in the STRF.
   Instead, obtain the offset rates by -rate + 2*mean(onset) during the middle of the signal.
3. This avoids a lot of isses with edge affects in the signal and attempts to put onset and offset on the same scale with similar sharpnes

Using new offset STRFs, we see improved sharpness in offset and both responses