CRITICAL CARE NURSING · LECTURE 4 · QUESTION BANK

Neurological Disorders

الاضطرابات العصبية — بنك الأسئلة

92	6	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

فاينال 2021/2022 ترم أول (صور الجروب) · فاينال 2021/2022 ترم تاني (صور الجروب) · كويز 1 كريتكال – ترم أول 2023 · كويز 1 كريتكال 2022 · ميدترم ترم أول 2023/2024 (صور الجروب بالإجابات) · أسئلة سابقة · كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 فاينال 2021/2022 ترم تاني (صور الجروب)

A patient who opens his eyes in response to speech, speaks inappropriate words and moves to localize pain site will receive a GCS of:

a. 9 **b.** 10
c. 11 **d.** 12

2 فاينال 2021/2022 ترم تاني (صور الجروب)

Decorticate posturing refers to abnormal:

a. Rigid pronation of upper and lower extremities
b. Flaccid paralysis of extremities
c. Rigid extension and pronation
d. Flexion of upper extremities and extension of lower extremities

3 فاينال 2021/2022 ترم تاني (صور الجروب)

An acute stroke patient unable to sign the consent form because of inability to write most probably has:

a. Alexia **b.** Apraxia
c. Agraphia **d.** Hemianopia

4 فاينال 2021/2022 ترم تاني (صور الجروب)

To prevent injury in a patient who has hemianopia, the nurse educates the patient to:

a. Consume soft foods **b.** Scan the room sides
c. Perform range of motion **d.** Use anti-embolism stockings

5 Signs and symptoms of meningeal irritation include all of the following EXCEPT: فاينال 2021/2022 ترم أول (صور الجروب)

a. Aphasia and hemiparesis
b. Photophobia and lacrimation
c. Nuchal rigidity and headache
d. Kernig's and Brudzinski's signs

6 Which statement regarding psychomotor seizure is TRUE? فاينال 2021/2022 ترم أول (صور الجروب)

a. They are psychological in origin
b. They involve acts of random violence
c. The patient usually becomes unconscious
d. They involve repetitive behavior patterns

7 Myasthenia gravis is mainly characterized by: فاينال 2021/2022 ترم أول (صور الجروب)

a. Aphasia and peripheral sensory deficit
b. Uncoordinated motor and sensory control
c. Ascending paralysis and generalized weakness
d. Muscular weakness with exercise, improved with rest

8 While inserting a peripheral cannula on the right hand, the patient reaches over with his left hand to remove the noxious stimulus. This response is called: ميدترم ترم أول 2023/2024 (صور الجروب بالإجابات)

a. Decortication
b. Decerebration
c. Withdrawal
d. Localization

9 You would allocate which Glasgow Coma Scale score for a patient who is in a coma? كويز 1 كريتكال 2022

a. 6
b. 9
c. 12
d. 15

10 You would allocate ..... GCS for a patient whose eyes are closed but open to speech, says words that don't make sense, can't obey commands and moves to locate a central stimulus. كويز 1 كريتكال – ترم أول 2023

a. 12 (E3, V4, M5)
b. 8 (E2, V4, M2)
c. 11 (E3, V3, M5)
d. 10 (E3, V3, M4)

11 A patient who experienced a transient ischemic attack is at high risk for: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Stroke
b. Diabetes
c. Hypertension
d. Hypercholesterolemia

12 When assisting a patient with right-side hemiparesis and dysphagia with eating, it is very important to: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Keep head of bed less than 30 degrees
b. Prevent aspiration by thinning the liquids
c. Check for food collected in the right cheek
d. Extend the patient's neck upward while eating

13 Hemorrhagic stroke occurs as a result of which of the following? كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Alcohol abuse
b. Ruptured intracranial aneurysm
c. Acquired or inherited coagulopathy
d. Hemorrhagic transformation of ischemic infarct

14 Paralysis of the arm and leg on the same side of the body due to stroke or CVA refers to: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Paraplegia
b. Hemiplegia
c. Quadriplegia
d. Cardioplegia

15 The acute phase of ischemic stroke ranges from: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. 12 to 24 hrs
b. 24 to 48 hrs
c. 24 to 72 hrs
d. 48 hrs to 1 week

16 When a patient with a stroke finds difficulty with writing, this is called: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Agraphia
b. Alexia
c. Apraxia
d. Agnosia

17 The clinical manifestations of a patient with hemorrhagic stroke include all of the following EXCEPT: كويز 2 كريتكال – ترم أول 2024 (ACS & Stroke)

a. Severe headache
b. Vomiting
c. Hypotension
d. Hemiparesis

18 In promoting health maintenance for prevention of strokes, the highest risk for the most common type of stroke is present in: أسئلة سابقة

a. African Americans
b. Women who smoke
c. Individuals with hypertension and diabetes
d. Those who are obese with high dietary fat intake

19 A thrombus that develops in a cerebral artery does not always cause loss of neurologic function because: أسئلة سابقة

a. The body can dissolve atherosclerotic plaques as they form
b. Some brain tissues do not require constant blood supply
c. Circulation via the circle of Willis may provide blood supply to the affected area
d. Deficits occur only when major arteries are occluded

20 A patient's facial numbness and inability to speak disappeared while awaiting examination and he requests discharge. Why should the nurse emphasize treatment before leaving? أسئلة سابقة

a. Probably an asymptomatic lacunar stroke
b. Symptoms will return and worsen in 24 hours
c. Transient deficits result from small hemorrhages that clot off
d. Probably a TIA, which is a sign of progressive cerebrovascular disease

21 What are characteristics of a stroke caused by intracerebral hemorrhage? (Select all that apply) أسئلة سابقة

a. Carries a poor prognosis
b. Caused by rupture of a vessel
c. Strong association with hypertension
d. Commonly occurs during or after sleep
e. Creates a mass that compresses the brain

22 Which type of stroke is associated with endocardial disorders, has a rapid onset and is unrelated to activity? أسئلة سابقة

a. Embolic
b. Thrombotic
c. Intracerebral hemorrhage
d. Subarachnoid hemorrhage

23 What primarily determines the neurologic functions affected by a stroke? أسئلة سابقة

a. The amount of tissue involved
b. The rapidity of onset
c. The brain area perfused by the affected artery
d. Presence of collateral circulation

24 The patient has a lack of comprehension of both verbal and written language. Which communication difficulty does he have? أسئلة سابقة

a. Dysarthria
b. Fluent dysphasia
c. Receptive aphasia
d. Expressive aphasia

25 A patient with left hemiplegia is admitted. To determine size, location and whether the stroke is ischemic or hemorrhagic, the nurse anticipates: أسئلة سابقة

a. Lumbar puncture
b. Cerebral arteriogram
c. Magnetic resonance imaging (MRI)
d. CT scan with contrast

26 A carotid endarterectomy is considered for a patient with several TIAs. The nurse explains that it: أسئلة سابقة

a. Joins a superficial extracranial artery to an intracranial artery
b. Restores circulation after obstruction of a cerebral artery
c. Removes an atherosclerotic plaque in the carotid artery to prevent stroke
d. Opens a stenosis with a balloon and stent

27 The incidence of ischemic stroke in patients with TIAs is reduced with: أسئلة سابقة

a. Furosemide
b. Lovastatin
c. Daily low-dose aspirin
d. Nimodipine

28 What is the priority intervention in the ED for the patient with a stroke? أسئلة سابقة

a. IV fluid replacement
b. Osmotic diuretics
c. Initiation of hypothermia
d. Maintenance of respiratory function with a patent airway and oxygen

29 A ruptured cerebral aneurysm has been diagnosed. The nurse anticipates which treatment? أسئلة سابقة

a. Hyperventilation therapy
b. Surgical clipping of the aneurysm
c. Hyperosmotic agents
d. Thrombolytic therapy

30 During the acute phase of a stroke, which cardiovascular sign is seen as the body attempts to increase cerebral blood flow? أسئلة سابقة

a. Hypertension
b. Fluid overload
c. Cardiac dysrhythmias
d. S3 and S4 heart sounds

31 During the secondary assessment of a patient with stroke, what should be included? (Select all that apply) أسئلة سابقة

a. Gaze
b. Sensation
c. Facial palsy
d. Proprioception
e. Current medications
f. Distal motor function

32 Which nursing intervention is indicated for the patient with hemiplegia? أسئلة سابقة

a. Footboard to prevent plantar flexion
b. Sling to immobilize the affected arm
c. Each joint lower than the proximal joint
d. Patient performs passive ROM of the affected limb with the unaffected limb

33 A right-brain stroke patient has homonymous hemianopsia. Early in care the nurse should: أسئلة سابقة

a. Place objects on the right side within the field of vision
b. Approach from the left side
c. Place objects on the left side
d. Patch the affected eye

34 Four days after a stroke a patient is to start oral feeding. Before feeding, the nurse should first: أسئلة سابقة

a. Check the patient's gag reflex
b. Order a soft diet
c. Raise the head of bed
d. Evaluate swallowing of crushed ice

35 What is an appropriate food for a stroke patient with mild dysphagia? أسئلة سابقة

a. Fruit juices
b. Pureed meat
c. Scrambled eggs
d. Fortified milkshakes

36 A wife asks why her husband with hemorrhagic stroke did not receive tPA. Best response: أسئلة سابقة

a. He didn't arrive within the timeframe
b. Not everyone is eligible; has he had surgery?
c. Discuss with his doctor
d. The medication dissolves clots and could cause more bleeding in his brain

37 What is an appropriate intervention to promote communication in a patient with aphasia? أسئلة سابقة

a. Use gestures, pictures and music
b. Talk about familiar ADLs
c. Structure statements so no verbal response needed
d. Use flashcards with simple words

38 A right-hemisphere stroke patient has unilateral neglect. During rehabilitation the nurse should: أسئلة سابقة

a. Avoid positioning on the affected side
b. Place objects on the unaffected side
c. Teach the patient to care consciously for the affected side
d. Protect the affected side with pillows

39 A patient with right-sided hemiplegia: what does the nurse teach the family about behavior changes? أسئلة سابقة

a. Ignore undesirable behaviors
b. Provide directions verbally in small steps
c. Distract from inappropriate emotional responses
d. Supervise all activities

40 The nurse helps the family cope with long-term effects of stroke by: أسئلة سابقة

a. Informing that the patient needs help with all ADLs
b. Explaining prestroke behavior will return
c. Encouraging family therapy or support groups
d. Helping them understand residual damage to promote problem solving

41 Which intervention should the nurse delegate to the LPN when caring for a patient after acute stroke? أسئلة سابقة

a. Assess neurologic status
b. Assess gag reflex
c. Administer ordered antihypertensives and platelet inhibitors
d. Teach caregivers strategies for neglect

42 The most common arrhythmia associated with stroke is: أسئلة سابقة

a. Premature ventricular contractions
b. Premature atrial contractions
c. Heart block
d. Atrial fibrillation

43 What is the primary feature of a coma? أسئلة سابقة

a. Inability to breathe without assistance
b. Prolonged unconsciousness and failure to respond to stimuli
c. Severe headache and nausea
d. Loss of motor function in one side of the body

44 Which of the following is a systemic cause of coma? أسئلة سابقة

a. Brain hemorrhage
b. Stroke
c. Hepatic encephalopathy
d. Brain tumor

45 Which central cause can lead to coma? أسئلة سابقة

a. Hypoglycemia
b. Stroke
c. Hypoxia
d. Uremic encephalopathy

46 What is the main presentation in the clinical diagnosis of coma? أسئلة سابقة

a. Decreased consciousness level (DCL)
b. Increased heart rate
c. Abnormal respiratory rate
d. High fever

47 Which of the following is commonly seen on a CT scan for coma diagnosis? أسئلة سابقة

a. Hemorrhage or edema in the brain
b. Elevated liver enzymes
c. Dehydration of brain tissues
d. Electrolyte imbalance

48 Which laboratory test can suggest hepatic encephalopathy in a coma patient? أسئلة سابقة

a. Elevated ammonia levels
b. Elevated creatinine levels
c. Low T3 and T4 levels
d. Positive autoantibodies

49 What could leukocytosis (elevated WBC count) indicate in a coma patient? أسئلة سابقة

a. Infection or inflammation
b. Hypoxia
c. Brain hemorrhage
d. Hypoglycemia

50 What does a low T3 and T4 level in the thyroid profile suggest? أسئلة سابقة

a. Hypothyroidism (myxedema coma)
b. Hyperthyroidism
c. Thyroid storm
d. Hypoglycemia

51 Elevated creatinine and urea levels in the kidney profile are indicative of: أسئلة سابقة

a. Liver failure
b. Uremic encephalopathy
c. Electrolyte imbalance
d. Stroke

52 Which condition may be suggested by elevated bilirubin levels in a coma patient? أسئلة سابقة

a. liver disfunction
b. Stroke
c. Brain tumor
d. Autoimmune disorder

53 Which autoimmune profile finding suggests an increased risk of stroke? أسئلة سابقة

a. Positive Anti-dsDNA antibodies
b. Elevated creatinine levels
c. Positive Anti-phospholipid antibodies
d. Elevated bilirubin

54 In coma management, what is the first step in conservative management? أسئلة سابقة

a. Airway maintenance
b. Fluid resuscitation
c. Brain surgery
d. Antibiotic administration

55 Which of the following is a critical component of conservative management for bedridden coma patients? أسئلة سابقة

a. Intravenous nutrition only
b. Respiratory care
c. Emergency surgery
d. Immediate neurological assessment

56 How should a bedridden coma patient be positioned to prevent complications? أسئلة سابقة

a. Keep the head flat
b. Regularly reposition the patient
c. Position them in a sitting posture
d. Place them on their back permanently

57 What is the recommended angle for the head of the bed in ventilated coma patients? أسئلة سابقة

a. 0 degrees
b. 15 to 30 degrees
c. 30 to 45 degrees
d. 60 to 90 degrees

58 What is the purpose of a "Ventilator Bundle" in coma management? أسئلة سابقة

a. To prevent skin ulcers
b. To reduce the need for sedation
c. To prevent respiratory and circulatory complications
d. To ensure proper hydration

59 When is surgical decompression needed for coma patients? أسئلة سابقة

a. Only if the patient is in stable condition
b. When there is brain swelling, bleeding, or mass effect
c. After failed conservative management
d. For any trauma-related coma

60 What does a brain MRI with diffusion-weighted imaging primarily help diagnose in coma patients? أسئلة سابقة

a. Brain hemorrhage
b. Stroke or ischemia
c. Electrolyte imbalance
d. Liver failure

61 What does a positive ANA test indicate in a coma patient? أسئلة سابقة

a. Stroke
b. Liver failure
c. Autoimmune disorders like lupus
d. Brain tumors

62 How can electrolyte imbalances contribute to coma? أسئلة سابقة

a. By impairing brain function
b. By increasing blood viscosity
c. By causing brain swelling
d. By inducing stroke

63 What is the purpose of an echocardiogram in coma management? أسئلة سابقة

a. To assess brain function
b. To evaluate heart function
c. To diagnose stroke
d. To identify liver dysfunction

64 In a coma patient, a CBC showing thrombocytosis or thrombocytopenia suggests: أسئلة سابقة

a. Stroke
b. Liver dysfunction
c. Bleeding disorder or inflammation
d. Brain hemorrhage

65 What does "ventilator bundle" refer to in the context of coma management? أسئلة سابقة

a. Measures for preventing respiratory and circulatory complications in ventilated patients
b. A treatment for electrolyte imbalance
c. A medication plan for coma patients
d. A nutritional intervention for coma patients

66 In the management of coma patients, what is the role of oral care? أسئلة سابقة

a. To reduce the risk of infection
b. To improve circulation
c. To enhance brain function
d. To prevent malnutrition

67 How does hypothyroidism contribute to coma? أسئلة سابقة

a. By increasing blood viscosity
b. By causing myxedema coma due to low thyroid hormones
c. By inducing electrolyte imbalance
d. By leading to brain hemorrhage

68 What is the significance of brain contusion in coma patients? أسئلة سابقة

a. It is a type of brain tumor
b. It suggests stroke
c. It results from traumatic brain injury and can lead to coma
d. It is an indication of systemic infection

69 What condition is associated with elevated liver enzymes (AST, ALT) in coma patients? أسئلة سابقة

a. Hepatic encephalopathy
b. Stroke
c. Uremic encephalopathy
d. Brain edema

أسئلة سابقة

70 **What is the primary goal in the management of coma patients?**

a. To administer sedatives
b. To treat the underlying cause
c. To prevent further brain swelling
d. To improve cardiac function

أسئلة سابقة

71 **Which of the following is a key part of managing coma patients who are on mechanical ventilation?**

a. Keeping the patient sedated at all times
b. Preventing pressure ulcers
c. Avoiding repositioning
d. Limiting oxygen intake

أسئلة سابقة

72 **A brain hemorrhage in a coma patient can lead to which of the following complications?**

a. Severe dehydration
b. Brain swelling or pressure
c. Hypoglycemia
d. Hepatic encephalopathy COPD

أسئلة سابقة

73 **Which of the following is a common sign of cardiogenic shock in ACS?**

a. High blood pressure
b. Low blood pressure with signs of organ dysfunction
c. Normal heart rate and rhythm
d. Lack of chest pain Acute Stroke:

أسئلة سابقة

74 **What is the primary cause of ischemic stroke?**

a. Increased intracranial pressure
b. Embolism from the left side of the heart or thrombus on the artery wall
c. Low oxygen levels in the brain
d. Infection

أسئلة سابقة

75 **Which of the following is a common symptom of Acute Stroke?**

a. Severe headache and disturbance of consciousness
b. Fever and chills
c. Nausea and vomiting without any neurological deficits
d. Sudden loss of appetite

أسئلة سابقة

76 **Which imaging modality is used as the first step in diagnosing stroke to differentiate between ischemic and hemorrhagic stroke?**

a. MRI
b. CT scan
c. Duplex Ultrasound
d. Electrocardiogram (ECG)

أسئلة سابقة

77 **What is the purpose of a Diffusion-Weighted Imaging (DWI) MRI in stroke diagnosis?**

a. To detect brain hemorrhages
b. To detect early ischemic stroke
c. To check for arrhythmias
d. To evaluate carotid artery stenosis

78 How does echocardiography contribute to stroke management? أسئلة سابقة

a. Detects brain tissue damage
b. Identifies arrhythmias
c. Identifies sources of embolism (e.g., LA appendage thrombus or LV smoke)
d. Detects high blood pressure

79 What is the significance of Duplex Ultrasound in stroke management? أسئلة سابقة

a. To evaluate brain tissue damage
b. To assess for stenosis or blockages in carotid and basilar arteries
c. To detect arrhythmias
d. To measure blood oxygen levels

80 What is the initial management approach for acute stroke patients? أسئلة سابقة

a. Immediate anticoagulation therapy
b. Keep the airway open and assess vital signs
c. Immediate thrombolysis
d. Surgical intervention

81 Which medication is commonly used in medical management of ischemic stroke to prevent further clot formation? أسئلة سابقة

a. Thrombolytics
b. Aspirin or clopidogrel
c. Corticosteroids
d. Diuretics

82 Within how many hours should thrombolytic therapy (e.g., tPA) be administered to ischemic stroke patients for optimal benefit? أسئلة سابقة

a. Within 1 hour
b. Within 3-4.5 hours
c. Within 12 hours
d. Within 24 hours

83 What is the purpose of interventional thrombectomy in stroke management? أسئلة سابقة

a. To dissolve clots in smaller arteries
b. To remove a clot from a large blood vessel to restore blood flow
c. To prevent brain swelling
d. To treat brain aneurysms

84 Which of the following procedures is used to treat brain aneurysms? أسئلة سابقة

a. Interventional thrombectomy
b. Interventional coiling
c. Surgical decompression
d. Carotid endarterectomy

85 What is the role of surgical decompression in stroke management? أسئلة سابقة

a. To remove a clot from an artery
b. To reduce brain swelling by removing part of the skull
c. To repair a damaged blood vessel
d. To stop brain hemorrhage

86 Which of the following is a characteristic of hemorrhagic stroke? أسئلة سابقة

a. Sudden severe headache and increase in intracranial pressure
b. No significant increase in intracranial pressure
c. Partial occlusion of a cerebral artery
d. Embolic blockage from the heart

87 What is a major complication associated with ischemic stroke that may require thrombolytic therapy? أسئلة سابقة

a. Severe brain edema
b. Embolism from the heart
c. Aneurysm rupture
d. Increased intracranial pressure

88 What is the significance of early administration of thrombolytic therapy in ischemic stroke? أسئلة سابقة

a. To prevent hemorrhagic transformation
b. To restore blood flow and minimize brain damage
c. To reduce blood pressure
d. To stabilize heart function

89 How does MRI with diffusion-weighted imaging (DWI) aid in early detection of ischemic stroke? أسئلة سابقة

a. By detecting brain hemorrhages
b. By identifying early tissue damage caused by reduced blood flow
c. By measuring oxygen levels in the blood
d. By detecting arrhythmias

90 What is the primary cause of ischemic stroke from the left side of the heart? أسئلة سابقة

a. Aneurysm rupture
b. Atrial fibrillation or other arrhythmias causing embolism
c. High blood pressure
d. Carotid artery stenosis

91 Which of the following is a benefit of using anticoagulants such as aspirin or clopidogrel in ischemic stroke management? أسئلة سابقة

a. To dissolve existing clots
b. To prevent further clot formation
c. To increase cerebral perfusion
d. To reduce blood pressure

92 In which case would interventional thrombectomy be most appropriate for stroke patients? أسئلة سابقة

a. When the stroke is hemorrhagic
b. In patients with mild symptoms
c. In patients with large vessel occlusion and ischemic stroke
d. When blood pressure is elevated

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 **A patient with a TIA is at high risk for:** NCLEX

a. Diabetes
b. Stroke
c. Hypotension
d. Seizure only

N2 **Before giving thrombolytics for ischemic stroke, which finding must be reported?** NCLEX

a. BP 195/110 mmHg
b. HR 88
c. Temperature 37.2
d. RR 18

N3 **Select ALL signs of meningeal irritation.** SELECT ALL

a. Nuchal rigidity
b. Kernig's sign
c. Brudzinski's sign
d. Photophobia
e. Hemiparesis

N4 **A stroke patient with homonymous hemianopia should be taught to:** NCLEX

a. Use eye patch
b. Scan the environment by turning the head
c. Avoid reading
d. Lie flat

N5 **Before feeding a patient after stroke, the nurse first:** NCLEX

a. Raises HOB only
b. Checks the gag/swallow reflex
c. Orders soft diet
d. Gives water

N6 **Myasthenia gravis is characterized by:** NCLEX

a. Ascending paralysis
b. Muscle weakness increasing with activity and improving with rest
c. Sensory loss
d. Aphasia

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

TIA	Transient deficit; warning of stroke.	**Ischemic stroke**	Thrombotic or embolic (AF).
Hemorrhagic stroke	Ruptured aneurysm/vessel; HTN; no tPA.	**Acute phase**	24–72 hours.
Hemiplegia	Paralysis of one side.	**Agraphia**	Inability to write.
Receptive aphasia	Can't understand speech/writing.	**Homonymous hemianopia**	Teach scanning; objects in visual field.
Dysphagia	Check gag reflex; thick foods.	**Meningeal signs**	Nuchal rigidity, Kernig, Brudzinski, photophobia.

Psychomotor seizure	Automatisms, repetitive behavior.
Myasthenia gravis	Weakness with activity, better with rest.

Answer Key

الإجابات

1. c — 11

2. d — Flexion of upper extremities and extension of lower extremities

3. c — Agraphia

4. b — Scan the room sides

5. a — Aphasia and hemiparesis

6. d — They involve repetitive behavior patterns
الورقة متظللة على psychological in origin — الـ psychomotor (complex partial) seizure بيكون فيها automatisms = حركات متكررة (repetitive behavior). إجابة مقترحة.

7. d — Muscular weakness with exercise, improved with rest

8. d — Localization
الورقة معلّمة decortication — الصح localization: المريض بيحرّك إيده التانية ناحية مصدر الألم عشان يشيله (M5).

9. a — 6

10. c — 11 (E3, V3, M5)

11. a — Stroke
أسئلة الـ ACS/MI (برّه القائمة) اتشالت في الكويز ده.

12. c — Check for food collected in the right cheek

13. b — Ruptured intracranial aneurysm

14. b — Hemiplegia

15. c — 24 to 72 hrs

16. a — Agraphia

17. c — Hypotension

18. c — Individuals with hypertension and diabetes

19. c — Circulation via the circle of Willis may provide blood supply to the affected area

20. d — Probably a TIA, which is a sign of progressive cerebrovascular disease

21. a, b, c — Carries a poor prognosis; Caused by rupture of a vessel; Creates a mass that compresses the brain

22. a — Embolic

23. c — The brain area perfused by the affected artery

24. c — Receptive aphasia

25. c — Magnetic resonance imaging (MRI)
الإجابات من مفتاح الدفعة المكتوب على الملف. سؤال right/left brain matching اتشال.

26. c — Removes an atherosclerotic plaque in the carotid artery to prevent stroke

27. c — Daily low-dose aspirin

28. d — Maintenance of respiratory function with a patent airway and oxygen

29. b — Surgical clipping of the aneurysm

30. a — Hypertension

31. a, b, c, d, f — Gaze; Sensation; Facial palsy; Proprioception; Distal motor function

32. d — Patient performs passive ROM of the affected limb with the unaffected limb

33. a — Place objects on the right side within the field of vision

34. a — Check the patient's gag reflex

35. c — Scrambled eggs

36. d — The medication dissolves clots and could cause more bleeding in his brain

37. b — Talk about familiar ADLs

38. c — Teach the patient to care consciously for the affected side

39. c — Distract from inappropriate emotional responses

40. d — Helping them understand residual damage to promote problem solving

41. c — Administer ordered antihypertensives and platelet inhibitors

42. d — Atrial fibrillation

43. b — Prolonged unconsciousness and failure to respond to stimuli

44. c — Hepatic encephalopathy

45. b — Stroke

46. a — Decreased consciousness level (DCL)

47. a — Hemorrhage or edema in the brain

48. a — Elevated ammonia levels

49. a — Infection or inflammation

50. a — Hypothyroidism (myxedema coma)

51. b — Uremic encephalopathy

52. a — liver disfunction

53. c — Positive Anti-phospholipid antibodies

54. a — Airway maintenance

55. b — Respiratory care

56. b — Regularly reposition the patient

57. c — 30 to 45 degrees

58. c — To prevent respiratory and circulatory complications

59. b — When there is brain swelling, bleeding, or mass effect

60. b — Stroke or ischemia

61. c — Autoimmune disorders like lupus

62. a — By impairing brain function

63. b — To evaluate heart function

64. c — Bleeding disorder or inflammation

65. a — Measures for preventing respiratory and circulatory complications in ventilated patients

66. a — To reduce the risk of infection

67. b — By causing myxedema coma due to low thyroid hormones

68. c — It results from traumatic brain injury and can lead to coma

69. a — Hepatic encephalopathy

70. b — To treat the underlying cause

71. b — Preventing pressure ulcers

72. b — Brain swelling or pressure

73. b — Low blood pressure with signs of organ dysfunction

74. b — Embolism from the left side of the heart or thrombus on the artery wall

75. a — Severe headache and disturbance of consciousness

76. b — CT scan

77. b — To detect early ischemic stroke

78. c — Identifies sources of embolism (e.g., LA appendage thrombus or LV smoke)

79. b — To assess for stenosis or blockages in carotid and basilar arteries

80. b — Keep the airway open and assess vital signs

81. b — Aspirin or clopidogrel

82. b — Within 3-4.5 hours

83. b — To remove a clot from a large blood vessel to restore blood flow

84. b — Interventional coiling

85. b — To reduce brain swelling by removing part of the skull

86. a — Sudden severe headache and increase in intracranial pressure

87. b — Embolism from the heart

88. b — To restore blood flow and minimize brain damage

89. b — By identifying early tissue damage caused by reduced blood flow

90. b — Atrial fibrillation or other arrhythmias causing embolism

91. b — To prevent further clot formation

92. c — In patients with large vessel occlusion and ischemic stroke

N1. b — Stroke
TIA is a warning sign of stroke.

N2. a — BP 195/110 mmHg
BP >185/110 must be controlled before tPA due to bleeding risk.

N3. a, b, c, d — Nuchal rigidity; Kernig's sign; Brudzinski's sign; Photophobia
Hemiparesis is a focal deficit, not meningeal irritation.

N4. b — Scan the environment by turning the head
Scanning compensates for the lost visual field.

N5. b — Checks the gag/swallow reflex
Aspiration prevention.

N6. b — Muscle weakness increasing with activity and improving with rest
Autoimmune attack on acetylcholine receptors.