CRITICAL CARE NURSING · LECTURE 6 · QUESTION BANK

# Specific Traumas

إصابات محددة — بنك الأسئلة

| 19 | 6 | 12 |
|---|---|---|
| Past exam questions | NCLEX-style | Flashcards |

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

## Part A · Past Exam Questions

أسئلة الدفعات السابقة

**1** أسئلة سابقة

**The nurse is handling a client with a chest tube. Suddenly, the chest drainage system is accidentally disconnected, what is the most appropriate action for the nurse to take ?**

**a.** Secure the chest tube using a tape
**b.** Clamp the chest tube immediately
**c.** Place the end of the chest tube in a container of normal sterile saline
**d.** Apply an occlusive dressing and notify the physician

**2** أسئلة سابقة

**.....is a sign of tension Pneumothorax**

**a.** Decreased cardiac output
**b.** Hypertension
**c.** Tracheal deviation
**d.** Increased cardiac output

**3** أسئلة سابقة

**Which intervention should the nurse implement for a male client who has had a left- sided chest tube for six (6) hours and refuses to take deep breaths because it hurts too much?**

**a.** Medicate the client and have the client take deep breaths
**b.** Encourage the client to take shallow breaths to help with the pain
**c.** Explain that deep breaths do not have to be taken at this time
**d.** Tell the client that if he doesn't take deep breaths, he could die

**4** أسئلة سابقة

**The immediate management of tension pneumothorax is**

**a.** Emergency department thoracotomy
**b.** Chest tube placement in 7th intercostal space
**c.** Needle thoracentesis in the 2nd intercostal space
**d.** A three-sided dressing placement over chest tube

**5** أسئلة سابقة

**The client is admitted to the emergency department with chest trauma. When assess- ing the client, which signs/ symptoms would the nurse expect to find that support the diagnosis of pneumothorax ?**

**a.** Bronchovesicular lung sounds and bradypnea
**b.** Unequal lung expansion and dyspnea
**c.** Frothy bloody sputum and consolidation
**d.** Barrel chest and polycythemia

**6** أسئلة سابقة

**..... causes mediastinal shift and compression of the lungs**

**a.** Primary pneumothorax
**b.** Secondary pneumothorax
**c.** Traumatic pneumothorax
**d.** Tension pneumothorax

**7** أسئلة سابقة

**The nurse assesses a patient with chest injury who had flail chest for which of the following most distinctive sign ?**

**a.** Massive Peripheral Cyanosis
**b.** Severe Postural Hypotension
**c.** Paradoxical chest movement
**d.** Dyspnea, especially on exhalation

**8** أسئلة سابقة

**Nurse Ruth assessing a patient for tracheal displacement should know that the trachea will deviate toward the:**

**a.** Contralateral side in a simple pneumothorax
**b.** Affected side in a hemothorax
**c.** Affected side in a tension pneumothorax
**d.** Contralateral side in hemothorax. - A patient is admitted with a left mid-axillary chest wound and experiencing extreme dyspnea, tachycardia, and hypoxia. On assessment, there was unequal rise and fall of the chest with absent breath sounds on the left side. Also, a "sucking" sound was noticed when the patient inhales and exhales. The patient's chest x-ray showed a pneumothorax. Questions from 1-2are related to the above situation

**9** أسئلة سابقة

**The above patient is expected to have which of the following types of pneumothorax?**

**a.** Open
**b.** Closed
**c.** Tension
**d.** Spontaneous

**10** أسئلة سابقة

**A client with a chest injury has suffered flail chest. The nurse assesses the client for which most distinctive sign of flail chest ?**

**a.** Cyanosis
**b.** Hypotension
**c.** Paradoxical chest movement
**d.** Dyspnea, especially on exhalation

**11** أسئلة سابقة

**A patient was admitted to the ER after a motor vehicle accident with multiple fractured ribs. Respiratory assessment includes signs/symptoms of secondary pneumothorax, which includes which of the following ?**

**a.** Crackles over lung bases of affected lung
**b.** Tracheal deviation toward the affected lung
**c.** Sharp pleuritic pain that worsens on inspiration
**d.** Increased diaphragmatic excursion on side of rib fractures

**12** A client reports left-sided chest pain after playing racquetball. The client is hospitalized and diagnosed with left pneumothorax. When assessing the client's left chest area, the nurse expects to identify which finding ? أسئلة سابقة

**a.** Dull sound on percussion
**b.** Vocal fremitus on palpation
**c.** Rales with rhonchi on auscultation
**d.** Absence of breath sounds on auscultation

**13** During percussion of the chest wall, hyperresonance suggests which of the following ? أسئلة سابقة

**a.** Pneumothorax
**b.** Haemothorax
**c.** Pneumonia
**d.** Pleural effusion

**14** What type of pressure or painful stimuli should the nurse avoid in a post-traumatic brain injury patient who has sustained multiple fractures to the face and eyes who is unable to perform a motor command based on a verbal stimulus ? أسئلة سابقة

**a.** Fingernailbedpressure
**b.** Trapezius squeeze
**c.** Supraorbital pressure
**d.** Toenailbedpressure

**15** You have responded to a motor vehicle crash involving a small bus carrying (6) passengers and a driver and a tractor trailer with a driver. The bus appears to have rolled over and landed on its wheels in the roadway and the tractor trailer is jackknifed in a nearby ditch. A middle-aged man who has an open head injury with exposed brain tissue. He is displaying agonal respirations أسئلة سابقة

**a.** Red
**b.** Yellow
**c.** Green
**d.** Black

**16** While working in the ED, the following four patients arrived at the same time. Indicate how to sort these patients utilizing triage categories أسئلة سابقة

**a.** A patient with a cut wound in the forehead, mild pain in the left leg and can walk
**b.** A patient with back & neck cyanosis, with abdominal stab wound & open skull fracture
**c.** A patient with left femur open fracture, cold extremities, shallow breathing, & weak pulse
**d.** A patient with severe bleeding, is fully conscious with a respiratory rate of 30 br./min
**e.** 4, 2, 3, 1
**f.** 2, 1,3,4
**g.** 4, 3, 1, 2
**h.** 3, 4, 1, 2

**17** A victim with which of the following conditions would be sorted using a GREEN triage ? أسئلة سابقة

**a.** Bilateralfemur fractures
**b.** Is cool, pale&diaphoretic
**c.** Unconscious whosebreathingis 40br/min
**d.** Has abrasions on the whole left posterior leg

**18** A 50-year-old patient, who has bilateral fractured femurs, weak radial pulse and a respiratory rate of 24 br/min will be receive ..... triage tag أسئلة سابقة

**a.** Black
**b.** Green
**c.** Red
**d.** Yellow

**19** While working in the ED, the following four patients arrived at the same time. Indicate how to sort these patients .utilizing triage categories أسئلة سابقة

**a.** Apatient is layingsupinecomplaining of inabilityto feelor moveextremities
**b.** Apatient withopenfemurfracturethat is bleedingprofusely
**c.** Apatient withdislocatedshoulder that controlledbyatourniquet
**d.** Apatient withopenskullfracture andexposedbrainmatter
**e.** 1,3,2,4
**f.** 4,3,1,2
**g.** 2,1,3,4
**h.** 3,4,1,2

## Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

**N1** Tracheal deviation, absent breath sounds on one side and hypotension suggest: NCLEX

**a.** Simple pneumothorax
**b.** Tension pneumothorax
**c.** Hemothorax
**d.** Rib fracture

**N2** Paradoxical chest movement is characteristic of: NCLEX

**a.** Flail chest
**b.** Pneumonia
**c.** Hemothorax
**d.** Pleural effusion

**N3** Select ALL signs of cardiac tamponade (Beck's triad). SELECT ALL

**a.** Hypotension
**b.** Distended neck veins
**c.** Muffled heart sounds
**d.** Bradypnea

**N4** Kehr's sign (left shoulder pain) after abdominal trauma suggests: NCLEX

**a.** Liver injury
**b.** Splenic rupture
**c.** Bladder injury
**d.** Kidney injury

**N5** Continuous bubbling in the water-seal chamber of a chest drain indicates: NCLEX

**a.** Normal function
**b.** Air leak in the system
**c.** Re-expanded lung
**d.** Blocked tube

**N6** Priority for a patient with suspected spinal cord injury is: NCLEX

**a.** Ambulate early
**b.** Immobilize the spine while maintaining airway (jaw thrust)
**c.** Remove collar
**d.** Head tilt-chin lift

# Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

| | | | |
|---|---|---|---|
| **Tension pneumothorax** | Tracheal shift, absent breath sounds, hypotension → needle decompression. | **Hemothorax** | Blood in pleura; dull percussion; chest tube. |
| **Flail chest** | Paradoxical movement. | **Tamponade** | Beck's triad; pericardiocentesis. |
| **Chest tube** | Continuous bubbling = air leak; tidaling normal. | **Splenic injury** | Kehr's sign; hypovolemia. |
| **Liver injury** | RUQ pain, shock. | **Spinal injury** | Immobilize; jaw thrust. |
| **Head injury** | Monitor GCS, ICP signs. | **Fracture** | Immobilize; check 5 Ps. |
| **Rib fracture** | Pain control, deep breathing. | **Lethal six** | Airway obstruction, tension PTX, open PTX, massive hemothorax, flail chest, tamponade. |

# Answer Key

الإجابات

**1. c** — Place the end of the chest tube in a container of normal sterile saline

**2. c** — Tracheal deviation

**3. a** — Medicate the client and have the client take deep breaths

**4. c** — Needle thoracentesis in the 2nd intercostal space

**5. b** — Unequal lung expansion and dyspnea

**6. d** — Tension pneumothorax

**7. c** — Paradoxical chest movement

**8. d** — Contralateral side in hemothorax. - A patient is admitted with a left mid-axillary chest wound and experiencing extreme dyspnea, tachycardia, and hypoxia. On assessment, there was unequal rise and fall of the chest with absent breath sounds on the left side. Also, a "sucking" sound was noticed when the patient inhales and exhales. The patient's chest x-ray showed a pneumothorax. Questions from 1-2are related to the above situation

**9. a** — Open

**10. c** — Paradoxical chest movement

**11. c** — Sharp pleuritic pain that worsens on inspiration

**12. d** — Absence of breath sounds on auscultation

**13. a** — Pneumothorax

**14. c** — Supraorbital pressure

**15. d** — Black

**16. h** — 3, 4, 1, 2

**17. d** — Has abrasions on the whole left posterior leg

**18. c** — Red

**19. g** — 2,1,3,4

**N1. b** — Tension pneumothorax
*Emergency needle decompression.*

**N2. a** — Flail chest
*A free segment moves inward on inspiration.*

**N3. a, b, c** — Hypotension; Distended neck veins; Muffled heart sounds
*Beck's triad.*

**N4. b** — Splenic rupture
*Diaphragmatic irritation by blood.*

**N5. b** — Air leak in the system
*Intermittent bubbling with expiration is normal for pneumothorax.*

**N6. b** — Immobilize the spine while maintaining airway (jaw thrust)
*Avoid neck extension.*