PEDIATRIC NURSING (THEORY) · COMPLETE QUESTION BANK

Pediatric Nursing — Theory All Lectures Question Bank

تمريض الأطفال (نظري) — مجمّع أسئلة المادة كلها

12	540	72	144
Lectures	Past exam questions	NCLEX-style	Flashcards

#	Lectures included in this file · المحاضرات اللي في الملف	Past Qs	NCLEX	Cards
1	Perspectives of Pediatric Nursing منظور تمريض الأطفال	60	6	12
2	Ethical Principles & Legal Issues المبادئ الأخلاقية والقضايا القانونية	57	6	12
3	Hospitalization of Children دخول الأطفال المستشفى	44	6	12
4	GIT Congenital Anomalies العيوب الخلقية في الجهاز الهضمي	51	6	12
5	High-Risk Neonate حديث الولادة عالي الخطورة	40	6	12
6	Play اللعب	37	6	12
7	Accident Prevention الوقاية من الحوادث	20	6	12
8	Breast Feeding الرضاعة الطبيعية	9	6	12
9	CNS Congenital Anomalies العيوب الخلقية في الجهاز العصبي	47	6	12
10	Rheumatic Fever الحمى الروماتيزمية	46	6	12
11	Nephrotic Syndrome & AGN المتلازمة الكلوية والتهاب الكلى الحاد	75	6	12
12	Respiratory Distress Syndrome (RDS) متلازمة الضائقة التنفسية عند حديثي الولادة	54	6	12

المصادر:

ميدترم 2020 · ميدترم 2021 · ميدترم 2022 · ميدترم 2024 (ترم أول 2023-2024) · كويز 1 – 2021 · كويز 2 – 2021 · كويز 1 – 2022 · كويز 1 – 2022 (نسخة بلاك بورد) · كويز 2 – 2022 · كويز 1 – 2023/2024 (شعبة 4) · كويز 1 إعادة – ترم ثاني 2023/2024 · كويز 1 – ترم ثاني 2025 · كويز 1 ملف (Quiz 1 and 2)

ملف (pediatric Quiz) كويز | ملف (Microsoft Forms) كويز 1 أطفال نظري | ملف (Quiz 1 and 2) كويز 2 شعبة 3

ملف (Quiz2 pediatric 2022) كويز 2 – 2022 | أسئلة سابقة | تجميعة صور ميدترمات سابقة | ملف (DOC-2022) كويز

ترقيم الأسئلة: رقم المحاضرة ثم رقم السؤال (مثلاً 4.12). أسئلة NCLEX عليها حرف N. الإجابات كلها في آخر الملف.

> اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

LECTURE 1

Perspectives of Pediatric Nursing

منظور تمريض الأطفال

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1.1 One of the following describes family-centered care: تكرر في 3 مصادر — ميدترم 2020

a. Recognizes that the family is the constant in a child's life
b. Reduces effect of cultural diversity on the family
c. Encourages family dependence on the health care system
d. Avoids families being part of the decision-making process

1.2 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر — ميدترم 2020

a. Establishing lifelong friendships
b. Providing financial assistance
c. Promoting disease prevention
d. Making decisions alone

1.3 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر — ميدترم 2021

a. Promoting disease prevention
b. Establishing life-long friendships
c. Providing financial assistance
d. Making decisions alone

1.4 Which of the following is the key component of family-centered care? تكرر في 2 مصادر — ميدترم 2022

a. Respect
b. Resources
c. Research
d. Reform

1.5 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر — ميدترم 2022

a. Promoting disease prevention
b. Providing financial assistance
c. Establishing life-long friendships
d. Making decisions alone

1.6 Which of the following responsibilities is one of the pediatric nurse's roles? — ميدترم 2024 (ترم أول 2023-2024)

a. Establishing life-long friendships
b. Providing financial assistance
c. Making decisions alone
d. Promoting disease prevention

1.7 Which of the following is NOT among the factors that affect the infant mortality rate? تكرر في 2 مصادر — كويز 1 – 2021

a. Short or long gestation
b. Birth weight of the newborn
c. Low level of father education
d. Low level of maternal education

1.8 A homeless school-age child came to the pediatric ER after a car accident; he stated that he lives in the street because his father is violent against him. This is an example of ..... morbidity. كويز 1 – 2021

a. Chronic
b. New
c. Acute
d. Chest

1.9 When a pediatric nurse teaches a mother to give a bath to her 6-month-old child, this reflects which concept of family-centered nursing care? كويز 1 – 2022

a. Enhancing
b. Respecting
c. Enabling
d. Collaborating

1.10 Which of the following problems is considered a new morbidity that may alter child health? كويز 1 – 2022

a. Obesity
b. Poverty
c. Isolation
d. Depression

1.11 Which of the following is most descriptive of family-centered care? كويز 1 – 2022 (نسخة بلاك بورد)
تكرر في 3 مصادر

a. Recognizes that the family is the constant in a child's life
b. Reduces effect of cultural diversity on the family
c. Avoids families being part of the decision-making process
d. Encourages family dependence on the health care system

1.12 One of the following responsibilities is included in the pediatric nurse's roles: كويز 1 – 2022 (نسخة بلاك بورد)
تكرر في 6 مصادر

a. Establishing life-long friendships
b. Making decisions alone
c. Promoting disease prevention
d. Providing financial assistance

1.13 The last cause of infant mortality in both developed and developing countries is ..... كويز 1 – 2023/2024 (شعبة 4)
تكرر في 2 مصادر

a. Accidents
b. Congenital anomalies
c. Diarrhea
d. Malnutrition

1.14 A nurse asks a mother to bathe her infant then teaches her the accurate bathing; this is an application of the concept of ..... in family-centered nursing care. كويز 1 – 2023/2024 (شعبة 4)

a. Empowerment
b. Enabling
c. Respecting
d. Collaboration

1.15 The second leading cause of infant mortality in developing countries is: كويز 1 إعادة – ترم تاني 2023/2024

a. Diarrheal disease
b. Chronic illness
c. Endocrine disorder
d. Blood disorder

1.16 One of the following is a key component of family-centered care: كويز 1 إعادة – ترم ثاني 2023/2024

a. Collaboration
b. Leadership
c. Professionalism
d. Decision making

1.17 Which of the following is a primary level of the pediatric nurse role? كويز 1 – ترم ثاني 2025

a. Provides nursing care for the ill child
b. Performs complex surgical procedures for a child
c. Assists children with disabilities following illness
d. Maintains the child's health and prevents diseases

1.18 Which of the following is a leading cause of infant mortality in developing countries? كويز 1 – ترم ثاني 2025

a. Low birth weight
b. Sepsis of newborn
c. Acute respiratory infections
d. Sudden infant death syndrome

1.19 A pediatric nurse makes an opportunity for a mother of a 6-month-old child in the pediatric ward to give a bath to her infant. This reflects which concept of family-centered nursing care? تكرر في 3 مصادر كويز 1 (ملف Quiz 1 and 2)

a. Enabling
b. Collaboration
c. Empowerment
d. Respect

1.20 Which of the following responsibilities is considered one of the pediatric nurse's roles? تكرر في 2 مصادر كويز 1 (ملف Quiz 1 and 2)

a. Establishing life-long friendships
b. Making decisions alone
c. Providing financial assistance
d. Promoting disease prevention

1.21 Which of the following is most descriptive of family-centered care? كويز 1 أطفال نظري (Microsoft Forms)
تكرر في 3 مصادر

a. Encourages family dependence on the health care system
b. Avoids families being part of the decision-making process
c. Recognizes that the family is the constant in a child's life
d. Reduces effect of cultural diversity on the family

1.22 Which of the following is the common cause of mortality among children in developing countries? كويز 1 أطفال نظري (Microsoft Forms)

a. Large birth weight
b. Female gender
c. Chronic illness
d. Acute lower respiratory infections

1.23 The most important factor affecting the infant mortality rate is ..... كويز (ملف pediatric Quiz)
تكرر في 2 مصادر

a. Birth weight
b. Gender
c. Maternal age
d. Prematurity

1.24 Which of the following represents "new morbidity" that may alter children's health? تكرر في 2 مصادر

كويز (ملف pediatric Quiz)

a. Psychological problems
b. Social problems
c. Physical problems
d. Mental problems

1.25 The most leading cause of newborn mortality in developed countries is ..... تكرر في 2 مصادر

كويز (ملف pediatric Quiz)

a. Infections
b. Accidents
c. Hypothyroidism
d. Jaundice

1.26 A pediatric nurse makes an opportunity for a mother of a 6-month-old child in the pediatric ward to give a bath to her infant. This reflects which concept of family-centered nursing care? تكرر في 3 مصادر

كويز (ملف pediatric Quiz)

a. Empowerment
b. Respect
c. Enabling
d. Collaboration

1.27 Factors that affect infant mortality:

كويز (ملف DOC-2022)

a. Place of house
b. Birth weight
c. Parent health
d. Family size

1.28 One of the key components of family-centered care is: تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. Belonging
b. Respect
c. Intimacy
d. Independency

1.29 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر

تجميعة صور ميدترمات سابقة

a. Promoting disease prevention
b. Establishing lifelong friendships
c. Making decisions alone
d. Providing financial assistance

1.30 Which of the following is the leading cause of death for children in developing countries? تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Acute lower respiratory tract infections
b. Childhood cancer
c. Sudden infant death syndrome
d. Heart diseases

1.31 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر

تجميعة صور ميدترمات سابقة

a. Providing financial assistance
b. Promoting disease prevention
c. Making decisions alone
d. Establishing lifelong friendships

1.32 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر

تجميعة صور ميدترمات سابقة

a. Establishing lifelong friendships
b. Providing financial assistance
c. Promoting disease prevention
d. Making decisions alone

1.33 Accidents are the ..... leading cause of infant mortality in both developing and developed countries. تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Last
b. Second
c. Third
d. First

1.34 One of the following describes family-centered care: تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Recognizes that the family is the constant in a child's life
b. Reduces effect of cultural diversity on the family
c. Encourages family dependence on health care systems
d. Avoids families being part of the decision-making process

1.35 A pediatric nurse makes an opportunity for a mother of a 6-month hospitalized child to give a bath to her infant. Which concept of family-centered nursing care does this reflect?

تجميعة صور ميدترمات سابقة

a. Enabling
b. Collaboration
c. Respect
d. Empowerment

1.36 Factors that affect the infant mortality:

أسئلة سابقة

a. Place of house
b. Birth weight
c. Parent health
d. Family size

1.37 The most important factor affecting the infant mortality rate is ..... تكرر في 2 مصادر

أسئلة سابقة

a. Birth weight
b. Gender
c. Maternal age
d. Prematurity

1.38 Which of the following represents "new morbidity" that may alter children' health ?

أسئلة سابقة

a. Psychological problems
b. Social problems
c. Physical problems
d. Mental problems

1.39 The most leading cause of newborn mortality in developed countries is ..... تكرر في 2 مصادر

أسئلة سابقة

a. Infections
b. Accidents
c. Hypothyroidism
d. Jaundice

1.40 A pediatric nurse makes an opportunity for a mother of 6month old child in pediatric ward to give a bath to her infant. This reflects which concepts of family centered nursing care ?

أسئلة سابقة

a. Empowerment
b. Respect
c. Enabling
d. Collaboration

1.41 Which of the following is the key component of family centered care ? تكرر في 2 مصادر

أسئلة سابقة

a. Respect.
b. Resources
c. Research.
d. Reform

1.42 One of the following responsibilities is included in the pediatric nurse's roles: تكرر في 6 مصادر

أسئلة سابقة

a. Promoting disease prevention.
b. Providing financial assistance.
c. Establishing life-long friendships.
d. Making decision alone.

1.43 Which of the following is not among the factors that affect the infant mortality rate ? تكرر في 2 مصادر

أسئلة سابقة

a. Short or long gestation
b. Birth weight of the newborn.
c. Low level of father education.
d. Low level of maternal education

1.44 A homeless school age child came to pediatric emergency room after car accident; in this stated that he lives in the street because his father is violent against him. This is an example of ..... morbidity.

أسئلة سابقة

a. Chronic.
b. New
c. Acute.
d. Chest.

1.45 Which of the following is the leading cause of death for children in developing countries ? تكرر في 2 مصادر

أسئلة سابقة

a. Acute lower respiratory tract infections
b. Childhood cancer
c. Sudden infant death syndrome
d. Heart diseases

1.46 Accidents are the ..... leading cause of infant mortality in both developing and developed countries. تكرر في 2 مصادر

أسئلة سابقة

a. Last
b. Second
c. Third
d. First

1.47 One of the following describes family centered care: تكرر في 3 مصادر

أسئلة سابقة

a. Recognizes that the family is the constant in a child's life
b. Reduces effect of cultural diversity on the family.
c. Encourages family dependence on health care systems
d. Avoids families to be part of the decision-making process

1.48 Factors that affects the infant mortality :

أسئلة سابقة

a. Place of house.
b. Birth weight.
c. Parent health.
d. Family size.

1.49 One of the key components of family centered care is : تكرر في 2 مصادر

أسئلة سابقة

a. Belonging.
b. Respect
c. Intimacy
d. Independency.

1.50 The last cause of infant mortality in both developed and developing countries is ..... تكرر في 2 مصادر

أسئلة سابقة

a. Accidents
b. Congenital anomalies
c. Diarrhea
d. Malnutrition

1.51 A nurse asks a mother to bath her infant then teach her the accurate bathing; this is an application of the concept of in family centered nursing care

أسئلة سابقة

a. Empowerment
b. Enabling
c. Respecting
d. Collaboration

1.52 A pediatric nurse makes an opportunity for a mother of a 6-month-old child in the pediatric ward to give a bath to her infant. This reflects which concept of family-centered nursing care? تكرر في 3 مصادر

أسئلة سابقة

a. Empowerment
b. Respect
c. Enabling
d. Collaboration

1.53 Which of the following is NOT among the factors that affect the infant mortality rate? تكرر في 2 مصادر

أسئلة سابقة

a. Short or long gestation
b. Birth weight of the newborn
c. Low level of father education
d. Low level of maternal education

1.54 Which of the following responsibilities is considered one of the pediatric nurse's roles? تكرر في 2 مصادر

أسئلة سابقة

a. Establishing life-long friendships
b. Providing financial assistance
c. Making decision alone
d. Promoting disease prevention

1.55 The most important factor affecting the infant mortality rate is: تكرر في 2 مصادر

أسئلة سابقة

a. Birth weight
b. Gender
c. Maternal age
d. Prematurity

1.56 Which of the following represents "new morbidity" that may alter children's health? تكرر في 2 مصادر

أسئلة سابقة

a. Psychological problems
b. Social problems
c. Physical problems
d. Mental problems

1.57 The most leading cause of newborn mortality in developed countries is: تكرر في 2 مصادر

أسئلة سابقة

a. Infections
b. Accidents
c. Hypothyroidism
d. Jaundice

1.58 Which of the following is most descriptive of family-centered care? تكرر في 3 مصادر أسئلة سابقة

a. Recognizes that the family is the constant in a child's life
b. Reduces effect of cultural diversity on the family
c. Avoids families being part of the decision-making process
d. Encourages family dependence on the health care system

1.59 Which of the following is the key component of family-centered care? تكرر في 2 مصادر أسئلة سابقة

a. Respect
b. Resources
c. Research
d. Reform

1.60 Which of the following is the leading cause of death for children in developing countries? تكرر في 2 مصادر أسئلة سابقة

a. Acute lower respiratory tract infections
b. Childhood cancer
c. Sudden infant death syndrome
d. Heart diseases

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

1.N1 The infant mortality rate is defined as the number of deaths: NCLEX

a. Under 5 years per 1000 population
b. In the first year of life per 1000 live births
c. In the first 28 days per 1000 live births
d. In the first week of life per 100 births

1.N2 Which is considered the most important single factor affecting infant mortality? NCLEX

a. Gender
b. Birth weight
c. Family size
d. Place of residence

1.N3 Select ALL examples of the 'new morbidity'. SELECT ALL

a. Poverty
b. Violence
c. School failure
d. Adjustment to divorce
e. Measles

1.N4 The nurse teaches a mother how to give insulin so she can do it at home. Which family-centered care concept is this? NCLEX

a. Respect
b. Enabling
c. Collaboration
d. Dignity

1.N5 Which is a PRIMARY prevention role of the pediatric nurse? NCLEX

a. Caring for a child with pneumonia
b. Immunization and health teaching
c. Rehabilitation after injury
d. Physiotherapy for cerebral palsy

1.N6 The leading cause of death in children older than 1 year is: NCLEX

a. Infections
b. Congenital anomalies
c. Injuries (accidents)
d. Cancer

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

IMR	Infant deaths in first year per 1000 live births.	**Neonatal mortality**	Deaths < 28 days of life.
Post-neonatal mortality	Deaths from 28 days to 11 months.	**Most important IMR factor**	Birth weight.
Leading infant death (developed)	Congenital anomalies/short gestation; newborn: infections (per bank).	**Leading infant death (developing)**	Acute lower respiratory infections, then diarrhea.
Accidents	Last cause of infant mortality in both developed & developing countries.	**New morbidity**	Social/behavioral problems: poverty, violence, school failure, divorce.
Family-centered care	Family is the constant in the child's life.	**FCC concepts**	Enabling, empowerment, respect, collaboration (key component: respect).
Enabling	Giving family opportunity to use/learn skills (e.g., bathing her infant).	**Nurse roles**	Health promotion, disease prevention, advocacy, therapeutic relationship, research, ethical decision making.

LECTURE 2

Ethical Principles & Legal Issues

المبادئ الأخلاقية والقضايا القانونية

Part A · Past Exam Questions

أسئلة الدفعات السابقة

2.1 Hospitalized children have the right to be admitted to hospital only when their: تكرر في 3 مصادر — ميدترم 2020

a. Diseases are not contagious
b. Socio-economic state is low
c. Conditions require in-patient care
d. Parents are tired of caring for them

2.2 The ethical principle that obligates the pediatric nurse to keep promises and high quality of nursing care to children/family is: تكرر في 3 مصادر — ميدترم 2020

a. Fidelity
b. Non-maleficence
c. Beneficence
d. Autonomy

2.3 Which of the following is considered a secondary ethical principle? تكرر في 4 مصادر — ميدترم 2020

a. Justice
b. Veracity
c. Beneficence
d. Autonomy

2.4 Informed consent before the surgical operation is an important outcome of the ethical principle of: تكرر في 4 مصادر — ميدترم 2021

a. Non-maleficence
b. Autonomy
c. Beneficence
d. Justice

2.5 The pediatric hospitals must offer suitable accommodations for the parents because: تكرر في 2 مصادر — ميدترم 2022

a. It is a right of hospitalized children
b. Parents like to spend time beside them
c. Rooms are plenty in the hospitals
d. Parents pay for the accommodations

2.6 Informed consent before the surgical operation is an important outcome of the ethical principle of: تكرر في 4 مصادر — ميدترم 2022

a. Justice
b. Autonomy
c. Non-maleficence
d. Beneficence

2.7 Informed consent is an important outcome of the ethical principle of: — ميدترم 2024 (ترم أول 2023-2024)

a. Beneficence
b. Autonomy
c. Justice
d. Non-maleficence

2.8 A pediatric nurse told the mother that her child will become as normal children of his age — not the truth, but because she felt sorry for the mother. This action violates the ethical principle of: تكرر في 2 مصادر

كويز 1 – 2021

a. Justice
b. Veracity
c. Beneficence
d. Fidelity

2.9 When a pediatric nurse discusses the child's information with her friend nurse who works in another hospital, this is a violation of the ethical principle of: تكرر في 2 مصادر

كويز 1 – 2021

a. Confidentiality
b. Sanctity of life
c. Beneficence
d. Justice

2.10 The pediatric hospitals must make suitable accommodations for the parents because ..... تكرر في 2 مصادر

كويز 1 – 2021

a. Parents pay for the accommodations
b. It is a right of hospitalized children
c. Rooms are plenty in the hospitals
d. It is normal Egyptian hospitality

2.11 When the nurse includes the family and the child in choosing the appropriate treatment for the child's condition, this reflects:

كويز 1 – 2022

a. Restorative and primitive
b. Researcher and developer
c. Therapeutic relationship
d. Ethical decision making

2.12 When a child refuses to take his medication, this is an example of which ethical principle?

كويز 1 – 2022

a. Justice
b. Autonomy
c. Confidentiality
d. Beneficence

2.13 Non-maleficence is a crucial basic ethical principle in caring for children, which means: تكرر في 2 مصادر

كويز 1 – 2022 (نسخة بلاك بورد)

a. Do no harm
b. Decide for self
c. Telling the truth
d. Fairness in care

2.14 When a preschooler asks the nurse before cannula insertion if it hurts and the nurse replies "no dear, it won't", this is a violation of the ethical principle of:

كويز 1 – 2023/2024 (شعبة 4)

a. Fidelity
b. Veracity
c. Beneficence
d. Justice

2.15 Which of the following reflects one of the hospitalized child's rights?

كويز 1 – 2023/2024 (شعبة 4)

a. Withhold information from the child's mother
b. Ask the mother to change her clothes into white
c. Discuss the child's condition with her friends
d. Accommodate the mother to stay with the child

2.16 Supporting an adolescent to make decisions about his/her health reflects which ethical principle? كويز 1 إعادة – ترم تاني 2023/2024

a. Confidentiality
b. Justice
c. Beneficence
d. Autonomy

2.17 Which ethical principle is MOST directly relevant to ensuring fair distribution of care for all children? كويز 1 – ترم تاني 2025

a. Beneficence
b. Non-maleficence
c. Justice
d. Fidelity

2.18 Which of the following is an example of non-maleficence in pediatric care? كويز 1 – ترم تاني 2025

a. Administering the least invasive treatment to avoid unnecessary harm
b. Allowing a child to make their own medical decisions without parental consent
c. Prescribing medication without considering possible side effects
d. Ignoring a child's discomfort during a procedure to save time

2.19 The ethical principle that encourages the nurse to manage the child's pain is ..... تكرر في 2 مصادر كويز 1 (ملف Quiz 1 and 2)

a. Autonomy
b. Veracity
c. Fidelity
d. Beneficence

2.20 Informed consent before the surgical operation is an important outcome of the ethical principle of: تكرر في 4 مصادر كويز 1 (ملف Quiz 1 and 2)

a. Justice
b. Beneficence
c. Non-maleficence
d. Autonomy

2.21 A pediatric nurse, to minimize harm (both physical and emotional), applies a topical anesthetic cream to numb the injection site before administering the shot. This reflects the principle of ..... كويز 1 أطفال نظري (Microsoft Forms)

a. Fidelity
b. Autonomy
c. Non-maleficence
d. Veracity

2.22 In pediatric nursing, the ethical principle of justice is most accurately demonstrated in which situation? كويز 1 أطفال نظري (Microsoft Forms)

a. Giving better treatment options to children whose parents can afford
b. Allocating equal time to each child during rounds, regardless of their needs
c. Providing extra attention to a child from a wealthy family
d. Prioritizing care for a critically ill child based on their need

2.23 When the nurse does the best interest of the child rather than her own self-interest, this ascertains the ethical principle of ..... تكرر في 2 مصادر كويز (ملف pediatric Quiz)

a. Veracity
b. Beneficence
c. Fidelity
d. Justice

2.24 Justice is a crucial basic ethical principle in caring for children, which means: كويز (ملف DOC-2022)
تكرر في 2 مصادر

a. The right to make one's own decisions
b. Often referred to as fairness
c. The duty to do no harm
d. Refers to telling the truth

2.25 Which of the following is considered a secondary ethical principle? كويز (ملف DOC-2022)
تكرر في 4 مصادر

a. Justice
b. Veracity
c. Autonomy
d. Beneficence

2.26 Hospitalized children have the right to be admitted to hospital only when their: تجميعة صور ميدترمات سابقة
تكرر في 3 مصادر

a. Diseases are not contagious
b. Conditions require in-patient care
c. Socio-economic state is low
d. Parents are tired of caring for them

2.27 The ethical principle that obligates the pediatric nurse to keep promises and high quality of nursing care to children/family is: تكرر في 3 مصادر تجميعة صور ميدترمات سابقة

a. Non-maleficence
b. Autonomy
c. Fidelity
d. Beneficence

2.28 Which of the following is considered a secondary ethical principle? تجميعة صور ميدترمات سابقة
تكرر في 4 مصادر

a. Veracity
b. Autonomy
c. Justice
d. Beneficence

2.29 Hospitalized children have the right to be admitted to hospital only when their: تجميعة صور ميدترمات سابقة
تكرر في 3 مصادر

a. Diseases are not contagious
b. Parents are tired of caring for them
c. Socio-economic state is low
d. Conditions require in-patient care

2.30 The ethical principle that obligates the pediatric nurse to keep promises and high quality of nursing care to children/family is: تكرر في 3 مصادر تجميعة صور ميدترمات سابقة

a. Fidelity
b. Non-maleficence
c. Beneficence
d. Autonomy

2.31 A pediatric nurse promised her patient that she will come in the morning with a toy and she forgot to bring it to him. This is the ethical principle of ..... تجميعة صور ميدترمات سابقة
تكرر في 2 مصادر

a. Fidelity
b. Justice
c. Veracity
d. Beneficence

2.32 Hospitalized children have the right to be admitted to hospital only when their: تجميعة صور ميدترمات سابقة

تكرر في 3 مصادر

a. Diseases are not contagious
b. Socio-economic state is low
c. Conditions require in-patient care
d. Parents are tired of caring for them

2.33 Which of the following is considered a secondary ethical principle? تجميعة صور ميدترمات سابقة

تكرر في 4 مصادر

a. Justice
b. Veracity
c. Beneficence
d. Autonomy

2.34 When the nurse do the best interest of the child rather than the self-interest of the herself, this ascertain the ethical principle of..... أسئلة سابقة

a. Veracity
b. Beneficence
c. Fidelity
d. Justice

2.35 The pediatric hospitals must offer suitable accommodations for the parents because: أسئلة سابقة

تكرر في 2 مصادر

a. It is a right of hospitalized children
b. Parents likes to spend time side them.
c. Rooms are plenty in the hospitals.
d. Parents pay for the accommodations.

2.36 Informed consent before the surgical operation is an important outcome of ethical principle of: أسئلة سابقة

a. Justice.
b. Autonomy.
c. Non-maleficence.
d. Beneficence.

2.37 A pediatric nurse told the mother that her child will become as normal children of his age, this is no truth but because she felt sorry for the mother. This action violates the ethical principle of: تكرر في 2 مصادر أسئلة سابقة

a. Justice
b. Veracity
c. Beneficence
d. Fidelity

2.38 When a pediatric nurse discusses the child information with her friend nurse who works in a hospital, this is a violation of ethical principle of: أسئلة سابقة

a. Confidentiality
b. Sanctity of life
c. Beneficence
d. Justice

2.39 Hospitalized children have the right to be admitted to hospital only when their: أسئلة سابقة

تكرر في 3 مصادر

a. Diseases are not contagious
b. Conditions require in-patient care
c. Socio-economic state is low
d. Parents are tiered of caring for them

2.40 The ethical principle that obligates the pediatric nurse to keep promises and high quality of nursing care to children / family is: تكرر في 3 مصادر

أسئلة سابقة

a. Non-maleficence
b. Autonomy
c. Fidelity
d. Beneficence

2.41 Which of the following considered as secondary ethical principle ?

أسئلة سابقة

a. Veracity
b. Autonomy
c. Justice
d. Beneficence

2.42 A pediatric nurse promised her patient that she will come in the morning with a toy and she forgot to bring it to him. This is the ethical principle of..... تكرر في 2 مصادر

أسئلة سابقة

a. Fidelity
b. Justice
c. Veracity
d. Beneficence

2.43 Non-maleficence is a crucial basic ethical principle in caring for children, which means: تكرر في 2 مصادر

أسئلة سابقة

a. Do no harm
b. Decide for self.
c. Telling the truth.
d. Fairness in care.

2.44 Justice is a crucial basic ethical principle in caring for children which means: تكرر في 2 مصادر

أسئلة سابقة

a. The right to make one's own decisions.
b. Often referred to as fairness.
c. Is the duty to do no harm.
d. Refers to telling the truth.

2.45 The ethical principle that encourages the nurse to manage the child's pain is ..... تكرر في 2 مصادر

أسئلة سابقة

a. Autonomy.
b. Veracity.
c. Fidelity.
d. Beneficence.

2.46 When a preschooler child asks the nurse before cannula insertion: if it hurts and the nurse reply: no dear it won't. This is violation of the ethical principle of:

أسئلة سابقة

a. Fidelity
b. Veracity
c. Beneficence
d. Justice

2.47 Which of the following reflects one of hospitalized child's rights ?

أسئلة سابقة

a. Withhold information from the child's mother
b. Ask mother to change her clothes into white
c. Discuss child condition with her friends
d. Accommodate the mother to stay with child

2.48 The ethical principle that encourages the nurse to manage the child's pain:

أسئلة سابقة

a. Veracity
b. Beneficence
c. Fidelity
d. Autonomy

2.49 Informed consent before the surgical operation is an important outcome of the ethical principle of: تكرر في 4 مصادر

أسئلة سابقة

a. Autonomy
b. Beneficence
c. Non-maleficence
d. Justice

2.50 A pediatric nurse told the mother that her child will become as normal children of his age — not true, but because she felt sorry for the mother. This action violates the ethical principle of: تكرر في 2 مصادر

أسئلة سابقة

a. Justice
b. Veracity
c. Beneficence
d. Fidelity

2.51 When a pediatric nurse discusses the child's information with her friend nurse who works in a hospital, this is a violation of the ethical principle of: تكرر في 2 مصادر

أسئلة سابقة

a. Confidentiality
b. Sanctity of life
c. Beneficence
d. Justice

2.52 The pediatric hospitals must make suitable accommodations for the parents because: تكرر في 2 مصادر

أسئلة سابقة

a. Parents pay for the accommodations
b. It is a right of hospitalized children
c. Rooms are plenty in the hospitals
d. It is normal Egyptian hospitality

2.53 When the nurse does the best interest of the child rather than her own self-interest, this ascertains the ethical principle of: تكرر في 2 مصادر

أسئلة سابقة

a. Veracity
b. Beneficence
c. Fidelity
d. Justice

2.54 The ethical principle that obligates the pediatric nurse to keep promises and high quality of nursing care to children/family is: تكرر في 3 مصادر

أسئلة سابقة

a. Fidelity
b. Non-maleficence
c. Beneficence
d. Autonomy

2.55 Which of the following is considered a secondary ethical principle? تكرر في 4 مصادر

أسئلة سابقة

a. Justice
b. Veracity
c. Beneficence
d. Autonomy

2.56 Non-maleficence is a crucial basic ethical principle in caring for children, which means: تكرر في 2 مصادر

أسئلة سابقة

a. Do no harm
b. Decide for self
c. Telling the truth
d. Fairness in care

2.57 A pediatric nurse promised her patient that she will come in the morning with a toy and she forgot to bring it. This is the ethical principle of: تكرر في 2 مصادر

أسئلة سابقة

a. Fidelity
b. Justice
c. Veracity
d. Beneficence

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

2.N1 A nurse tells a school-age child honestly that the injection will hurt for a moment. Which principle is applied? NCLEX

a. Fidelity
b. Veracity
c. Justice
d. Autonomy

2.N2 An adolescent is involved in decisions about his treatment plan. This respects: NCLEX

a. Beneficence
b. Autonomy
c. Non-maleficence
d. Fidelity

2.N3 Select ALL that are basic (primary) ethical principles. SELECT ALL

a. Autonomy
b. Beneficence
c. Non-maleficence
d. Justice
e. Veracity
f. Confidentiality

2.N4 The nurse uses EMLA cream before venipuncture to minimize harm. This reflects: NCLEX

a. Non-maleficence
b. Justice
c. Fidelity
d. Veracity

2.N5 Which action violates confidentiality? NCLEX

a. Reporting suspected child abuse to authorities
b. Discussing a child's diagnosis with a nurse friend from another hospital
c. Sharing information with the care team
d. Documenting in the medical record

2.N6 Justice in pediatric nursing is best demonstrated by: NCLEX

a. Giving more time to families who pay more
b. Distributing care fairly according to the child's need
c. Letting the child decide everything
d. Keeping promises

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Autonomy	Right of self-determination; basis of informed consent.	**Beneficence**	Doing good — acting in the child's best interest (e.g., pain management).
Non-maleficence	Do no harm (e.g., minimize pain of procedures).	**Justice**	Fairness — care distributed according to need.
Veracity	Telling the truth (secondary principle).	**Fidelity**	Keeping promises and commitments; high-quality care.

Confidentiality	Not sharing the child's information with people not involved in care.
Primary principles	Autonomy, beneficence, non-maleficence, justice.
Secondary principles	Veracity, fidelity, confidentiality.
Informed consent for minors	Given by parent/guardian; child gives assent according to age.
Hospitalized child right	Admission only when care cannot be provided at home; parents stay with child.
Ethical decision making	Involving the child and family in choosing treatment.

LECTURE 3

Hospitalization of Children

دخول الأطفال المستشفى

Part A · Past Exam Questions

أسئلة الدفعات السابقة

3.1 A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر — ميدترم 2020

a. Despair
b. Detachment
c. Protest
d. Regression

3.2 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر — ميدترم 2020

a. Concerned with appearance
b. Separation from peer group
c. Stranger anxiety
d. Fear of pain

3.3 A hospitalized toddler, for two days, cries because he was left alone. The nurse realizes this behavior is typical of: — ميدترم 2021

a. Regression
b. Despair
c. Protest
d. Stranger anxiety

3.4 Which of the following represents the major stressor of hospitalization for adolescents? تكرر في 2 مصادر — ميدترم 2022

a. Separation anxiety
b. Fear of pain
c. Concerned with appearance
d. Loss of control

3.5 Separation anxiety is more drastic among children aged: تكرر في 2 مصادر — ميدترم 2024 (ترم أول 2023-2024)

a. 2 months to 6 months
b. 6 months to 12 months
c. 6 months to 6 years
d. 6 months to 12 years

3.6 The pediatric nurse should realize that guided imagery is successfully used to decrease pain of: — ميدترم 2024 (ترم أول 2023-2024)

a. Toddlers
b. Preschoolers
c. School-agers
d. Adolescents

3.7 A child hospitalized for a chronic illness showed symptoms of depression but later started interacting with others. What does the nurse infer? The child is ..... — كويز 1 – 2021

a. Detached from both parents
b. Getting used to the surroundings
c. Showing improved social skills
d. Content with the care provided

3.8 Which of the following stressors would the nurse recognize as having the greatest impact on the school-age hospitalized child? تكرر في 2 مصادر

كويز 1 – 2022

a. Separation from family
b. Fear of death
c. Loss of control and privacy
d. Concerned with appearance

3.9 Which of the following child's behaviors leads the nurse to believe that the child is exhibiting the stage of protest?

كويز 1 – 2022

a. Screams and hits the nurse
b. Withdrawn from others
c. Returns to the habit of bed-wetting
d. Sits in a corner with a toy

3.10 Which of the following represents the major stressor of hospitalization for children from middle infancy throughout the preschool years? تكرر في 2 مصادر

كويز 1 – 2022 (نسخة بلاك بورد)

a. Separation anxiety
b. Fear of bodily injury
c. Fear of pain
d. Loss of control

3.11 Which of the following should the nurse do to prevent separation anxiety in a hospitalized toddler? تكرر في 2 مصادر

كويز 1 – 2022 (نسخة بلاك بورد)

a. Assume the parental role when parents are not present
b. Limit visits by siblings and relatives
c. Establish a routine similar to the child's home
d. Rotate nursing staff so the child becomes comfortable

3.12 Attendance of parents during painful procedures on their toddlers should be:

كويز 1 – 2023/2024 (شعبة 4)

a. Based on individual assessment of parents
b. Based on the type of procedure to be performed
c. Discouraged for the benefit of parents and child
d. Encouraged and permitted freely

3.13 Which of the following substances increases pain sensation?

كويز 1 – 2023/2024 (شعبة 4)

a. Histamine
b. Endorphins
c. Enkephalins
d. Dynorphins

3.14 Regression is a reaction to hospitalization; ..... are at high risk for it.

كويز 1 – 2023/2024 (شعبة 4)

a. Infants
b. Toddlers
c. Preschoolers
d. School-agers

3.15 Which one of the following stressors most affects a 2-year-old toddler?

كويز 1 – 2023/2024 (شعبة 4)

a. Separation from peers
b. Fear of the dark
c. Separation from parents
d. Fear of death

3.16 The nurse best describes intermittent pain as pain that:

كويز 1 إعادة – ترم تاني 2023/2024

a. Stops and starts again
b. Moves from one area to another
c. Covers a large area
d. Is sticking and intense

3.17 Which of the following stressors would the nurse recognize as having the greatest impact on the school-age hospitalized child? تكرر في 2 مصادر

كويز 1 إعادة – ترم تاني 2023/2024

a. Fear of death
b. Loss of control and privacy
c. Concerned with appearance
d. Separation from family

3.18 In the despair phase of separation anxiety the child behaves as follows:

كويز 1 إعادة – ترم تاني 2023/2024

a. Shows signs of interest in the surroundings
b. Avoids and rejects contact with strangers
c. Loud cry, scream and search for parent with eyes
d. Inactive, withdrawn, depressed and sad

3.19 Which of the following child's behaviors exhibits the stage of protest in separation anxiety?

كويز 1 – ترم تاني 2025

a. Cries, screams and clings to parents
b. Withdrawn from others, sad and depressed
c. Interested in surroundings, active and communicative
d. Interacts with strangers and any other children

3.20 Which of the following is a primary stressor for a child during hospitalization?

كويز 1 – ترم تاني 2025

a. Boredom from lack of activities
b. Unfamiliar hospital food
c. The noise of medical equipment
d. Separation from parents and siblings

3.21 A behavior observed in the despair phase of separation anxiety as a hospitalization reaction:

كويز 1 أطفال نظري (Microsoft Forms)

a. Forms new but superficial relationships and appears happy
b. Regresses to earlier behavior (e.g. thumb sucking, bed wetting)
c. Shows increased interest in surroundings, interacts with strangers
d. Physically attacks strangers (kicks, bites, hits, pinches)

3.22 Which of the following is the period in which hospitalization reactions are obvious and risky?

كويز 1 أطفال نظري (Microsoft Forms)

a. 6 months to 18 months
b. 3 months to 3 years
c. 6 months to 6 years
d. 3 months to 12 months

3.23 A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Detachment
b. Regression
c. Protest
d. Despair

3.24 A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Despair
b. Detachment
c. Protest
d. Regression

3.25 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Concerned with appearance
b. Separation from peer group
c. Stranger anxiety
d. Fear of pain

3.26 The characteristics of the despair phase of separation anxiety are:

تجميعة صور ميدترمات سابقة

a. Rejects contact with strangers
b. The child shows signs of interest in the surroundings
c. Loud cry, scream and searches for parent with eyes
d. Depressed, regressive behavior, withdrawal

3.27 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Fear of pain
b. Concerned with appearance
c. Stranger anxiety
d. Separation from peer group

3.28 What is the appropriate nursing intervention to minimize separation anxiety in a hospitalized child?

تجميعة صور ميدترمات سابقة

a. Encourage contact with children the same age
b. Provide for privacy
c. Allow mother's rooming-in
d. Explain procedures and routines

3.29 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Stranger anxiety
b. Separation from peer group
c. Concerned with appearance
d. Fear of pain

3.30 A hospitalized toddler newly admitted and screams because he was left alone. The nurse realizes this behavior as typical of:

أسئلة سابقة

a. Despair
b. Detachment
c. Protest
d. Regression

3.31 What is the appropriate nursing intervention minimize separation anxiety in a hospitalized child ?

أسئلة سابقة

a. Encourage contact with children the same age
b. Provide for privacy.
c. Allow mother's rooming in.
d. Explain procedures and routines.

3.32 The pediatric hospitals must offer suitable accommodations for the parents because:

أسئلة سابقة

a. It is a right of hospitalized children
b. Parents likes to spend time side them
c. Rooms are plenty in the hospitals.
d. Parents pay for the accommodations.

3.33 Which of the following represents the major stressor of hospitalization for adolescents ? تكرر في 2 مصادر

أسئلة سابقة

a. Separation anxiety.
b. Fear of pain.
c. Concerned with appearance.
d. Loss of control.

3.34 Separation anxiety is more drastic among children aged..... تكرر في 2 مصادر

أسئلة سابقة

a. 2 months to 6 months
b. 6 months to 12 months
c. 6 months to 6 years
d. 6 months to 12 years

3.35 Which of the following is considered more stressful for hospitalized school age children? تكرر في 3 مصادر

أسئلة سابقة

a. Stranger anxiety
b. Loss of control and privacy
c. Considered with appearance
d. Fear of pain .

3.36 Which of the following represents the major stressor of hospitalization for children from middle infancy throughout the preschool years? تكرر في 2 مصادر

أسئلة سابقة

a. Separation anxiety
b. Fear of bodily injury
c. Fear of pain
d. Loss of control

3.37 Regression is a reaction to hospitalization that is ..... are at high risk for it.

أسئلة سابقة

a. Infants
b. Toddlers
c. Preschoolers
d. School agers

3.38 Which one of the following stressors most affect a 2-year-old toddler child?

أسئلة سابقة

a. Separation from peers
b. Fear of the dark
c. Separation from parents
d. Fear of death

3.39 Which of the following reflects one of hospitalized child's rights?

أسئلة سابقة

a. Withhold information from the child's mother
b. Ask mother to change her clothes into white
c. Discuss child condition with her friends
d. Accommodate the mother to stay with child

3.40 A hospitalized toddler, for two days, cries because he left alone. The nurse realizes this behavior is typical of:

أسئلة سابقة

a. Regression.
b. Despair.
c. Protest.
d. Stranger anxiety.

3.41 Which of the following should the nurse do to prevent separation anxiety in a hospitalized toddler? تكرر في 2 مصادر

أسئلة سابقة

a. Assume the parental role when parents are not present.
b. Limit visits by siblings and relatives.
c. Establish a routine similar to child's home.
d. Rotate nursing staff so the child becomes comfortable

3.42 Hospitalized children have the right to be admitted to hospital only when their: أسئلة سابقة

a. Diseases are not contagious
b. Socio-economic state is low
c. Conditions require in-patient care
d. Parents are tiered of caring for them

3.43 Which of the following is considered more stressful for hospitalized school age children? أسئلة سابقة
تكرر في 3 مصادر

a. Fear of pain
b. Considered with appearance
c. Stranger anxiety
d. Separation from peer group

3.44 A child is hospitalized for a chronic illness. Initially, the child showed symptoms of depression but later started interacting with others. What does the nurse infer from the patient's behavior? The child is: أسئلة سابقة

a. Content with the care provided.
b. Showing improved social skills.
c. Getting used to the surroundings.
d. Detached from both parents.

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

3.N1 A 2-year-old cries loudly, clings to the mother and refuses the nurse. This is the stage of: NCLEX

a. Despair
b. Detachment
c. Protest
d. Denial

3.N2 Which is the major stressor for a hospitalized adolescent? NCLEX

a. Separation from parents
b. Altered body image and loss of peer contact
c. Fear of the dark
d. Stranger anxiety

3.N3 Select ALL interventions that reduce separation anxiety in a hospitalized toddler. SELECT ALL

a. Encourage rooming-in of the parent
b. Keep the home routine (bedtime rituals)
c. Bring a favorite toy or blanket from home
d. Rotate different nurses every shift
e. Limit parental visits

3.N4 A child who previously cried now appears happy and interacts superficially with everyone, ignoring the parent on visits. This is: NCLEX

a. Protest
b. Despair
c. Detachment (denial)
d. Regression

3.N5 Regression during hospitalization is most common among: NCLEX

a. Infants
b. Preschoolers
c. School-age children
d. Adolescents

3.N6 **Which statement best describes the effect of loss of control on a school-age child?** NCLEX

a. It has no effect

b. They fear losing control of their bodies and roles and may react with depression or hostility

c. They develop stranger anxiety

d. They return to parallel play

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Separation anxiety age	Most marked from 6 months to about 30 months (bank: 6 months–6 years).	**Protest**	Cries, screams, searches for parent, rejects strangers.
Despair	Inactive, withdrawn, sad, depressed; may regress.	**Detachment (denial)**	Superficial happiness, interacts with everyone, ignores parent — most harmful.
Toddler major stressor	Separation from parents.	**Preschooler reaction**	Regression; fear of bodily injury/mutilation.
School-age stressor	Loss of control & privacy; separation from peers.	**Adolescent stressor**	Body image/appearance; separation from peers.
Reducing separation anxiety	Rooming-in, home routines, favorite objects, consistent nurse.	**Pain substances**	Histamine, bradykinin, prostaglandins increase pain; endorphins decrease it.
Guided imagery	Pain technique effective for school-age children.	**Parents during painful procedures**	Encouraged to be present (supportive role).

LECTURE 4

GIT Congenital Anomalies

العيوب الخلقية في الجهاز الهضمي

Part A · Past Exam Questions

أسئلة الدفعات السابقة

4.1 Which of the following is the nurse's priority in the plan of care for an infant admitted with pyloric stenosis? تكرر في 4 مصادر — ميدترم 2020

a. Evaluate hydration status
b. Assess tissue perfusion
c. Assess respiratory status
d. Maintain thermoregulation

4.2 A newborn's failure to pass meconium within 24 hours after birth indicates which of the following? تكرر في 2 مصادر — ميدترم 2020

a. Intussusception
b. Abdominal wall defect
c. Imperforate anus
d. Aganglionic megacolon

4.3 When the nurse observes that the infant has respiratory distress and the apical pulse is auscultated on the right side, the nurse suspects the infant has: تكرر في 3 مصادر — ميدترم 2020

a. Pyloric stenosis
b. Cleft palate
c. Diaphragmatic hernia
d. Biliary atresia

4.4 For an infant undergoing surgery to repair tracheo-esophageal fistula, it is essential that nursing care is directed toward prevention of: تكرر في 2 مصادر — ميدترم 2020

a. Projectile vomiting
b. Dehydration
c. Aspiration
d. Abdominal distension

4.5 The position of choice postoperatively for a child with cleft palate is: تكرر في 2 مصادر — ميدترم 2020

a. Upright
b. On one side
c. Prone
d. Supine

4.6 A nurse is assessing a neonate with esophageal atresia. Which finding should the nurse expect to encounter? تكرر في 2 مصادر — ميدترم 2021

a. Presence of feeding reflexes
b. Increase in saliva
c. Gastric tube easily passed
d. Feeding without difficulty

4.7 A priority post-operative care for an infant with cleft lip and palate is: تكرر في 3 مصادر — ميدترم 2022

a. Protect the operative site from infection and injury
b. Provide family support and education
c. Spend good time and play with the infant
d. Provide a safe environment to prevent falling

4.8 Preoperatively, an infant with biliary atresia must be provided with ..... formula: تكرر في 2 مصادر

ميدترم 2022

a. Intravenous routine for age
b. Water-soluble vitamin containing
c. High-caloric fatty total parenteral
d. A regular routine age group

4.9 When planning care for the infant with cleft lip and palate, the priority nursing action is: تكرر في 2 مصادر

ميدترم 2024 (ترم أول 2023-2024)

a. Burp the baby frequently
b. Temporarily avoid breastfeeding
c. Put the child in supine position
d. Prevent the baby from crying

4.10 The position of choice postoperatively for a child with cleft palate is: تكرر في 2 مصادر

كويز 1 – 2021

a. Supine
b. Prone
c. Upright
d. One side

4.11 The main goal of preoperative nursing care for a newborn with esophageal atresia with tracheoesophageal fistula is: تكرر في 2 مصادر

كويز 1 – 2021

a. Promote growth
b. Maintain skin intact
c. Prevent aspiration
d. Prevent pain

4.12 Which of the following is expected to be found in a child with congenital diaphragmatic hernia?

كويز 1 – 2022

a. A palpable mass in the right upper quadrant
b. Scaphoid abdomen and diminished breath sounds
c. Weight loss and bloody diarrhea
d. Severe, crampy abdominal pain and lethargy

4.13 Which data does the nurse expect to find when assessing a child with pyloric stenosis?

كويز 1 – 2022

a. Watery diarrhea
b. Projectile vomiting
c. Increased urine output
d. Vomiting large amounts of bile

4.14 Which clinical manifestations would the nurse expect to observe while caring for the child diagnosed with intussusception? تكرر في 2 مصادر

كويز 1 – 2023/2024 (شعبة 4)

a. Currant jelly stool
b. Foul-smelling stool
c. Dark brown stool
d. Watery stool

4.15 Which clinical manifestations would the nurse expect to assess in the child diagnosed with Hirschsprung's disease? تكرر في 2 مصادر

كويز 1 – 2023/2024 (شعبة 4)

a. Bloody diarrhea, fever, and vomiting
b. Irritability, severe abdominal cramps and fecal soiling
c. Constipation and passage of ribbon-like stool
d. Respiratory distress and scaphoid abdomen

4.16 **Preoperative care for an infant with biliary atresia includes providing formula with:** كويز 1 إعادة – ترم ثاني 2023/2024

a. Artificial milk
b. Low sugar
c. Water-soluble vitamins
d. High salt

4.17 **A priority post-operative care for an infant with cleft lip and palate is to:** كويز 1 إعادة – ترم ثاني 2023/2024

a. Protect the operative site from infection or injury
b. Keep a safe environment to prevent falling
c. Provide family support and education
d. Spend good time and play with the infant

4.18 **Which of the following is expected to be found while assessing a child with congenital diaphragmatic hernia?** كويز 1 – ترم ثاني 2025

a. Severe abdominal distension
b. Diminished breathing sounds
c. Weight loss and bloody diarrhea
d. Severe, crampy abdominal pain

4.19 **Which of the following is an important preoperative nursing intervention for a child with Hirschsprung disease?** كويز 1 – ترم ثاني 2025

a. Limiting the child's fluid intake
b. Avoid giving stool softeners or enemas
c. Providing large frequent feedings
d. Measure abdominal girth periodically

4.20 **Which clinical manifestations would the nurse expect to assess in a child diagnosed with Hirschsprung's disease?** كويز 1 (ملف Quiz 1 and 2)

a. Constipation and passage of ribbon-like stool
b. Irritability, severe abdominal cramps and fecal soiling
c. Respiratory distress and scaphoid abdomen
d. Bloody diarrhea, fever, and vomiting

4.21 **Which of the following is the nurse's priority in the plan of care for an infant admitted with pyloric stenosis?** تكرر في 4 مصادر كويز 1 أطفال نظري (Microsoft Forms)

a. Assesses respiratory status
b. Evaluates hydration status
c. Assesses tissue perfusion
d. Maintains thermoregulation

4.22 **A priority post-operative care for an infant with cleft lip and palate is:** تكرر في 3 مصادر كويز 1 أطفال نظري (Microsoft Forms)

a. Provides family support and education
b. Provides a safe environment to prevent falling
c. Spends good time and plays with infants
d. Protects operative sites from infection and injury

4.23 The nurse is caring for a 4-month-old who has just had an isolated cleft lip repaired. Select the best position in the immediate post-operative period: تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. Prone
b. Supine
c. Left side-lying
d. Right side-lying

4.24 The nurse monitors a newborn with a suspected diagnosis of imperforate anus. Which is a clinical manifestation associated with this disorder? تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. The passage of currant jelly-like stools
b. Palpated mass in the abdomen
c. Failure to pass meconium immediately after birth
d. Bile-stained fecal emesis

4.25 Which of the following is the nurse's priority in the plan of care for an infant admitted with pyloric stenosis? تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Assess respiratory status
b. Maintain thermoregulation
c. Evaluate hydration status
d. Assess tissue perfusion

4.26 For an infant undergoing surgery to repair tracheo-esophageal fistula, it is essential that nursing care is directed toward prevention of: تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Projectile vomiting
b. Abdominal distension
c. Dehydration
d. Aspiration

4.27 When the nurse observes that the infant has respiratory distress and the apical pulse is auscultated on the right side, the nurse suspects the infant has: تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Pyloric stenosis
b. Biliary atresia
c. Cleft palate
d. Diaphragmatic hernia

4.28 For an infant undergoing surgery to repair tracheo-esophageal fistula, it is essential that nursing care is directed toward prevention of: تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Projectile vomiting
b. Abdominal distension
c. Dehydration
d. Aspiration

4.29 The nurse is reviewing the record of a child with a diagnosis of Hirschsprung disease. Which data should the nurse expect to note?

تجميعة صور ميدترمات سابقة

a. Ribbon-like, foul-smelling stools
b. Jaundice, pruritus, irritability
c. Increased urine output
d. Projectile vomiting

4.30 For an infant undergoing surgery to repair tracheo-esophageal fistula, it is essential that nursing care is directed toward prevention of: تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Projectile vomiting
b. Dehydration
c. Aspiration
d. Abdominal distension

4.31 Which of the following is the nurse's priority in the plan of care for an infant admitted with pyloric stenosis? تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Evaluate hydration status
b. Assess tissue perfusion
c. Assess respiratory status
d. Maintain thermoregulation

4.32 A newborn's failure to pass meconium within 24 hours after birth indicates which of the following? تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Intussusception
b. Abdominal wall defect
c. Imperforate anus
d. Aganglionic megacolon

4.33 A newborn's failure to pass meconium within 24 hours after birth indicates which of the following? تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Intussusception
b. Imperforate anus
c. Abdominal wall defect
d. Aganglionic megacolon

4.34 When the nurse observes that the infant has respiratory distress and the apical pulse is auscultated on the right side, the nurse suspects the infant has: تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. Pyloric stenosis
b. Cleft palate
c. Diaphragmatic hernia
d. Biliary atresia

4.35 When the child has forceful projectile vomiting and visible gastric peristalsis, the nurse anticipates that the infant has .....

تجميعة صور ميدترمات سابقة

a. Intussusception
b. Extrahepatic biliary atresia
c. Hypertrophic pyloric stenosis
d. Hirschsprung disease

4.36 A priority post-operative care for an infant with cleft lip and palate is: تكرر في 3 مصادر

أسئلة سابقة

a. Protect operative site from infection and injury.
b. Provide family support and education.
c. Spend good time and play with infant.
d. Provide safe environment to prevent falling.

4.37 Preoperatively, an infant with biliary atresia must be provided with.....formula: تكرر في 2 مصادر

أسئلة سابقة

a. Intravenous routine for age.
b. Water soluble vitamin containing.
c. High caloric fatty total parenteral.
d. A regular routine age group.

4.38 Position of choice postoperatively to child with cleft palate is:

أسئلة سابقة

a. Supine
b. Prone
c. Upright
d. One side

4.39 The main goal of preoperative nursing care for a new born with esophageal atresia with tracheoesophageal fistula تكرر في 2 مصادر

أسئلة سابقة

a. Promote growth
b. Maintain skin intact
c. Prevent aspiration
d. Prevent pain

4.40 Which of the following is the nurses' priority in the plan of care for an infant admitted with pyloric stenosis? تكرر في 4 مصادر

أسئلة سابقة

a. Assess respiratory status
b. Maintain thermo-regulation
c. Evaluate hydration status
d. Assess tissue perfusion

4.41 For an infant undergoes surgery to repair tracheo-esophageal fistula, it is essential that the nursing care directed toward prevention of:

أسئلة سابقة

a. Aspiration
b. Abdominal distension
c. Projectile vomiting
d. Dehydration

4.42 When the nurse observes that the infant has respiratory distress and the apical pulse auscultated on the right side; the nurse suspect the infant has: تكرر في 3 مصادر

أسئلة سابقة

a. Pyloric stenosis
b. Biliary atresia
c. Cleft palate
d. Diaphragmatic hernia

4.43 The nurse is reviewing the record of a child with a diagnosis of Hirschsprung's Disease. Which data should the nurse expect to note?

أسئلة سابقة

a. Rkibbon like, foul-smelling stools
b. Jaundice. Pruritus, irritability
c. Increased urine output
d. Projectile vomiting

4.44 A newborn's failure to pass meconium within 24 hours after birth indicate which of the following?

أسئلة سابقة

a. Intussusception
b. Abdominal wall defect
c. Imperforated anus
d. A ganglionic mega colon

4.45 When the child has forceful projectile vomiting and visible gastric peristalsis. Based on these findings, the nurse anticipates that the infant has.....

أسئلة سابقة

a. Intussusception
b. Extra hepatic biliary atresia
c. Hypertrophic pyloric stenosis
d. Hirschsprngs' disease.

4.46 A nurse is assessing a neonate with esophageal atresia. Which finding should the nurse expect to encounter? تكرر في 2 مصادر

أسئلة سابقة

a. Presence of feeding reflexes.
b. Increase in saliva.
c. Gastric tube easily passed.
d. Feeding without difficulty.

4.47 The nurse is caring for a 4-month-old who has just had an isolated cleft lip repaired. Select the best position for the child in the immediate post-operative period تكرر في 2 مصادر

أسئلة سابقة

a. Prone
b. Supine.
c. Left side-lying
d. Right side-lying.

4.48 The nurse monitors a newborn with a suspected diagnosis of imperforate anus. Which is a clinical manifestation associated with this disorder ? تكرر في 2 مصادر

أسئلة سابقة

a. The passage of currant jelly-like stools.
b. Palpated mass in the abdomen.
c. Failure to pass meconium immediately after birth.
d. Bile-stained fecal emesis

4.49 Which clinical manifestations would the nurse expect to assess in the child diagnosed with Hirschsprung's disease? تكرر في 2 مصادر

أسئلة سابقة

a. Constipation and passage of ribbon- like stool.
b. Irritability, severe abdominal cramps and fecal soiling.
c. Respiratory distress and scaphoid abdomen.
d. Bloody diarrhea, fever, and vomiting.

4.50 Which clinical manifestations would the nurse expect to observe while caring for the child diagnosed with intussusception? تكرر في 2 مصادر

أسئلة سابقة

a. Currant jelly stool
b. Foul-smelling stool
c. Dark brown stool
d. Watery stool

4.51 When planning care for the infant with cleft lip and palate, the priority nursing action is: تكرر في 2 مصادر

أسئلة سابقة

a. Burp the baby frequently.
b. Temporarily avoid breastfeed
c. Put the child in supine position.
d. Prevent the baby from crying

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

4.N1 A newborn has frothy saliva, choking and cyanosis with the first feed. The nurse suspects: NCLEX

a. Pyloric stenosis
b. Esophageal atresia with TEF
c. Hirschsprung disease
d. Cleft lip

4.N2 Preoperative positioning of an infant with EA/TEF is: NCLEX

a. Flat supine
b. Head elevated 30–40° (prone or supine) with continuous suction of the pouch
c. Trendelenburg
d. Left side with head down

4.N3 Select ALL expected findings in hypertrophic pyloric stenosis. SELECT ALL

a. Projectile non-bilious vomiting
b. Olive-shaped mass in the right upper quadrant
c. Visible peristaltic waves
d. Currant-jelly stools
e. Hypochloremic metabolic alkalosis

4.N4 After cleft lip repair, which device protects the suture line? NCLEX

a. Mittens only
b. Elbow restraints and a Logan bow
c. Abdominal binder
d. Prone positioning

4.N5 Which finding is typical of Hirschsprung disease in an older child? NCLEX

a. Bloody diarrhea
b. Chronic constipation with ribbon-like, foul-smelling stools and abdominal distension
c. Projectile vomiting
d. Jaundice and clay stools

4.N6 A newborn with congenital diaphragmatic hernia is in respiratory distress. The nurse should position the infant: NCLEX

a. On the unaffected side
b. With head elevated and on the affected side
c. Prone with head down
d. Trendelenburg

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Cleft lip post-op position	Supine or side-lying (never prone) + elbow restraints + Logan bow.	**Cleft palate post-op position**	Prone/side to drain secretions.
Cleft feeding	Upright, special nipple, burp frequently.	**EA/TEF signs**	3 C's: coughing, choking, cyanosis + excessive drooling; NG tube can't pass.
EA/TEF pre-op	NPO, head up 30–40°, continuous suction of upper pouch, gastrostomy.	**Pyloric stenosis**	Projectile non-bilious vomiting, olive mass RUQ, visible peristalsis, metabolic alkalosis.
Pyloric stenosis priority	Correct dehydration/electrolytes before pyloromyotomy.	**Hirschsprung**	Aganglionic colon; no meconium in 24–48 h; ribbon-like foul stools; distension.
Intussusception	Telescoping bowel; colicky pain, sausage mass, currant-jelly stool.	**Imperforate anus**	No anal opening; failure to pass meconium; check with first temperature.
Diaphragmatic hernia	Scaphoid abdomen, respiratory distress, bowel sounds in chest, heart sounds shifted right.	**Biliary atresia**	Jaundice, clay stools, dark urine; formula with fat-soluble vitamins/MCT pre-op (Kasai).

LECTURE 5

High-Risk Neonate

حديث الولادة عالي الخطورة

Part A · Past Exam Questions

أسئلة الدفعات السابقة

5.1 An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر — ميدترم 2020

a. True
b. False

5.2 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر — ميدترم 2020

a. Extremely low
b. Appropriate
c. Moderately low
d. Very low

5.3 Which of the following is a characteristic of prematurity? تكرر في 2 مصادر — ميدترم 2020

a. Well-developed genitalia
b. Sole full of creases
c. Strong grasp reflex
d. Excessive lanugo hair

5.4 The nurse is caring for an infant born at 40 weeks and weighing 1800 g. What would the nurse expect in the chart? — ميدترم 2021

a. Preterm, small for gestational age
b. Full term, large for gestational age
c. Preterm, large for gestational age
d. Full term, small for gestational age

5.5 The nurse is teaching a mother of a newborn about developmental care. The nurse should instruct the mother to: — ميدترم 2021

a. Provide more than one stimulus at a time
b. Stop if there is evidence of overstimulation
c. Offer it during a period of sleeping
d. Provide it for a long period

5.6 The nurse observes a preterm infant with fine hair covering the forehead, back and shoulders. This finding correctly describes: — ميدترم 2021

a. Vernix
b. Milia
c. Lanugo
d. Mottling

5.7 The best intervention to prevent infection in a high-risk neonate begins with: — ميدترم 2022

a. Not wearing a fresh scrub on care
b. Monitoring the mortality rate of neonates
c. Observing continually for signs of infection
d. Meticulous hand washing of all persons in contact

5.8 The best nursing intervention to deal with a preterm neonate in the NICU: تكرر في 2 مصادر

ميدترم 2022

a. Minimal handling during care
b. Use pen in length measurement
c. Cover the neonate with linen
d. Leave the window of the incubator open

5.9 The best nursing intervention to deal with a preterm neonate in the NICU: تكرر في 2 مصادر

ميدترم 2024 (ترم أول 2023-2024)

a. Cover the neonate with linen
b. Minimal handling during care
c. Leave the window of the incubator open
d. Use pen in length measurement

5.10 To prevent heat loss in high-risk neonates resulting from evaporation, the nurse should:

ميدترم 2024 (ترم أول 2023-2024)

a. Warm the crib pad
b. Turn on the radiant warmer
c. Close the room's doors
d. Dry the infant in a warm towel

5.11 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر

كويز 1 – 2022 (نسخة بلاك بورد)

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

5.12 The best nursing intervention to conserve energy in a high-risk neonate is to:

كويز 1 إعادة – ترم ثاني 2023/2024

a. Minimize neonate handling
b. Increase environmental noise
c. Encourage oral feeding
d. Maintain patent airway

5.13 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

كويز 1 (ملف Quiz 1 and 2)

a. Low birth weight
b. Very very low birth weight
c. Very low birth weight
d. Appropriate birth weight

5.14 Which one of the following is considered a characteristic of being a preterm infant? تكرر في 2 مصادر

كويز 1 (ملف Quiz 1 and 2)

a. Low-set ears that will rise later
b. Lack of sole creases on his feet
c. Continuous irritability or crying
d. Scant lanugo on her back and arms

5.15 Extremely low birth weight infants — their weight is: تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. 1500 to 2499 gm
b. 1000 to 1499 gm
c. More than 2500 gm
d. Less than 1000 gm

5.16 Which of the following interventions should be included in the preterm infant's care plan? تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. Use scissors to remove tape
b. Promote developmental care
c. Feed dextrose water immediately after birth
d. Excess handling

5.17 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Moderately low
b. Very low
c. Appropriate
d. Extremely low

5.18 Which of the following is a characteristic of prematurity? تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Well-developed genitalia
b. Sole full of creases
c. Strong grasp reflex
d. Excessive lanugo hair

5.19 The nurse should prevent heat loss in the newborn infant resulting from radiation by:

تجميعة صور ميدترمات سابقة

a. Drying the infant with a warm blanket
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Closing the doors to the room

5.20 An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

5.21 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Extremely low
b. Appropriate
c. Moderately low
d. Very low

5.22 An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

5.23 The nurse should prevent heat loss in the newborn infant resulting from radiation by:

تجميعة صور ميدترمات سابقة

a. Closing the doors to the room
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Drying the infant with a warm blanket

5.24 The best intervention to prevent infection in high-risk neonate begins with:

أسئلة سابقة

a. Not wearing a fresh scrub on care.
b. Monitor mortality rate of neonate.
c. Observing continually for signs of infection.
d. Meticulous hand washing of all persons in contact.

5.25 The best nursing intervention to deal with preterm neonate in NICU:

أسئلة سابقة

a. Minimal handling during care.
b. Use pen in length measurement
c. Cover neonate with linen
d. Leave window of incubator open

5.26 The neonate whose birth weight is 2000 gm. is considered: تكرر في 4 مصادر

أسئلة سابقة

a. Moderately low
b. Very low
c. Appropriate
d. Extremely low

5.27 Which of the following is characteristics of prematurity? أسئلة سابقة

a. Well developed genitalias
b. Sole full with creases
c. Strong grasp reflex
d. Excessive lanugo hair

5.28 The nurse assisted with the delivery of a newborn. Which nursing action is most effective in preventing heat loss by evaporation? أسئلة سابقة

a. Warming the crib pad
b. Closing the doors to the room
c. Drying the infant with a warm blanket,
d. Turning on the overhead radiant warmer

5.29 An infant whose weight falls above the 90th. percentile on the intrauterine growth charts considered large for gestational age تكرر في 3 مصادر أسئلة سابقة

a. True
b. False

5.30 The nurse should prevent the heat loss in the new born infant resulting from radlation by: أسئلة سابقة

a. Closing the doors to the room
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Drying the infant with a warm blanket

5.31 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر أسئلة سابقة

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

5.32 The nurse caring for an infant born at 40 weeks and weighing 1800g. What statement would nurse expect in chart أسئلة سابقة

a. Preterm a small for gestational age.
b. A full term a large for gestational age.
c. A preterm a large for gestational age.
d. A full term a small for gestational age.

5.33 The nurse observes a preterm infants with fine hair covered the forehead, back and shoulders. This finding correctly describes: أسئلة سابقة

a. Vernix.
b. Millia.
c. Lanugo.
d. Mottling.

5.34 Extremely low birth weight infants, their weight is: تكرر في 2 مصادر أسئلة سابقة

a. 1500 to 2499 gm.
b. 1000 to 1499 gm.
c. More than 2500 gm.
d. Less than 1000 gm.

5.35 Which of the following interventions should be included in the preterm ?infant's care plan أسئلة سابقة تكرر في 2 مصادر

a. Use scissors to remove tape
b. Promote developmental care.
c. Feed dextrose water immediately after birth.
d. Excess handling.

5.36 The neonate infant whose birth weight is 2000 gm is considered: أسئلة سابقة

a. Low birth weight.
b. Very very low birth weight.
c. Very low birth weight.
d. Appropriate birth weight.

5.37 Which one of the following is considered a characteristic of being a preterm infant? تكرر في 2 مصادر أسئلة سابقة

a. Low-set ears that will rise later.
b. Lack of sole creases on his feet.
c. Continuous irritability or crying
d. Scant lanugo on her back and arms.

5.38 To prevent heat loss in the high risk neonates resulting from evaporation the nurse should... أسئلة سابقة

a. Warm the crib pad
b. Turn on the radiant warmer
c. Close the room' doors
d. Dry the infant in a warm towel

5.39 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر أسئلة سابقة

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

5.40 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر أسئلة سابقة

a. Appropriate birth weight
b. Low birth weight
c. Very low birth weight
d. Very very low birth weight

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

5.N1 A neonate born at 39 weeks weighs 2100 g. The correct classification is: NCLEX

a. Preterm, AGA
b. Term, SGA (low birth weight)
c. Post-term, LGA
d. Term, AGA

5.N2 Which finding is characteristic of a preterm infant? NCLEX

a. Deep sole creases
b. Abundant lanugo and few sole creases
c. Strong suck and cry
d. Well-formed ear cartilage

5.N3 Select ALL physiological handicaps of the premature infant. SELECT ALL

a. Poor temperature regulation
b. Respiratory difficulties and apnea
c. Susceptibility to infection
d. Strong immunity from maternal antibodies
e. Immature liver (jaundice, bleeding)

5.N4 To prevent heat loss by CONVECTION the nurse should: NCLEX

a. Dry the baby immediately
b. Keep the baby away from drafts and keep incubator portholes closed
c. Warm the stethoscope
d. Place the baby under a radiant warmer

5.N5 **Very low birth weight (VLBW) refers to a birth weight of:** NCLEX

a. Less than 2500 g
b. Less than 1500 g
c. Less than 1000 g
d. 2500–4000 g

5.N6 **Developmental care in the NICU includes:** NCLEX

a. Bright continuous lighting
b. Clustering care, minimal handling and reducing noise
c. Frequent unnecessary stimulation
d. Separating parents from the baby

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Preterm	Born before completion of 37 weeks.	**LBW / VLBW / ELBW**	<2500 g / <1500 g / <1000 g.
SGA / LGA	Below 10th / above 90th percentile for gestational age.	**Preterm features**	Excess lanugo, thin skin, few sole creases, weak reflexes, soft ear cartilage.
Heat loss - evaporation	Prevent by drying immediately with warm towel.	**Heat loss - radiation**	Prevent with radiant warmer/warming surroundings.
Heat loss - convection	Avoid drafts; close incubator portholes.	**Heat loss - conduction**	Warm surfaces (crib pad, scale, stethoscope).
Developmental care	Minimal handling, clustered care, dim light, low noise, nesting.	**Infection prevention**	Meticulous hand washing of everyone in contact.
Premature handicaps	Poor thermoregulation, respiratory problems, infection, immature liver/kidney, poor feeding.	**Complication of prematurity**	Respiratory distress syndrome (surfactant deficiency).

LECTURE 6

Play

اللعب

Part A · Past Exam Questions

أسئلة الدفعات السابقة

6.1 The nurse is observing children playing in the hospital playroom. She would expect to see 4-year-old children playing: تكرر في 2 مصادر — ميدترم 2020

a. Alone with hand-held computer games
b. With their own toys alongside other children
c. Pretend play with other children
d. Competitive board games with older children

6.2 The nurse develops a plan of care including play activities. How does play benefit children? تكرر في 2 مصادر — ميدترم 2020

a. It helps to improve their physical development
b. It increases their immune system
c. It develops only their normal mental abilities
d. It encourages destructive behavior

6.3 (Matching) Play beside, but not with, each other is called: — ميدترم 2020

a. Parallel
b. Solitary
c. Constructive
d. Day dreaming
e. Dramatic

6.4 (Matching) A form of mental play is: — ميدترم 2020

a. Parallel
b. Solitary
c. Constructive
d. Day dreaming
e. Dramatic

6.5 (Matching) Play with own body parts is: — ميدترم 2020

a. Parallel
b. Solitary
c. Constructive
d. Day dreaming
e. Dramatic

6.6 When children start to interact with others during play it is called parallel play. (True / False) — ميدترم 2021

a. True
b. False

6.7 A 2-year-old child likes to play: ميدترم 2022

a. Beside other children, but not with them
b. With other children and borrow and loan toys
c. In groups and share games with each other
d. With himself and his body

6.8 The pediatric nurse should consider, when selecting a toy for a hospitalized child, that it is: ميدترم 2022

a. Large numbers of toys
b. Suitable to age and health
c. High technology and expensive
d. Has one value

6.9 In which type of play do children play beside one another, but not with each other? ميدترم 2024 (ترم أول 2023-2024)

a. Solitary play
b. Parallel play
c. Associative play
d. Constructive play

6.10 While selecting a toy for a hospitalized child, the pediatric nurse considers that the toy should be: تكرر في 2 مصادر ميدترم 2024 (ترم أول 2023-2024)

a. Suitable for age and health
b. Of one value
c. Large in size
d. High technology

6.11 You're developing a plan of care for a 4-year-old that includes play activities. What type will he participate in at this age? كويز 1 – 2021

a. Solitary
b. Cooperative
c. Sociodramatic
d. Parallel

6.12 The nurse develops a plan of care including play activities. How does play benefit children? تكرر في 2 مصادر كويز 1 – 2021

a. It develops only their normal mental capacity
b. It increases their immune system
c. It encourages destructive behavior
d. It helps to improve their physical growth

6.13 How does play benefit children? It ..... كويز 1 – 2022

a. Increases their immune system
b. Develops only their normal mental capacity
c. Helps to improve their physical growth
d. Encourages destructive behavior

6.14 Play for the hospitalized child is necessary to ..... تكرر في 2 مصادر كويز 1 – 2022

a. Improve competition
b. Improve cognitive abilities
c. Improve language abilities
d. Release tension

6.15 The benefit of play for the hospitalized child is: تكرر في 2 مصادر

كويز 1 – 2022 (نسخة بلاك بورد)

a. Provides diversion and relaxation
b. Improves language abilities
c. Feeling of homesickness
d. Improves competition

6.16 At which age group does the child start to interact with others during play?

كويز 1 – 2022 (نسخة بلاك بورد)

a. Infancy
b. Toddlerhood
c. Adolescence
d. Preschool

6.17 Play for the hospitalized child is necessary to: تكرر في 2 مصادر

كويز 1 إعادة – ترم تاني 2023/2024

a. Improve language abilities
b. Release tension
c. Improve cognitive abilities
d. Improve competition

6.18 What is the main characteristic of parallel play?

كويز 1 – ترم تاني 2025

a. Playing together in a shared activity
b. Engaging in pretend play scenarios
c. Observing others without participating
d. Playing alongside other children

6.19 When children sit in a group but each one plays alone, this type of play is:

كويز 1 (ملف Quiz 1 and 2)

a. Sensory
b. Pretend
c. Parallel
d. Solitary

6.20 The nurse is observing a four-year-old child playing in the hospital playroom. Which of the following would she expect to see?

كويز 1 (ملف Quiz 1 and 2)

a. Dramatic play with other children
b. Competitive board games with older children
c. With their own toys alongside other children
d. Alone with hand-held computer games

6.21 As the nurse in charge prepares a playroom for ill children in a pediatric hospital, which play value should the selected toys mostly reflect?

كويز 1 أطفال نظري (Microsoft Forms)

a. Self-awareness
b. Socialization
c. Therapeutic
d. Sensory-motor

6.22 A mother tells the nurse that her preschool girl's toy is an empty box with a stove in it and she calls it "the house". Based on this information, this child's play type is:

كويز 1 أطفال نظري (Microsoft Forms)

a. Parallel play
b. Socio-dramatic play
c. Solitary play
d. Cooperative play

6.23 In which type of play would you see children playing with puzzles alongside, but with no actual sharing?

كويز (ملف DOC-2022)

a. Fantasy
b. Constructive
c. Solitary
d. Parallel

6.24 The type of play that allows children to try out a variety of roles, such as husband, wife, doctor, or police officer: كويز (ملف DOC-2022)

a. Constructive
b. Solitary
c. Sociodramatic
d. Parallel

6.25 Choosing a suitable child's toy is influenced by the child's culture. (True / False) تجميعة صور ميدترمات سابقة

a. True
b. False

6.26 The nurse is observing children playing in the hospital playroom. She would expect to see 4-year-old children playing: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Pretend play with other children
b. Alone with hand-held computer games
c. Competitive board games with older children
d. With their own toys alongside other children

6.27 When a nurse gives a toddler with secretions in the lung a whistle, this will fulfill which play value? تجميعة صور ميدترمات سابقة

a. Therapeutic
b. Socialization
c. Cognitive
d. Creativity

6.28 Which of the following examples represents parallel play? تجميعة صور ميدترمات سابقة

a. Heba plays with a rattle while sitting on her mother's lap
b. Ahmed plays with a car beside Heba, who plays with a doll
c. Ahmed and Aly share their toys
d. Ahmed plays a game with Aly and Hany

6.29 The nurse is observing children playing in the hospital playroom. She would expect to see 4-year-old children playing: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Alone with hand-held computer games
b. With their own toys alongside other children
c. Pretend play with other children
d. Competitive board games with older children

6.30 Which of the following examples represents parallel play? تجميعة صور ميدترمات سابقة

a. Ahmed and Aly share their toys
b. Heba plays with a rattle while sitting on her mother's lap
c. Ahmed plays a game with Aly and Hany
d. Ahmed plays with a car beside Heba, who plays with a doll

6.31 When caregivers choose a toy for their children, they must consider that the ..... تجميعة صور ميدترمات سابقة

a. Toy has removable parts
b. Toy is suitable to developmental age
c. Toy has one value

6.32 When a nurse gives a toddler with secretions in the lung a whistle, this will fulfill which play value? تجميعة صور ميدترمات سابقة

a. Cognitive
b. Socialization
c. Creativity
d. Therapeutic

6.33 Choosing a suitable child's toy is influenced by the child's culture. (True / False) تجميعة صور ميدترمات سابقة

a. True
b. False

6.34 Children are playing with toys, but they are not playing together. This type of play is known as: تجميعة صور ميدترمات سابقة

a. Associative
b. Parallel
c. Cooperative
d. Solitary

6.35 Constructive play is the type of play for preschool children. (True / False) تجميعة صور ميدترمات سابقة

a. True
b. False

6.36 The benefit of play for the hospitalized child is: تكرر في 2 مصادر أسئلة سابقة

a. Provides diversion and relaxation
b. Improves language abilities
c. Feeling of homesickness
d. Improves competition

6.37 While selecting a toy for a hospitalized child, the pediatric nurse considers that the toy should be: تكرر في 2 مصادر أسئلة سابقة

a. Suitable for age and health
b. Of one value
c. Large in size
d. High technology

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

6.N1 A 3-year-old plays with blocks beside another child who plays with cars, without interacting. This is: NCLEX

a. Solitary play
b. Parallel play
c. Associative play
d. Cooperative play

6.N2 Which play activity is most suitable for a 4-year-old? NCLEX

a. Board games with strict rules
b. Dramatic/pretend play such as playing 'doctor'
c. Rattle and mobile
d. Competitive team sports

6.N3 Select ALL functions of play in the hospital. SELECT ALL

a. Provides diversion and relaxation
b. Helps the child feel more secure
c. Lessens stress of separation
d. Allows expression of feelings
e. Increases fear of procedures

6.N4 **Blowing bubbles or a whistle for a child with lung secretions is an example of which play value?** NCLEX

a. Socialization
b. Therapeutic
c. Creativity
d. Moral

6.N5 **Which toy is safest for an 8-month-old?** NCLEX

a. Small marbles
b. Large soft blocks with no detachable parts
c. Balloons
d. Toy with long strings

6.N6 **Constructive play is typical of:** NCLEX

a. Infants
b. Toddlers
c. School-age children
d. Neonates

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Solitary play	Infants; play alone.	**Parallel play**	Toddlers; beside but not with others.
Associative play	Preschool; group play without organization.	**Cooperative play**	School-age; organized with rules and roles.
Sociodramatic play	Preschool; acting roles (doctor, mother) and pretend.	**Constructive play**	School-age; building/creating, problem solving.
Sense-pleasure play	Infants; stimulation of senses, play with body.	**Therapeutic play**	Achieves treatment goals (e.g., blowing whistle for lungs).
Toy selection	Suitable for age, health and safe; influenced by culture.	**Functions of play**	Sensorimotor, intellectual, social, creative, self-awareness, moral, therapeutic.
Play in hospital	Diversion, relaxation, release of tension, sense of security.	**Day dreaming**	Mental form of play (adolescents/fantasy).

LECTURE 7

Accident Prevention

الوقاية من الحوادث

Part A · Past Exam Questions

أسئلة الدفعات السابقة

7.1 Toddlers are more prone to accidents because of their: (ميدترم 2024 (ترم أول 2023-2024)

a. Curiosity
b. Rank
c. Race
d. Growth

7.2 When the mother keeps all medicines and household products out of the reach of children, this represents: (ميدترم 2024 (ترم أول 2023-2024)

a. Passive preventive strategy
b. Posttraumatic care role
c. Prevent complication
d. Active preventive strategy

7.3 The common site of accidental injuries in children under the age of 15 years is: (كويز 2 – 2021) تكرر في 2 مصادر

a. Home
b. School
c. Street
d. Club

7.4 Which of the following is the most common injury in infants? تكرر في 2 مصادر (كويز 2 – 2021)

a. Motor vehicle injuries
b. Burns
c. Suffocation
d. Drowning

7.5 The incidence of injuries among 2–6-year-old children in Egypt is as follows: (كويز 2 – 2022)

a. Boys higher than girls
b. Girls higher than boys
c. Boys are equal to girls
d. Boys are twice girls

7.6 Which one of the following is considered the most common cause of injuries among rural children? (كويز 2 – 2022)

a. Falls
b. Poisons
c. Drowning
d. Fractures

7.7 The most common indoor accident among 6-month-old infants is: (كويز 2 شعبة 3 (ملف Quiz 1 and 2) تكرر في 2 مصادر

a. Drowning
b. Poisoning
c. Burning
d. Falling

7.8 **Which of the following measures should be followed at home to ensure child safety?** كويز 2 شعبة 3 (ملف Quiz 1 and 2)

a. Medicines are to be kept on the dinner table
b. Electrical plugs must be covered and at low level
c. Kitchen curtains are better to be used than doors
d. Furniture is to be away from the window

7.9 **When the mother keeps all medicines and household products out of the reach from children, this represents** أسئلة سابقة

a. Massive preventive strategy
b. Posttraumatic care role
c. Prevent complication
d. Active preventive strategy

7.10 **Which patients are most at risk for sudden infant death syndrome? (Select all that apply)** أسئلة سابقة

a. A newborn whose mother had 1 drink of alcohol early in pregnancy
b. A 2-month-old infant who sleeps in the bed with his mother
c. A 1-month-old infant who prefers to sleep on his stomach
d. A 1-month-old infant who prefers to sleep on her back
e. A newborn whose mother smokes 2 packs/day

7.11 **What is the leading cause of death in children older than 1 year?** أسئلة سابقة

a. Suicide
b. Chronic respiratory illness
c. Unintentional injuries
d. Congenital anomalies

7.12 **What should the nurse teach parents of a toddler about prevention of accidental poisoning? (Select all that apply)** أسئلة سابقة

a. Keep toxic substances in the garage
b. Remove colorful labels from containers of toxic materials
c. Know the number of the nearest poison control center
d. Caution the child against eating nonedible items such as plants
e. Discard empty poison containers

7.13 **Which groups of infants have a higher risk for SIDS? (Select all that apply)** أسئلة سابقة

a. Male gender
b. Low Apgar scores
c. First-born infants
d. Full-term infants
e. Low birth weight

7.14 **For what injuries are crawling infants most at risk?** أسئلة سابقة

a. Aspiration/poisoning
b. Falls
c. Motor vehicle collisions
d. Burns

7.15 Which statement is true regarding injury prevention in infants and toddlers? أسئلة سابقة

a. Repeated "No" is needed to be effective with this age group.
b. Injury prevention and parent education should be handled on a growth and developmental basis.
c. Falls are the most common injuries for these age groups.
d. All injuries can be prevented with proper education regarding safety measures.

7.16 The most common indoor accident among 6-month-old infants is: تكرر في 2 مصادر أسئلة سابقة

a. Drowning
b. Poisoning
c. Burning
d. Falling

7.17 Which of the following measures should at home to ensure child safety? أسئلة سابقة

a. Medicines are to be kept on dinner table
b. Electrical plugs must be covered and at low level
c. Kitchen curtains are better to be used than doors
d. Furnitures are to be away from the window

7.18 Which of the following is the most common injury in infants? تكرر في 2 مصادر أسئلة سابقة

a. Motor vehicle injuries
b. Burns
c. Suffocation
d. Drowning

7.19 The common site of accidental injuries in children under the age of 15 years is: أسئلة سابقة
تكرر في 2 مصادر

a. Home
b. School
c. Street
d. Club

7.20 The incidence of injuries among 2 – 6 years old children at Egypt is as following أسئلة سابقة

a. Boys higher than girls
b. Girls higher than boys
c. Boys are equal girls
d. Boys are twice girls

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

7.N1 A mother of a 9-month-old asks which injury her crawling infant is most at risk for. The nurse answers: NCLEX

a. Motor vehicle injury as a pedestrian
b. Aspiration of small objects and poisoning
c. Sports injuries
d. Firearm injuries

7.N2 Which is an example of an ACTIVE (behavioral) preventive strategy? NCLEX

a. Child-resistant medicine caps made by the factory
b. Building codes for window guards
c. Parent stores medicines and cleaners in a locked high cabinet
d. Automatic airbags in cars

7.N3 Select ALL teaching points that reduce the risk of SIDS. SELECT ALL

a. Place the infant on the back to sleep
b. Use a firm mattress without pillows or soft toys
c. Bed-share with parents every night
d. Avoid smoking around the infant
e. Overdress the infant during sleep

7.N4 Why are toddlers especially prone to accidents? NCLEX

a. Fully developed judgment
b. Curiosity and increasing mobility with poor awareness of danger
c. Peer pressure
d. Risk-taking to impress friends

7.N5 In the epidemiological model of injury, a hot cup of tea that scalds a toddler represents the: NCLEX

a. Host
b. Agent
c. Environment
d. Vector-free factor

7.N6 Which statement about injuries among Egyptian children is correct? NCLEX

a. Girls have more injuries than boys
b. Drowning and animal bites occur mainly among rural children
c. Injuries are rare under 6 years
d. Burns are the least common injury

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Accident	Unintentional event causing injury from external forces or environmental hazards, without deliberate intent.	**Injury**	Damage to the body from acute exposure to thermal, mechanical, electrical or chemical energy, or absence of heat/oxygen.
Impairment	Losing part/all of a limb or having a defective organ or body mechanism.	**Disability**	Loss or reduction of functional ability.
Handicap	Permanent disadvantage or restriction of activity caused by disability.	**Epidemiological model**	Host (child) + Agent (energy/vector) + Environment.
Active prevention	Requires repeated action by the person (e.g., parent locks medicines).	**Passive prevention**	Protects automatically without individual action (e.g., child-resistant caps, airbags).
Most common non-fatal injury in young children	Falls.	**Leading cause of death after age 1**	Unintentional injuries.

Egypt: injuries 2–6 years	Higher in boys than girls; drowning & animal bites mostly in rural children.
Crawling infant hazard	Aspiration of small objects and poisoning.

LECTURE 8

Breast Feeding

الرضاعة الطبيعية

Part A · Past Exam Questions

أسئلة الدفعات السابقة

8.1 **The nurse plans to instruct the mother about measures to prevent breast engorgement. The nurse should tell her to:** ميدترم 2022

a. Breastfeed the neonate frequently
b. Wear a supportive brassiere with nipple shields
c. Restrict fluid intake for 24 to 48 hours
d. Feed the neonate for only 5 minutes per side on the first day

8.2 **A new mother asks why she should feed her newborn colostrum. The nurse's most appropriate answer is because it is:** ميدترم 2022

a. Helping the mother learn how to breastfeed before she goes home
b. High in antibodies, protein, vitamins, and minerals
c. Not real milk
d. Lower in calories than milk and should be supplemented by formula

8.3 **The nurse should instruct the mother that colostrum provides the neonate with:** ميدترم 2024 (ترم أول 2023-2024)

a. More fats than breast milk has
b. Required vitamin K for the neonate
c. Iron needed till age of 6 months
d. Antibodies that the neonate lacks

8.4 **Which of the following statements is correct when teaching a mother about breastfeeding?** ميدترم 2024 (ترم أول 2023-2024)

a. Initiate breastfeeding within 4 hours after birth
b. Encourage breastfeeding daily every 3 hours
c. Give only breast milk for the first 6 months of age
d. Allow artificial teats or pacifiers to breastfed infants

8.5 **A new mother asks whether she should feed her newborn colostrum. The nurse's most appropriate answer is:** كويز 2 – 2021

a. Colostrum is high in antibodies, protein, vitamins, and minerals
b. Colostrum is lower in calories than milk and should be supplemented by formula
c. Colostrum is important as it is rich in iron
d. Colostrum is unnecessary for newborns

8.6 The nurse plans to instruct the mother about the measures to prevent breast engorgement: كويز 2 – 2021

a. Wear a supportive brassiere with nipple shields
b. Administer fluid intake for 24 to 48 hours
c. Breastfeed the neonate frequently
d. Feed the neonate for only 5 minutes per side on the first day

8.7 The indicators of successful breastfeeding include the following EXCEPT: كويز (ملف pediatric Quiz)

a. Urinary output and defecation
b. Excessive sweating
c. Increase in body weight
d. Calm sleeping

8.8 Which of the following statements is most relevant to exclusive breastfeeding? كويز (ملف pediatric Quiz)

a. Early initiation and for six months
b. Day and night, scheduled
c. On demand and allowing water
d. For six months at daytime only

8.9 To ensure successful breastfeeding the child must: كويز (ملف pediatric Quiz)

a. Play with his mother
b. Sleep before feeding
c. Latch on well to the breast
d. Be bathed before feeding

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

8.N1 A mother asks why her baby should get colostrum. The best answer is that colostrum is: NCLEX

a. Lower in protein than mature milk
b. Rich in antibodies (IgA) and acts as the baby's first immunization
c. Mainly water and has no nutritional value
d. High in fat to help weight gain

8.N2 Which hormone is responsible for the milk let-down (ejection) reflex? NCLEX

a. Prolactin
b. Oxytocin
c. Estrogen
d. Progesterone

8.N3 Select ALL signs that a breastfed infant is receiving enough milk. SELECT ALL

a. 6 or more wet diapers per day
b. Steady weight gain
c. Calm sleeping after feeds
d. Excessive sweating during feeds
e. Crying immediately after every feed

8.N4 Which nutrient is NOT sufficient in breast milk and should be supplemented? NCLEX

a. Vitamin C
b. Vitamin D
c. Lactose
d. Water

8.N5 According to WHO/UNICEF, breastfeeding should be initiated: NCLEX

a. Within 1 hour of birth
b. After 6 hours
c. After the first bath
d. When milk 'comes in' on day 3

8.N6 Complementary feeding should start at: NCLEX

a. 2 months
b. 4 months
c. 6 months while breastfeeding continues
d. 12 months after stopping breastfeeding

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Colostrum	First thick yellow milk; rich in protein and IgA; first immunization; laxative effect.	**Prolactin reflex**	Anterior pituitary; milk production in response to suckling (supply = demand).
Oxytocin reflex	Posterior pituitary; let-down/ejection; reduced by worry, pain, stress.	**Foremilk vs hindmilk**	Foremilk: watery, quenches thirst. Hindmilk: fat-rich, gives satiety and calories.
Breast milk whey:casein	About 90:10 early — easy to digest.	**Vitamin lacking in breast milk**	Vitamin D.
Exclusive breastfeeding	Only breast milk (no water) for the first 6 months.	**Initiation**	Within 1 hour of birth (WHO step 4).
Signs of good attachment	Mouth wide open, lower lip turned out, chin touches breast, more areola above than below.	**Adequate intake signs**	≥6 wet diapers/day, weight gain, calm sleep after feeds.
Complementary feeding	Start at 6 months; does not replace breast milk.	**Benefits to mother**	Lactational amenorrhea, uterine involution, less breast/ovarian cancer, less anemia.

LECTURE 9

CNS Congenital Anomalies

العيوب الخلقية في الجهاز العصبي

Part A · Past Exam Questions

أسئلة الدفعات السابقة

9.1 ميدترم 2020

Khalid is a 3-day-old infant admitted to the pediatric surgical ward with a myelomeningocele and hydrocephalus. While caring for Khalid the nurse should:

a. Assist with the insertion of a central line
b. Position him on the right side
c. Support with mechanical ventilator
d. Cover the sac with a moist and sterile dressing

9.2 ميدترم 2020

Khalid is a 3-day-old infant admitted with a myelomeningocele and hydrocephalus. The recommended position for Khalid is: تكرر في 2 مصادر

a. Prone
b. Trendelenburg's
c. Fowler's
d. Supine

9.3 ميدترم 2020

(Matching) Anencephaly is best described as:

a. Incomplete fusion of the vertebrae that may be signaled by an overlying tuft of hair
b. A protrusion of nerve tissue and meninges through a skull defect
c. Absence of the cerebral hemispheres
d. A sac of meninges and CSF protruding through a vertebral defect without nerve tissue

9.4 ميدترم 2020

(Matching) Encephalocele is best described as:

a. Incomplete fusion of the vertebrae that may be signaled by an overlying tuft of hair
b. A protrusion of nerve tissue and meninges through a skull defect
c. Absence of the cerebral hemispheres
d. A sac of meninges and CSF protruding through a vertebral defect without nerve tissue

9.5 ميدترم 2020

(Matching) Spina bifida occulta is best described as:

a. Incomplete fusion of the vertebrae that may be signaled by an overlying tuft of hair
b. A protrusion of nerve tissue and meninges through a skull defect
c. Absence of the cerebral hemispheres
d. A sac of meninges and CSF protruding through a vertebral defect without nerve tissue

9.6 ميدترم 2020

(Matching) Meningocele is best described as:

a. Incomplete fusion of the vertebrae that may be signaled by an overlying tuft of hair
b. A protrusion of nerve tissue and meninges through a skull defect
c. Absence of the cerebral hemispheres
d. A sac of meninges and CSF protruding through a vertebral defect without nerve tissue

9.7 When caring for an infant with spina bifida, which technique is most important in recognizing possible hydrocephalus? — ميدترم 2021

a. Obtaining skull X-ray
b. Measuring head circumference
c. Performing a lumbar puncture
d. Magnetic resonance imaging

9.8 The nurse informed doctors that the WBC in CSF culture is up to 1000 and the CSF is turbid for a child with a VP shunt. What is the nurse's interpretation? — ميدترم 2021

a. Normal shunt function
b. Need for repeated CSF culture
c. Infection in CSF is indicated
d. Normal WBC and color of CSF

9.9 The major cause of congenital hydrocephalus is brain tumor. (True / False) — ميدترم 2021

a. True
b. False

9.10 What is the major cause of congenital hydrocephalus that interrupts ventricular circulation in the brain? — ميدترم 2022

a. Meningitis
b. Head injury
c. Brain tumor
d. Aqueduct stenosis

9.11 The nurse informed doctors that the WBC in CSF culture is up to 1000 for a child connected to a VP shunt. What is the nurse's interpretation? — ميدترم 2022

a. Indicative of CSF infection
b. Need for repeated CSF culture
c. Normal WBC of CSF
d. Normal shunt function

9.12 The most important preoperative nursing action for a child with myelomeningocele is: — ميدترم 2024 (ترم أول 2023-2024)

a. Examination of the membranous cyst
b. Diapering and sponge bath for the child
c. Apply a sterile, moist, non-adherent dressing
d. Keep the infant in the prone position

9.13 Which of the following nursing actions is most important in caring for a child with hydrocephalus? — ميدترم 2024 (ترم أول 2023-2024)

a. Measure vital signs
b. Assess weight
c. Monitor urinary output
d. Measure head circumference

9.14 One of the following is the most important nursing intervention for an infant born with spina bifida cystica: — كويز 2 – 2022

a. Provide psychological support to the family
b. Assess neurologic function
c. Assess growth measurement
d. Cover the defect with a saline-soaked dressing

9.15 Postoperative nursing care of a child after insertion of a ventriculoperitoneal shunt is ..... كويز 2 – 2022

a. Administer narcotics for pain control
b. Check the urine for glucose and protein
c. Monitor for increased temperature
d. Assess body weight

9.16 When caring for an infant with spina bifida, which measure is most important in recognizing possible hydrocephalus? كويز 2 شعبة 3 (ملف Quiz 1 and 2)

a. Obtain skull X-ray
b. Measure head circumference
c. Perform a lumbar puncture
d. Magnetic resonance imaging

9.17 Which priority intervention should the nurse perform for the child's position after insertion of a shunt to relieve hydrocephalus? كويز 2 شعبة 3 (ملف Quiz 1 and 2)

a. Supine position
b. On the non-operative site
c. Semi-Fowler position
d. Upright position

9.18 Khalid is a 3-day-old infant admitted with a myelomeningocele and hydrocephalus. The recommended position for Khalid is: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Fowler's
b. Trendelenburg's
c. Prone
d. Supine

9.19 Khalid is a 3-day-old infant admitted with a myelomeningocele and hydrocephalus. While caring for Khalid the nurse should: تجميعة صور ميدترمات سابقة

a. Assist with the insertion of a central line
b. Position him on the right side
c. Support with mechanical ventilator
d. Cover the sac with a moist and sterile dressing

9.20 Khalid is a 3-day-old infant admitted with a myelomeningocele and hydrocephalus. The recommended position for Khalid is: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Prone
b. Trendelenburg's
c. Fowler's
d. Supine

9.21 Khalid is a 3-day-old infant admitted with a myelomeningocele and hydrocephalus. The recommended position for Khalid is: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Fowler's
b. Trendelenburg's
c. Prone
d. Supine

9.22 The most common complication for VP shunt is: أسئلة سابقة

a. Shunt disconnection
b. Seizures
c. Fever
d. Obstruction

9.23 **An infant is born with a sac protruding through the spine, containing CSF, a portion of meninges and nerve roots. the condition is referred to as** أسئلة سابقة

a. Meningocele
b. Myelomeningocele
c. Spina bifida occulta
d. Anencephaly

9.24 **The nurse is caring for an infant with a myelomeningocele. the parents ask the nurse why they keep measuring the babys head circumference. select the nurses best response** أسئلة سابقة

a. Babies heads are measured to ensure growth is on track
b. Babies with a myelomeningocele are at risk for hydrocephalus, which shows up as an increase in head size
c. Because your baby has an opening on the spinal cord, your infant is at risk for meningitis, which can show up as an increase in head size
d. Many infants with myelomeningocele have microcephaly, which can show up as a decrease in head size

9.25 **A parent of a newborn diagnosed with myelomeningocele asks what is a common long term complication. the nurses best response is** أسئلة سابقة

a. Learning disabilities
b. UTI
c. Hydrocephalus
d. Decubitus ulcers and skin breakdown

9.26 **Which would be included in the plan of care for a hospitalized newborn following surgical repair of a myelomeningocele? (Select all that apply)** أسئلة سابقة

a. Skull X-rays
b. Daily head circumference measurements
c. MRI scan
d. Vital signs every 6 hours
e. Holding to breastfeed

9.27 **Which is included in the plan of care for a newborn who has myelomeningocele** أسئلة سابقة

a. Place the child in the prone position with a sterile dry dressing over the defect. slowly begin oral gastric feeds to prevent the development of nectrozing enterocolitis
b. Place the child in the prone position with a sterile dry dressing over the defect. begin IV fluids to prevent dehydration
c. Place the child in the prone position with a sterile moist dressing over the defect. slowly begin oral gastric feeds to prevent the development of necrotizing enterocolitis
d. Place the child in the prone position with a sterile moist dressing over the defect. begin IV fluids to prevent dehydrations

9.28 **Which does the nurse include in the postoperative plan of care of a child with myelomeningocele following ligament release?** أسئلة سابقة

a. Encourage resuming a regular diet, beginning slowly with bland foods such as bananas
b. Encourage the child to blow balloons to increase deep breathing
c. Assist the child to change positions to avoid skin breakdown
d. Provide education on dietary requirements to prevent obesity and skin breakdown

9.29 **The nurse is working in a school health clinic and a teen mentions that her older sister just had a baby born with a myelomeningocele. the teen is wondering if there is anything she can do to prevent this from happening to her baby when she decides to have children. which is the best response** أسئلة سابقة

a. Take a multivtaim with folic acid daily
b. Eat more fruits and vegetables daily
c. Have breakfast every morning
d. There is nothing that can be done to decrease the risk

9.30 **The nurse is caring for an infant with myelomeningocele who is going to surgery later today for closure of the sac. which would be a priority nursing diagnosis before surgery** أسئلة سابقة

a. Alteration in parent infant bonding
b. Altered growth and development
c. Risk of infection
d. Risk for weight loss

9.31 **Which should be included in the plan of care for a newborn with myelomeningocele who will have a surgical repair tomorrow** أسئلة سابقة

a. Offer formula every 3 hours
b. Turn the infant back to front every 2 hours
c. Place a wet dressing on the sac
d. Provide pain medication every 4 hours

9.32 **Which should the nurse do first when caring for an infant who just had a repair of a myelomeningocele** أسئلة سابقة

a. Weight diapers for 24 hour urine output
b. Measure head circumference
c. Offer clear fluids
d. Assess for infection

9.33 **Which should be the priority ND for a 12 hour old newborn with a myelomeningocele at L2** أسئلة سابقة

a. Altered bowel elimination r/t neuro deficits
b. Potential for infection r/t the physical defect
c. Altered nutrition r/t neuro deficit
d. Disturbance in self concept r/t physical disability

9.34 **Over the last week an infant with repiared myelomeningocele has had a high pitched cry and been irritable. length, weight and head circumference have been at the 50th percentile. today length is at the 50th percentile, weight is at the 70th percentile and head circumference is at the 90th percentile. the nurse should do which of the following** أسئلة سابقة

a. Tell the parent this is normal for an infant with a repaired myelomeningocele
b. Tell the parent this might mean the baby has increased icp
c. Suspect the baby's icp is low because of a leak
d. Refer the baby to the nuerologist for follow up care

أسئلة سابقة

9.35 **Which should the nurse prepare the parents of an infant for the following surgical repair and closure of a myelomeningocele shortly after birth. the infant will:**

a. Not need any long term management and should be considered cured
b. Not bo at risk for UTI or movement problems
c. Have continual drainage of CSF needed frequent dressing changes
d. Need lifelong management of urinary, orthopedic and neurological problems

أسئلة سابقة

9.36 **A newborn with a repaired myelomeningocele is assessed for hydrocephalus. which would the nurse expect in an infant with hydrocephalus**

a. Low pitched cry and depressed fontanel
b. Low pitched cry and bulging fontanel
c. Bulging fontanel and downwardly rotated eyes
d. Depressed fontanel and upwardly rotated eyes

أسئلة سابقة

9.37 **A nurse is caring for a neonate born 12 hours ago with a myelomeningocele. Which assessment finding should be reported immediately to the PCP**

a. Axillary temp of 102.2 (fever)
b. Dribbling of urine
c. No lower limb movement
d. Talipes equinavaraus

أسئلة سابقة

9.38 **An infant has returned to the peds unit after repair of a meylomeningocele. The nruse notices that the infant has had no urine output in the past 2 hours. Which nursing intervention would be most appropriate**

a. Perform credes maneuver on the infants bladder
b. Catheterize the infants bladder
c. Ask the mother to breast feed the infant
d. Increase the IV fluid rate

أسئلة سابقة

9.39 **The parents of a child newly dx with myelomeningocele ask the nurse why surgical repair needs to be done immediately. Which reponse would be most accurate**

a. Its done to rapidly restore neural pathways to the legs
b. Its done to decrease the possibility of infection and further cord damage
c. Its done to expose the spinal cord defect and allow an individualized therapeutic strategy
d. Its done to remove excess nerve tissue from the vertebral canal and to decrease pressure on the cord

أسئلة سابقة

9.40 **The most important pre-operative nursing action for child with myelomeningocele is:**

a. Examine of the membranous cyst
b. Diapering and sponged path for the child
c. Apply sterile, moist, non-adherent dressing
d. Keep infant in the prone position

أسئلة سابقة

9.41 **The nurse is working in the PICU caring for an infant who has just returned from having a shunt placed for hydrocephalus. Which position initially will be most beneficial for this child?**

a. Semi-Fowler in an infant seat
b. Flat in a crib
c. Trendelenberg
d. In crib with HOB elevated 90 degrees

9.42 Early signs and symptoms of hydrocephalus in an infant include which of the following? أسئلة سابقة

a. Confusion, headache, diplopia
b. Rapid head growth, poor feeding, confusion
c. Papilledema, irritability, headache
d. Full fontanels, poor feeding, rapid head growth

9.43 Which of the following nursing action is most important in caring of child with hydrocephalus? أسئلة سابقة

a. Measure vital signs
b. Assess weight
c. Monitor urinary output
d. Measure head circumference

9.44 The nurse informed doctors about the WBC result in CSF culture up to 1000 for child connected with VP shunt, what the nurse interpretation of this result أسئلة سابقة

a. Indicted of CSF infection
b. Need for repeated CSF culture
c. Normal WBC of CSF
d. Normal shunt function

9.45 Khalid is 3 days old infant admitted to the pediatric surgical ward with a myelomeningocele and hydrocephalus while caring for Khalid the nurse should أسئلة سابقة

a. Assist with the insertion of a central line
b. Position him on the right side
c. Support with mechanical ventilator
d. Cover sac with a moist and sterile dressing

9.46 Khalid is 3 days old infant admitted to the pediatric surgical ward with a myelomeningocele and hydrocephalus the recommended position for Khalid is أسئلة سابقة

a. Prone
b. Trendelburg's
c. Fowler's
d. Supine

9.47 What is the major cause for congenital hydrocephalus that interrupted ventricle circulation in the brain? أسئلة سابقة

a. Meningitis
b. Head injury
c. Brain tumor
d. Aqueduct stenosis

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

9.N1 The priority preoperative care for a newborn with myelomeningocele is: NCLEX

a. Supine positioning
b. Covering the sac with a sterile, moist, non-adherent dressing
c. Applying a dry pressure dressing
d. Allowing the parents to hold the baby upright

9.N2 **Which is an early sign of hydrocephalus in an infant?** NCLEX

a. Headache and diplopia
b. Rapidly increasing head circumference with bulging fontanel
c. Papilledema
d. Decreased head circumference

9.N3 **Select ALL signs of VP shunt malfunction or infection.** SELECT ALL

a. Vomiting and irritability
b. Bulging fontanel
c. Fever
d. Increased appetite
e. High-pitched cry

9.N4 **Which advice helps prevent neural tube defects?** NCLEX

a. Avoid vitamin supplements
b. Folic acid 0.4 mg daily before conception and in early pregnancy
c. High-protein diet in the 3rd trimester
d. Iron supplements only

9.N5 **Spina bifida occulta is often signaled by:** NCLEX

a. A large sac on the back
b. A tuft of hair or dimple over the lumbosacral spine
c. Absent cerebral hemispheres
d. Open skull defect

9.N6 **Immediately after VP shunt insertion, the infant is usually positioned:** NCLEX

a. On the operative side
b. Flat, on the non-operative side
c. Head down
d. Sitting upright

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Anencephaly	Absence of cerebral hemispheres; incompatible with life.	**Encephalocele**	Protrusion of brain tissue and meninges through a skull defect.
Spina bifida occulta	Hidden vertebral defect; tuft of hair or dimple.	**Meningocele**	Sac with meninges and CSF only; no nerve tissue.
Myelomeningocele	Sac with meninges, CSF and spinal cord/nerve roots.	**MMC pre-op**	Prone, sterile moist non-adherent dressing, protect from feces/urine, measure head circumference.
MMC complications	Hydrocephalus, UTI, neurogenic bladder, paralysis.	**Hydrocephalus cause**	Commonly aqueduct stenosis (obstructive).
Hydrocephalus signs	Large head, bulging fontanel, sunset eyes, high-pitched cry, irritability.	**Most important hydrocephalus assessment**	Daily head circumference.

VP shunt complications	Obstruction (most common) and infection.
Prevention	Folic acid before conception and in early pregnancy.

LECTURE 10

Rheumatic Fever

الحمى الروماتيزمية

Part A · Past Exam Questions

أسئلة الدفعات السابقة

10.1 Which of the following nursing diagnoses is most appropriate for a child with rheumatic fever? — ميدترم 2022

a. High risk for injury
b. Imbalanced nutrition: more than requirements
c. Delayed growth and development
d. Impaired gas exchange

10.2 A nurse is preparing a child for discharge after being diagnosed with rheumatic fever. Which is the correct instruction? — ميدترم 2022

a. Only give penicillin for dental work
b. Give aspirin for management of chorea
c. Give penicillin until 21 years old
d. Give penicillin for one month total

10.3 Which of the following instructions would the nurse include in a teaching plan that focuses on prevention of rheumatic fever? — ميدترم 2024 (ترم أول 2023-2024)

a. Use corticosteroids to reduce inflammation
b. Provide antibiotic before dental work
c. Give penicillin to patients for treatment
d. Seek immediate treatment for throat infection

10.4 One minor manifestation of rheumatic fever is: — ميدترم 2024 (ترم أول 2023-2024)

a. Erythema marginatum
b. Polyarthritis and chorea
c. Increased sedimentation rate
d. Painless subcutaneous nodules

10.5 Symptoms of rheumatic fever typically appear how long after a streptococcal infection? — أسئلة سابقة

a. 1–2 days
b. 3–5 days
c. 2–3 weeks
d. 1 month

10.6 Which joints are most commonly affected by pain or swelling in rheumatic fever? — أسئلة سابقة

a. Shoulders and hips
b. Knees, ankles, elbows and wrists
c. Fingers and toes
d. Spine and neck

10.7 What type of joint pain is often seen in rheumatic fever? — أسئلة سابقة

a. Persistent pain in one joint
b. Pain that moves from joint to joint (migratory)
c. Pain relieved by rest
d. Pain without swelling or redness

10.8 Which is NOT a typical manifestation of rheumatic fever? أسئلة سابقة

a. Fever
b. Joint swelling
c. Skin rash
d. Headache

10.9 Under the 2015 Modified Jones Criteria, which symptom most likely indicates rheumatic fever? أسئلة سابقة

a. Severe headache
b. Hot or tender joints
c. Abdominal pain
d. Nausea and vomiting

10.10 Which body systems are typically affected in rheumatic fever? أسئلة سابقة

a. Respiratory, digestive and immune
b. Musculoskeletal, cardiovascular and nervous
c. Digestive, endocrine and respiratory
d. Cardiovascular, digestive and reproductive

10.11 Under the 2015 Modified Jones Criteria, which is a common sign? أسئلة سابقة

a. Skin lesions
b. Fever and joint pain
c. Gastrointestinal distress
d. Shortness of breath

10.12 Which is the characteristic skin manifestation of rheumatic fever? أسئلة سابقة

a. Erythema nodosum
b. Erythema marginatum
c. Psoriasis
d. Vitiligo

10.13 What is the primary treatment of rheumatic fever (eradicating the streptococcus)? أسئلة سابقة

a. Corticosteroids
b. Aspirin
c. Penicillin
d. Sulfadiazine

10.14 Which is NOT a sign or symptom of rheumatic fever? أسئلة سابقة

a. Joint pain
b. Fatigue
c. Chest pain
d. Nausea

10.15 One minor manifestation for rheumatic fever is أسئلة سابقة

a. Erythema marginatum
b. Poly arthritis and chorea
c. Increased sedimentation rate
d. Painless subcutaneous nodules

10.16 A nurse is preparing child for discharge after diagnosed with rheumatic fever that is the following instruction أسئلة سابقة

a. Only give penicillin for dental work
b. Give aspirin for management chorea
c. Give penicillin until 21 years old
d. Give penicillin for one month total

أسئلة سابقة

10.17 Which of the following nursing diagnosis is most appropriate for child with rheumatic fever?

a. High risk for injury
b. Imbalance munition more than requirement
c. Delayed growth and development
d. Impaired gas exchange

أسئلة سابقة

10.18 In case of rheumatic heart fever with Myocarditis, the chest x-ray reveals normal heart size

a. True **b.** False

أسئلة سابقة

10.19 Rheumatic fever is inflammation of the joints, heart, skin, and nervous system that occurs in children as a complication of untreated streptococcal infection of the throat. While most people with rheumatic fever recover, which part of the body may be permanently damaged in a small percentage of people?

a. Digestive tract **b.** Ears
c. Heart **d.** Lungs

أسئلة سابقة

10.20 Which of the following is usually the first symptom of rheumatic fever?

a. Chest pain or palpitations **b.** Jerky, uncontrollable movements
c. Joint pain and fever **d.** Rash

أسئلة سابقة

10.21 Penicillin is safely given to patients by the IV route. (True / False)

a. True **b.** False

أسئلة سابقة

10.22 Each of the following drugs is used to treat rheumatic fever in children EXCEPT:

a. Acetaminophen **b.** Antibiotics
c. Anti-inflammatory drugs **d.** Corticosteroids

أسئلة سابقة

10.23 Patient with polyarthritis and carditis without evident of CHF (congestive heart failure) is treated with

a. IV salicylates **b.** Oral salicylates
c. Corticosteroid **d.** Both B and C

أسئلة سابقة

10.24 Which symptom presented is NOT a sign of rheumatic fever?

a. Truncal rash **b.** Joint tenderness
c. Changes in vision **d.** Nausea

أسئلة سابقة

10.25 Rheumatic fever is caused by which of the following bacteria?

a. Pseudomonas **b.** Staphylococcus
c. Streptococcus **d.** Shingella

10.26 Which of the following factors in the patient's history and physical is LEAST relevant to the diagnosis of Rheumatic fever? أسئلة سابقة

a. Sore throat from 3 weeks previous
b. Living in the current conditions of Flint, MI
c. The mother's history of type-2 diabetes
d. A possible genetic factor from the father who's history is unknown

10.27 The provider suspects rheumatic fever and orders blood work. Which results indicate inflammation and cardiac involvement? (Select all that apply) أسئلة سابقة

a. Elevated C-reactive protein
b. Decreased erythrocyte sedimentation rate
c. Elevated WBC count
d. Decreased hemoglobin and hematocrit

10.28 According to the Jones criteria, which sets of findings would indicate a positive diagnosis of rheumatic fever? (Select all that apply) أسئلة سابقة

a. Carditis, fever, and an elevated WBC count
b. Positive strep throat culture, arthritis, chorea
c. Recent scarlet fever, carditis, fever, arthralgia
d. Elevated C-reactive protein, carditis, fever

10.29 Acute Rheumatic Fever (ARF) is primarily caused by which bacterial species? أسئلة سابقة

a. Staphylococcus aureus
b. Streptococcus pneumoniae
c. Streptococcus agalactiae
d. Streptococcus pyogenes (Group A Streptococcus)

10.30 Molecular mimicry in Acute Rheumatic Fever refers to: أسئلة سابقة

a. The ability of bacteria to mimic host cells.
b. The interaction between antibodies and antigens.
c. The cross-reactivity between bacterial antigens and host tissues.
d. The production of toxins by Streptococcus pyogenes.

10.31 Which of the following clinical manifestations is NOT a major criterion for diagnosing Acute Rheumatic Fever? أسئلة سابقة

a. Carditis
b. Chorea
c. Elevated C-reactive protein
d. Subcutaneous nodules

10.32 Patient with polyarthritis and carditis with evident cardiomegaly is treated with أسئلة سابقة

a. IV salicylates
b. Oral salicylates
c. Corticosteroid
d. Adrenaline

10.33 Which of the following in not considered among supportive therapies for patient with carditis

أسئلة سابقة

a. Digoxin
b. Diuretics
c. Oxygen
d. Increasing fluid and salt intake

10.34 Which clinical manifestation of Acute Rheumatic Fever involves rapid, jerky movements and behavioral changes?

أسئلة سابقة

a. Carditis
b. Arthritis
c. Chorea
d. Subcutaneous nodules

10.35 Which genetic marker can influence an individual's susceptibility to developing Acute Rheumatic Fever after a streptococcal infection?

أسئلة سابقة

a. HLA-B27
b. Rh factor
c. ABO blood type
d. Genetic markers associated with the immune response

10.36 Which of the following is a major clinical manifestation of Acute Rheumatic Fever that involves inflammation of the heart?

أسئلة سابقة

a. Chorea
b. Arthritis
c. Carditis
d. Subcutaneous nodules

10.37 Which of the following is NOT a potential complication of Acute Rheumatic Fever?

أسئلة سابقة

a. Rheumatic heart disease
b. Glomerulonephritis
c. Meningitis
d. Chorea

10.38 Which of the following is a minor criterion for diagnosing Acute Rheumatic Fever?

أسئلة سابقة

a. Carditis
b. Arthritis
c. Elevated erythrocyte count
d. Subcutaneous nodules

10.39 What is the gold standard for diagnosing Group A Streptococcus infection in the laboratory?

أسئلة سابقة

a. Rapid strep test
b. Gram staining
c. Throat culture
d. Molecular PCR test

10.40 Which of the following medications is NOT commonly used in the treatment of Acute Rheumatic Fever?

أسئلة سابقة

a. Penicillin
b. Nonsteroidal anti-inflammatory drugs (NSAIDs)
c. Corticosteroids
d. Antibiotics targeting Gram-negative bacteria

10.41 Rheumatic fever primarily affects which of the following body systems? أسئلة سابقة

a. Digestive system
b. Musculoskeletal system
c. Respiratory system
d. Nervous system

10.42 Which of the following is the characteristic feature of rheumatic fever? أسئلة سابقة

a. Joint pain and swelling
b. Skin rash
c. Heart murmurs
d. All of the above

10.43 Which of the following age groups are at higher risk for rheumatic fever? Select the correct answer أسئلة سابقة

a. Adults 50 and above
b. Children ages 5-15
c. Infants
d. Women ages 20-40

10.44 The best way to prevent rheumatic fever is: أسئلة سابقة

a. Avoiding contact with infected individuals
b. Getting vaccinated
c. Early diagnosis and treatment of streptococcal infections
d. Washing hands frequently

10.45 Rheumatic fever can lead to long-term complications, primarily affecting the..... أسئلة سابقة

a. Heart
b. Kidneys
c. Liver
d. Lungs

10.46 The condition characterized by scarring and damage to the heart valves as a consequence of rheumatic fever is called..... أسئلة سابقة

a. Coronary artery disease
b. Rheumatic heart disease
c. Rheumatoid arthritis
d. Septicemia

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

10.N1 Rheumatic fever is an autoimmune reaction that follows infection with: NCLEX

a. Staphylococcus aureus
b. Group A beta-hemolytic streptococcus
c. Pneumococcus
d. E. coli

10.N2 Select ALL MAJOR Jones criteria. SELECT ALL

a. Carditis
b. Polyarthritis
c. Chorea
d. Erythema marginatum
e. Subcutaneous nodules
f. Elevated ESR

10.N3 **Secondary prophylaxis after rheumatic fever usually consists of:** NCLEX

a. Oral aspirin for one month
b. Benzathine penicillin IM every 3–4 weeks for years
c. Corticosteroids for life
d. Antibiotics only during dental work

10.N4 **A child with RF has chorea. The priority nursing diagnosis is:** NCLEX

a. Impaired gas exchange
b. Risk for injury
c. Constipation
d. Excess fluid volume

10.N5 **Which finding indicates carditis in a child with rheumatic fever?** NCLEX

a. New heart murmur and tachycardia out of proportion to fever
b. Migratory joint pain
c. Ring-shaped rash
d. Painless nodules on extensor surfaces

10.N6 **The joint involvement in rheumatic fever is typically:** NCLEX

a. Small joints, symmetric and deforming
b. Migratory polyarthritis of large joints, non-deforming
c. Spinal only
d. Single joint with permanent damage

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Cause	Autoimmune reaction 2–3 weeks after group A beta-hemolytic strep pharyngitis.	**Molecular mimicry**	Cross-reactivity between streptococcal antigens and host tissues.
Major Jones criteria	Carditis, polyarthritis, chorea, erythema marginatum, subcutaneous nodules.	**Minor criteria**	Fever, arthralgia, raised ESR/CRP, prolonged PR interval.
Evidence of strep	Positive throat culture (gold standard) or raised ASO titer.	**Arthritis**	Migratory, large joints, non-deforming.
Chorea (Sydenham)	Involuntary jerky movements, emotional lability; risk for injury.	**Carditis**	Only manifestation causing permanent damage → rheumatic heart disease (mitral).
Treatment	Penicillin, salicylates for arthritis, steroids for carditis with cardiomegaly/CHF, bed rest.	**Secondary prophylaxis**	Benzathine penicillin IM every 3–4 weeks (often until 21 years).
Primary prevention	Prompt treatment of strep throat.	**Age group**	Children 5–15 years.

LECTURE 11

Nephrotic Syndrome & AGN

المتلازمة الكلوية والتهاب الكلى الحاد

Part A · Past Exam Questions

أسئلة الدفعات السابقة

11.1 The nurse assessing a child with acute post-streptococcal glomerulonephritis should be alert for which finding? — ميدترم 2022

a. Weight gain
b. Increased urine output
c. Hypotension
d. Tea-colored urine

11.2 A nurse is assigned to care for a child with nephrotic syndrome. Which important parameter should be assessed daily? تكرر في 2 مصادر — ميدترم 2022

a. Blood urea nitrogen
b. Weight
c. Height
d. Activity tolerance

11.3 A parent of a child with glomerulonephritis asks why the urine is an abnormal color. The nurse's best response is: — ميدترم 2024 (ترم أول 2023-2024)

a. The urine is discolored because the child is receiving antihypertensives
b. There is blood in the child's urine that causes it to be tea-colored
c. Change of diet often causes the urine to be tea-colored
d. The child's urine is very concentrated, so it appears abnormal

11.4 Which area of potential disturbance should be a prime consideration when planning ongoing nursing care for a child with nephrotic syndrome? — ميدترم 2024 (ترم أول 2023-2024)

a. Body image
b. Sexual maturation
c. Muscle coordination
d. Intellectual development

11.5 A parent asks the nurse how to prevent the child from having minimal change nephrotic syndrome again. The nurse's best response is: تكرر في 2 مصادر — كويز 2 – 2021

a. Keep the child away from sick playmates
b. It is rare for a child to have a relapse after recovery
c. Decrease sodium in the child's diet
d. Unfortunately, relapses cannot be prevented

11.6 Which of the following signs and symptoms are characteristic of nephrotic syndrome? — كويز 2 – 2022

a. Hypertension, edema, hematuria
b. Hypertension, edema, proteinuria
c. Gross hematuria, fever, proteinuria
d. Poor appetite, edema, proteinuria

11.7 A parent asks the nurse how to prevent the child from having nephrotic syndrome again. The nurse's best response is: كويز 2 – 2022

a. Unfortunately, relapses cannot be prevented
b. Decrease sodium in the child's diet
c. Keep the child away from sick playmates
d. It is rare for a child to have a relapse after recovery

11.8 Which finding typically is the earliest sign of improvement in a child with acute glomerulonephritis? كويز 2 – 2022

a. Increased urine output
b. Increased appetite
c. Increased energy level
d. Decreased diarrhea

11.9 Nephrotic syndrome is primarily characterized by: أسئلة سابقة

a. Hematuria
b. High creatinine levels
c. Massive proteinuria
d. Urinary tract obstruction

11.10 The most common cause of nephrotic syndrome in children is: أسئلة سابقة

a. FSGS
b. Membranous nephropathy
c. Minimal change disease
d. Lupus nephritis

11.11 A key difference between nephritis and nephrosis is: أسئلة سابقة

a. Foamy urine in nephritis
b. Inflammation in nephrosis
c. Edema in nephritis only
d. Inflammation is a main feature of nephritis

11.12 Which of the following is a complication of nephrotic syndrome? أسئلة سابقة

a. Hypotension
b. Weak immune system (infection)
c. Hyperkalemia
d. Anemia

11.13 Which diagnostic tool confirms kidney tissue damage? أسئلة سابقة

a. CT scan
b. Urinalysis
c. Blood test
d. Kidney biopsy

11.14 Nephritis is commonly caused by: أسئلة سابقة

a. Hyperlipidemia
b. Autoimmune diseases
c. Minimal change disease
d. Nephrotoxic drugs only

11.15 Which medication is used to manage high cholesterol in nephrosis? أسئلة سابقة

a. Furosemide
b. Atorvastatin
c. Penicillin
d. Prednisone

أسئلة سابقة

11.16 In nephritis, which symptom is common?

a. Massive proteinuria
b. Foamy urine
c. Hypertension
d. Severe hyperlipidemia

أسئلة سابقة

11.17 Which dietary recommendation is common for both nephritis and nephrosis?

a. High-sodium intake
b. High-protein diet
c. Low-protein, low-sodium diet
d. No dietary restrictions

أسئلة سابقة

11.18 What is the purpose of corticosteroids in nephritis treatment?

a. Lower blood pressure
b. Control fluid retention
c. Treat infections
d. Reduce inflammation

أسئلة سابقة

11.19 The nurse a child with acute post streptococcal glomerulonephritis should be alert for which finding

a. Weight gain
b. Increased urine output
c. Hypotension
d. Tea colored urine

أسئلة سابقة

11.20 Which finding typically is the earliest sign of improvement in case of child with acute glomerulonephritis

a. Increased urine output
b. Increased appetite
c. Increased energy level

أسئلة سابقة

11.21 Glomerulonephritis refers to inflammation of the glomeruli of the kidneys. (True / False)

a. True
b. False

أسئلة سابقة

11.22 Acute post-streptococcal GN (APSGN) is the less common post-infectious renal disease. (True / False)

a. True
b. False

أسئلة سابقة

11.23 Acute glomerulonephritis incidence in males to females is:

a. 2:1
b. 1:2
c. 1:1

أسئلة سابقة

11.24 The following are considered methods of diagnosing AGN:

a. Urinalysis
b. History and clinical manifestations
c. Physical examination
d. Laboratory findings
e. All of the above

أسئلة سابقة

11.25 Therapeutic treatment of AGN includes all of the following except

a. Antibiotic such as penicillin
b. Antihypertensive
c. Anticonvulsant
d. Diuretics
e. Antihistaminic

11.26 **A parent of a child with glomerulonephritis asks the nurse why the urine is abnormal color the nurse's best response is** أسئلة سابقة

a. The urine is discolored because the child is receiving antihypertensive
b. There is blood in the child's urine that causes it to be tea colored
c. Change diet often causes the urine to be tea colored
d. Child's urine is very concentrated so it appears to be abnormal

11.27 **Nursing management of AGN includes** أسئلة سابقة

a. Restrict protein intake
b. Maintain potassium and fluid restrictions
c. Monitor heart rate and RR

11.28 **A 5-year-old male with AGN has mild facial edema, tea-colored urine and BP 165/110. Which interventions will you include in his plan of care? (Select all that apply)** أسئلة سابقة

a. Initiate and maintain a high-sodium diet daily
b. Monitor intake and output hourly
c. Encourage ambulation every 2 hours while awake
d. Assess color of urine after every void
e. Weigh the patient daily on a standing scale
f. Encourage the patient to consume 4 L of fluid per day

11.29 **While providing care to a pediatric patient with acute glomerulonephritis, you note the urine output to be 10 mL/hr. The patient weighs 30 lbs. As the nurse, you will want to limit what type of foods from the patient's diet?** أسئلة سابقة

a. Calcium-rich foods
b. Potassium-rich foods
c. Purine -rich foods
d. None of the above because the patient's urinary output is normal based on the patient's weight.

11.30 **A patient who is experiencing poststreptococcal glomerulonephritis has edema mainly in the face and around the eyes. As the nurse, you know to expect the edema to be more prominent during the?** أسئلة سابقة

a. Evening
b. Afternoon
c. Morning
d. Bedtime

11.31 **Which patient below is at MOST RISK for developing acute glomerulonephritis?** أسئلة سابقة

a. A 3 year old male who has a positive ASO titer.
b. A 5 year old male who is recovering from an appendectomy.
c. An 18 year old male who is diagnosed with HIV.
d. A 6 year old female newly diagnosed with measles.

أسئلة سابقة

11.32 **Which of the following are NOT signs and symptoms of acute post-streptococcal glomerulonephritis? (Select all that apply)**

a. Hypotension
b. Increased glomerular filtration rate
c. Cola-colored urine
d. Massive proteinuria
e. Elevated BUN and creatinine
f. Mild swelling in the face or eyes

أسئلة سابقة

11.33 **Post-streptococcal glomerulonephritis is a type of NEPHROTIC syndrome, meaning both RBCs and protein leak from the inflamed glomerulus. (True / False)**

a. True
b. False

أسئلة سابقة

11.34 **Within the past month, the admission rate of patients with poststreptococcal glomerulonephritis has doubled on your unit. You are proving an in-service to your colleagues about this condition. Which statement is CORRECT about this condition?**

a. "This condition tends to present 6 months after a strep infection of the throat or skin."
b. "It is important the patient consumes a diet rich in potassium based foods due to the risk of hypokalemia."
c. "Patients are less likely to experience hematuria with this condition."
d. "This condition is not caused by the streptococcal bacteria attacking the glomerulus, but by the immune system's response to the bacteria by creating an antigen-antibody complex which inflames the glomerulus."

أسئلة سابقة

11.35 **Glomerulonephritis (nephritic syndrome) is a kidney disorder. This disorder affects the glomeruli (clusters of microscopic blood vessels with small pores through which blood is filtered). Glomerulonephritis is characterized by swelling (edema), high blood pressure, and which of the following?**

a. Red blood cells in urine
b. An inability to urinate
c. An increased desire for salt
d. Extreme thirst

أسئلة سابقة

11.36 **Acute glomerulonephritis most often occurs as a complication of which of the following?**

a. A back injury
b. Poor nutrition
c. A streptococcal infection
d. A viral infection

أسئلة سابقة

11.37 **Urinalysis results include proteinuria, many red blood cells and red blood cell casts, and 1-2 white blood cells per high power field. Which of the following diseases best fits with these findings?**

a. Acute cystitis
b. Acute pyelonephritis
c. Chronic pyelonephritis
d. Acute post-streptococcal glomerulonephritis
e. Minimal change disease

أسئلة سابقة

11.38 **The nurse is caring for a child admitted with acute glomerulonephritis. Which of the following clinical manifestations would likely have been noted in the child with this diagnosis?**

a. Smoky colored urine
b. Jaundiced skin
c. Strawberry red tongue
d. Loose, dark stools

11.39 A child diagnosed acute glomerulonephritis will most likely have a history of which of the following? أسئلة سابقة

a. Recent illness such as strep throat
b. Hemorrhage or history of bruising easily
c. Sibling diagnosed with the same disease
d. Hearing loss with impaired speech development

11.40 A school nurse is trying to prevent poststreptococcal glomerulonephritis in children. Which of the following would be the best way to prevent this? أسئلة سابقة

a. Prophylactic antibiotics after strep throat are important.
b. Tell parents to give ibuprofen if their child has a sore throat.
c. All children in the child's class should be tested for strep throat if there is a positive.
d. Encourage the child to take all the antibiotics if diagnosed with strep throat.

11.41 You obtain a history from the mother of a child with glomerulonephritis about how he became ill. Which of the following would you expect her to tell you she noticed? أسئلة سابقة

a. Headache, loss of appetite
b. Dark brown urine
c. Loss of weight, oliguria
d. Diuresis and pallor

11.42 The caregiver of a child being treated at home for acute glomerulonephritis calls the nurse reporting that her daughter has just had a convulsion. The child is resting comfortably but the caregiver would like to know what to do. The nurse would instruct the caregiver to do which of the following actions? أسئلة سابقة

a. Give the child fluids and report back to the nurse in a few hours.
b. Take the child's blood pressure and report the findings to the nurse while the nurse is still on the phone.
c. Give the child a diuretic and report back to the nurse in a few hours.
d. Weigh the child in the same clothes

11.43 A symptom often seen in acute glomerulonephritis is edema. The most common site the edema is first noted is in which area of the body? أسئلة سابقة

a. Sacrum
b. Eyes
c. Hands
d. Ankles

11.44 A child has been admitted to the hospital with a diagnosis of acute glomerulonephritis. During history-taking the nurse first asks the child's parents about a recent history of: أسئلة سابقة

a. Bleeding ulcer
b. Deep vein thrombosis
c. Myocardial infarction
d. Streptococcal infection

11.45 The child has been diagnosed to have glomerulonephritis. What should the nurse observe in the urine? أسئلة سابقة

a. Blood
b. Pus
c. White blood cells
d. Glucose

أسئلة سابقة

11.46 You're collecting a urine sample on a patient who is experiencing proteinuria due to nephrotic syndrome. As the nurse, you know the urine will appear:

a. Tea-colored
b. Orange and frothy
c. Dark and foamy
d. Straw-colored

أسئلة سابقة

11.47 The characteristic features of nephrotic syndrome are?

a. Heavy proteinuria (greater than 3.5 g/24 hr in adults or 40 mg/sq.m/hr in children)
b. Hypoalbunemia (less than 2.5 g/dL)
c. Edema
d. Hyperlipidemia
e. All the above

أسئلة سابقة

11.48 Which of the following are uncommon findings in a patient with idiopathic nephrotic syndrome?

a. Abdominal pain
b. Anorexia and Irritability
c. Hypertension
d. Gross hematuria
e. Both C and D

أسئلة سابقة

11.49 Which of the following statements about nephrotic syndrome are true?

a. Most children (90%) with nephrotic syndrome have a form of idiopathic nephrotic syndrome
b. Minimal change disease (85%) is the most common cause of idiopathic nephrotic syndrome
c. The remaining 10% who do not have idiopathic NS have secondary NS related to systemic or glomerular diseases such as membranous nephropathy or membranoproliferative glomerulonephritis
d. All the above statements are true

أسئلة سابقة

11.50 A patient is diagnosed with nephrotic syndrome. Which signs and symptoms are common in this condition?

a. Foamy, frothy urine
b. Massive proteinuria
c. Edema
d. Hypoalbuminemia
e. All of the above
f. None of the above

أسئلة سابقة

11.51 Select the most common type of medications that may be ordered by a physician to treat nephrotic syndrome:

a. Corticosteroids
b. Antihypertensives
c. Diuretics
d. Both a and c

أسئلة سابقة

11.52 Each of the following features are characteristic of the nephrotic syndrome except:

a. Marked proteinuria
b. Hypoalbuminemia
c. Edema
d. Hypertension

أسئلة سابقة

11.53 The child with nephrotic syndrome who has ascites and difficulty breathing is probably most comfortable sleeping in which position?

a. Supine
b. Sims' position
c. Prone
d. Fowler's

أسئلة سابقة

11.54 **In caring for a child with nephrotic syndrome, which of the following interventions will be included in the child's plan of care?**

a. Ambulating three to four times a day
b. Testing the urine for glucose levels regularly
c. Increasing fluid intake by 50 cc an hour
d. Weighing on the same scale each day

أسئلة سابقة

11.55 **The nurse is caring for a child with nephrotic syndrome. The child is noted to have edema. The edema would most likely be seen where on this child?**

a. Sacrum
b. Abdomen
c. Eyes and ankles
d. Fingers

أسئلة سابقة

11.56 **The nurse is caring for an 8-year-old hospitalized child with nephrotic syndrome. Which of the following nursing interventions would be appropriate for this child?**

a. Test the urine for ketones twice a day
b. Weigh the child once a week.
c. Administer antipyretics as needed.
d. Measure the abdominal girth daily.

أسئلة سابقة

11.57 **An 11-year-old boy presents to your office with pitting edema and proteinuria exceeding 3.5g in 24 hours. You suspect that this patient has experienced a loss of polyanions in his glomerular basement membranes. Which of the following findings would confirm your diagnosis?**

a. WBC casts in the urine
b. RBC casts in the urine
c. Selective albuminuria
d. Negatively birefringent crystals in the urine
e. Bence-Jones proteinuria

أسئلة سابقة

11.58 **Three-year-old Carlo has been admitted to the pediatric unit with a tentative diagnosis of nephrotic syndrome. The diagnosis of Idiopathic Nephrotic Syndrome has been confirmed. Which unexpected finding would the nurse report?**

a. Proteinuria
b. Distended abdomen
c. Blood in the urine
d. Elevated serum lipid levels

أسئلة سابقة

11.59 **Three-year-old Carlo has been admitted to the pediatric unit with a tentative diagnosis of nephrotic syndrome. Carlo's potential for impairment of skin integrity is related to:**

a. Joint inflammation
b. Drug therapy
c. Edema
d. Generalized body rash

أسئلة سابقة

11.60 **Prednisone is prescribed for Carlo. The nurse evaluate its effectiveness by**

a. Checking his BP every 4 hours
b. Checking his urine for protein
c. Weighing him each morning before breakfast
d. Observing him for behavioral changes

11.61 A 6 year old male is diagnosed with nephrotic syndrome. In your nursing care plan you will include which of the following as a nursing diagnosis for this patient? أسئلة سابقة

a. Risk for infection
b. Deficient fluid volume
c. Constipation
d. Overflow urinary incontinence

11.62 Which patient below is NOT at risk for developing nephrotic syndrome? أسئلة سابقة

a. An 8 year old male with diabetes mellitus.
b. A 5 year old female diagnosed with minimal change disease.
c. A 10 year old male with Lupus.
d. A 7 year old male recently diagnosed with Goodpasture's Syndrome.

11.63 The mother of a child recently diagnosed with nephrotic syndrome asks how to identify early signs of relapse. You tell her to monitor the child for: أسئلة سابقة

a. Weight loss
b. Protein in the urine using an over-the-counter kit
c. Tea-colored urine
d. Swelling in the legs, hands, face, or abdomen
e. A and C
f. B and D

11.64 A student describes the signs and symptoms of nephrotic syndrome. Which statements require re-education? (Select all that apply) أسئلة سابقة

a. Slight proteinuria
b. Hypoalbuminemia
c. Edema
d. Hyperlipidemia
e. Tea-colored urine
f. Hypertension

11.65 You're providing care to a 6-year-old male patient who is receiving treatment for nephrotic syndrome. Which assessment finding below requires you to notify the physician immediately? أسئلة سابقة

a. Frothy, dark urine
b. Redden area on the patient's left leg that is swollen and warm
c. Elevated lipid level on morning labs
d. Urine test results that shows proteinuria

11.66 As the nurse, you know that it is important to implement a low sodium diet for a patient with nephrotic syndrome. However, it is important to implement what other type of diet due to another complication associated with this syndrome? أسئلة سابقة

a. Low-phosphate
b. Low-fat
c. High-carbohydrate
d. Low-potassium

11.67 A parent asks the nurse how to prevent the child from having minimal change nephrotic syndrome again the nurse's best response تكرر في 2 مصادر

أسئلة سابقة

a. Keep the child away from sick playmates
b. It is rare for a child to have a relapse after recovery
c. Decrease sodium in child
d. Unfortunately, relapses, cannot be prevented

11.68 A nurse is assigned to care for a child with nephrotic syndrome which important parameter should be assessed daily تكرر في 2 مصادر

أسئلة سابقة

a. Blood urea nitrogen
b. Weight
c. Height
d. Activity tolerance

11.69 Which of the following signs and symptoms are characteristics of the nephrotic syndrome?

أسئلة سابقة

a. Hypertension, edema, hematuria
b. Hypertension, edema, proteinuria
c. Gross hematuria, fever, proteinuria
d. Por appetite, edema, proteinuria

11.70 A parent asks the nurse how to prevent the child form having nephrotic syndrome again the nurse's best response is

أسئلة سابقة

a. Unfortunately, relapses, cannot be prevented
b. Decrease sodium in child diet
c. Keep the child away from sick playmates
d. It is rare a child to have a relapse after recovery

11.71 Which of the following areas of potential disturbances should be a prime consideration when planning ongoing nursing care for child with nephrotic syndrome

أسئلة سابقة

a. Body image
b. Sexual maturation
c. Muscle coordination
d. Intellectual development

11.72 Nephrotic syndrome is a clinical state that characterized by

أسئلة سابقة

a. Massive proteinuria
b. Hypoproteinemia
c. Sever edema
d. Hyperlipidemia
e. All of the above

11.73 Secondary NS is idiopathic nephrosis or minimal change nephrotic syndrome

أسئلة سابقة

a. True
b. False

11.74 All of the following considered therapeutic management of nephrotic syndrome except

أسئلة سابقة

a. Corticosteroids
b. Antibiotic
c. Alkylating agent
d. Analgesic
e. I.V albumin

11.75 Relapse considered one of complication of NS

أسئلة سابقة

a. True
b. False

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

11.N1 The classic triad of nephrotic syndrome includes: NCLEX

- **a.** Hematuria, hypertension, oliguria
- **b.** Massive proteinuria, hypoalbuminemia, edema
- **c.** Fever, rash, arthritis
- **d.** Polyuria, polydipsia, weight loss

11.N2 A child with AGN has tea-colored urine. The nurse explains this is due to: NCLEX

- **a.** Medication
- **b.** Hematuria
- **c.** Bilirubin
- **d.** Concentrated urine

11.N3 Select ALL appropriate nursing actions for a child with nephrotic syndrome. SELECT ALL

- **a.** Weigh daily on the same scale
- **b.** Measure abdominal girth daily
- **c.** Protect from infection
- **d.** Encourage a high-sodium diet
- **e.** Provide meticulous skin care

11.N4 AGN most commonly follows which infection? NCLEX

- **a.** Viral hepatitis
- **b.** Group A beta-hemolytic streptococcal throat or skin infection
- **c.** Urinary tract infection
- **d.** Measles

11.N5 The earliest sign of improvement in AGN is: NCLEX

- **a.** Increased appetite
- **b.** Increased urine output (diuresis)
- **c.** Normal ESR
- **d.** Weight gain

11.N6 The first-line drug for idiopathic (minimal change) nephrotic syndrome is: NCLEX

- **a.** Penicillin
- **b.** Prednisone (corticosteroids)
- **c.** Digoxin
- **d.** Furosemide alone

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Nephrotic syndrome	Massive proteinuria, hypoalbuminemia, edema, hyperlipidemia.	**Minimal change disease**	Most common cause of childhood NS (~85%); steroid-responsive.
NS urine	Dark, frothy/foamy.	**NS edema**	Periorbital (morning) → generalized, ascites.
NS key nursing	Daily weight, abdominal girth, skin care, infection prevention, body image.	**NS relapse prevention**	Avoid sick contacts (infection triggers relapse).

AGN	Immune-complex glomerulonephritis 1–3 weeks after group A strep infection.
AGN urine	Tea/cola-colored (hematuria), oliguria, mild proteinuria.
AGN signs	Periorbital edema, hypertension, oliguria, hematuria.
AGN improvement sign	Increased urine output.
AGN with convulsion	Suspect hypertensive encephalopathy — check BP.
Nephritic vs nephrotic	Nephritic: RBCs + protein, HTN. Nephrotic: massive protein only.

LECTURE 12

Respiratory Distress Syndrome (RDS)

متلازمة الضائقة التنفسية عند حديثي الولادة

Part A · Past Exam Questions

أسئلة الدفعات السابقة

12.1 (Matching) Grade I respiratory distress is manifested by: ميدترم 2020

a. Rapid respiration and nasal flaring
b. Intercostal and subcostal retractions
c. Expiratory grunting
d. Cyanosis and altered conscious level
e. Wheezy chest and tachypnea

12.2 (Matching) Grade II respiratory distress is manifested by: ميدترم 2020

a. Rapid respiration and nasal flaring
b. Intercostal and subcostal retractions
c. Expiratory grunting
d. Cyanosis and altered conscious level
e. Wheezy chest and tachypnea

12.3 (Matching) Grade III respiratory distress is manifested by: ميدترم 2020

a. Rapid respiration and nasal flaring
b. Intercostal and subcostal retractions
c. Expiratory grunting
d. Cyanosis and altered conscious level
e. Wheezy chest and tachypnea

12.4 (Matching) Grade IV respiratory distress is manifested by: ميدترم 2020

a. Rapid respiration and nasal flaring
b. Intercostal and subcostal retractions
c. Expiratory grunting
d. Cyanosis and altered conscious level
e. Wheezy chest and tachypnea

12.5 Which of the following findings would alert the nurse to notify the primary health care provider when assessing the neonate? تكرر في 2 مصادر ميدترم 2021

a. Red reflex of the eyes
b. Equal bilateral air entry
c. Respiratory rate 40 br/min
d. Sternum retraction

12.6 Which of the following is a specific treatment of respiratory distress syndrome? تكرر في 2 مصادر ميدترم 2021

a. Antibiotic medication
b. Mechanical ventilation
c. Exchange transfusion
d. Surfactant replacement

12.7 A neonate's oral cavity is filled with mucus and he became cyanosed. What is the first action of the nurse? تكرر في 2 مصادر ميدترم 2024 (ترم أول 2023-2024)

a. Recording the incident
b. Insert nasogastric tube
c. Frequent suctioning
d. Administering O2 therapy

12.8 The nurse could teach the parents that the main function of surfactant is to: تكرر في 2 مصادر

ميدترم 2024 (ترم أول 2023-2024)

a. Decrease lung secretion
b. Prevent lung collapse
c. Produce lung collapse
d. Decrease lung inflation

12.9 Which assessment signs alert the nurse to the possibility of respiratory distress syndrome? تكرر في 2 مصادر

كويز 2 – 2022

a. Hypotension and bradycardia
b. Tachypnea and retractions
c. Acrocyanosis and grunting
d. The presence of a barrel chest

12.10 Which nursing diagnosis is priority for a preterm infant with respiratory distress syndrome?

كويز 2 – 2022

a. Fluid volume deficit
b. Risk for infection
c. Impaired gas exchange
d. Altered body temperature

12.11 While developing a plan of care for a newborn with RDS, which is considered the main nursing diagnosis?

كويز 2 شعبة 3 (ملف Quiz 1 and 2)

a. Body image disturbance
b. Altered body temperature
c. Impaired gas exchange
d. Altered bowel elimination

12.12 An early sign of neonatal respiratory distress syndrome is grunting. (True / False) تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

12.13 An early sign of neonatal respiratory distress syndrome is grunting. (True / False) تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

12.14 The nurse could teach the parents that the main function of surfactant is to تكرر في 2 مصادر

أسئلة سابقة

a. Decrease lung secretion
b. Prevent the lung collapse
c. Produce lung collapse
d. Decrease lung inflation

12.15 A neonate's oral cavity is filled with mucus and he became cyanosed what is the first action of the nurse to do تكرر في 2 مصادر

أسئلة سابقة

a. Recording the incident
b. Insert nasogastric tube
c. Frequently suctioning
d. Administering O2 therapy

12.16 Which assessment signs alert the nurse to the possibility of respiratory distress syndrome تكرر في 2 مصادر

أسئلة سابقة

a. Hypotension and bradycardia
b. Tachypnea and retractions
c. Acrocyanosis and grunting
d. The presence of a barrel chest

12.17 The major main aim of surfactant therapy is as follow

أسئلة سابقة

a. Increase air leaks of lung
b. Decreased oxygen requirement
c. Decrease surface tension of alveoli
d. Increase surface tension of alveoli

12.18 An early sign of neonatal respiratory distress syndrome is grunting. (True / False)

تكرر في 2 مصادر

أسئلة سابقة

a. True
b. False

12.19 In which weeks of gestation of the following for starting corticosteroid therapy for pregnant women to overcome fetal lung prematurity

أسئلة سابقة

a. 20 - 30
b. 22 – 32
c. 24 – 34
d. 25 – 35

12.20 Which of the following nursing diagnosis is priority for preterm infant with respiratory distress syndrome?

أسئلة سابقة

a. Fluid volume deficit
b. Risk for infection
c. Impaired gas exchange
d. Altered body temperature

12.21 Which of the following findings would alert the nurse to notify the primary health care provider when assessing the neonates? تكرر في 2 مصادر

أسئلة سابقة

a. Red reflex of the eyes
b. Equal bilateral air entry
c. Respiratory rate 40br/min
d. Sternum retraction

12.22 The preterm infant is diagnosed with respiratory distress syndrome which is caused by alteration in the body's secretion of

أسئلة سابقة

a. Progesterone
b. Adrenaline
c. Surfactant
d. Somatotropin

12.23 Which of the following is a specific treatment of respiratory distress syndrome

تكرر في 2 مصادر

أسئلة سابقة

a. Antibiotic medication
b. Mechanical ventilation
c. Exchange transfusion
d. Surfactant replacement

12.24 Hyaline membrane disease also called RDS

أسئلة سابقة

a. True
b. False

12.25 All of the following considered risk factors of RDS except

أسئلة سابقة

a. Cold stress
b. Maternal diabetes
c. Limited birth
d. Prematurity

12.26 Which of the following is considered a physiological abnormality of RDS?

أسئلة سابقة

a. Decreased alveolar surface tension
b. Decreased alveolar ventilation
c. Increased lung volume

12.27 Intercostal retraction is a sign to diagnose which grade of RDS

أسئلة سابقة

a. 3
b. 2
c. 1
d. 4

12.28 Antenatal corticosteroid therapy are indicated in pregnant women 24 – 34 week's gestation at high risk of preterm delivery within the next 7 days

أسئلة سابقة

a. True
b. False

12.29 Type of surfactant that is an extract of calf lung:

أسئلة سابقة

a. Curosurf
b. Survanta
c. Infasurf

12.30 What of the following consider nursing diagnosis of patient with RDS

أسئلة سابقة

a. Impaired gas exchange
b. Imbalanced nutrition less than body requirements
c. Ineffective family coping
d. All of the above

12.31 RDS is the most common cause of respiratory distress in premature infants, correlating with structural and functional immaturity of the heart. (True / False)

أسئلة سابقة

a. True
b. False

12.32 The more the gestational age and birth weight, the more the incidence of RDS. (True / False)

أسئلة سابقة

a. True
b. False

12.33 It's alse known as hyaline membrane disease:

أسئلة سابقة

a. Rheumatic fever
b. Acute kidney injury
c. Heart failure
d. Respiratory distress syndrome

12.34 Pulmonary surfactant production begins at

أسئلة سابقة

a. 34 weeks
b. 26 weeks
c. 24 weeks
d. 32 weeks

12.35 Infants born after 32 weeks have a much higher chance of survival than those born at 24 weeks.

أسئلة سابقة

a. True
b. False

12.36 Adequate amounts of pulmonary surfactant is produced to prevent atelectasis at about

أسئلة سابقة

a. 34 weeks
b. 26 weeks
c. 24 weeks
d. 32 weeks

أسئلة سابقة

12.37 All of the following are considered predisposing and risk factor of RDS, Except:

a. Perinatal asphyxia
b. Cold stress
c. Female gender
d. Genetic disorders

أسئلة سابقة

12.38 In RDS, Mortality rate increases with decreasing gestational age and in LBW newborns.

a. True
b. False

أسئلة سابقة

12.39 Mature levels of pulmonary surfactant are usually present after

a. 34 weeks
b. 26 weeks
c. 35 weeks
d. 24 weeks

أسئلة سابقة

12.40 In RDS, the interference with gas exchange results in which of the following except:

a. Hypoxia (decreased O2 saturation)
b. Increased PH
c. Increased PaCO2
d. Increased work of breathing

أسئلة سابقة

12.41 A newborn infant with RDS may be presented with which of the following, except:

a. Tachypnea
b. Nasal flaring
c. Hypertension
d. Intercostal retraction

أسئلة سابقة

12.42 In most cases of RDS, symptoms and signs reach a peak within 3 days

a. True
b. False

أسئلة سابقة

12.43 Which of the following electrolytes imbalance associated with RDS:

a. Respiratory acidosis
b. Metabolic alkalosis
c. Respiratory alkalosis
d. Metabolic acidosis

أسئلة سابقة

12.44 A preterm infant presented with dyspnea, fatigue, nasal flaring, expiratory retraction, his respiratory rate is 85br/min, according to the S&S of the infant, the infant is in which grade of RD:

a. Grade I
b. Grade II
c. Grade III
d. Grade IIII

أسئلة سابقة

12.45 Neonatal respiratory distress syndrome (NRDS):

a. Occurs in ~50% of premature births with a weight of <1500 g
b. Always follows premature births before 30-week gestation
c. Is commonly treated by giving the neonate corticosteroids
d. Occurs in 1 in 300 of all births.

أسئلة سابقة

12.46 Four 38-week-gestation gravidas have just delivered. Which of the babies should be monitored closely by the nurse for respiratory distress?

a. The baby whose mother has diabetes mellitus.
b. The baby whose mother has lung cancer.
c. The baby whose mother has hypothyroidism.
d. The baby whose mother has asthma.

12.47 A newborn infant is presented by the following S&S, RR 70 br/min, mild retraction, no cyanosis, mild decrease in air entry, and audible grunting by stethoscope, according to Downe's score, the evaluation of this infant and his diagnosis is: أسئلة سابقة

a. 5 & respiratory distress
b. 3 & no respiratory distress
c. 4 & respiratory distress
d. 8 & impeding respiratory failure

12.48 For the prevention of RDS, the following measures are done, except: أسئلة سابقة

a. Administering tocolytics
b. Surfactant replacement therapy
c. Administering corticosteroids
d. Give beta blockers

12.49 A 1,000-gram neonate is being admitted to the neonatal intensive care unit. The surfactant Survanta (beractant) has just been prescribed to prevent respiratory distress syndrome. Which of the following actions should the nurse take while administering this medication? أسئلة سابقة

a. Flush the intravenous line with normal saline solution.
b. Assist the neonatologist during the intubation procedure.
c. Inject the medication deep into the vastus lateralis muscle.
d. Administer the reconstituted liquid via an oral syringe.

12.50 A 30-week-gestation neonate, 2 hours old, has received Survanta (beractant). Which of the following would indicate a positive response to the medication? أسئلة سابقة

a. Axillary temperature 98.0°F
b. Oxygen saturation 96%
c. Apical heart rate 154 bpm
d. Serum potassium 4.0 mEq/L

12.51 Administration of corticosteroids to women 48 hours before delivery of fetuses helps: أسئلة سابقة

a. Stimulate the production of surfactant
b. Accelerate fetal lung maturation
c. Reduce the incidence of RDS
d. All of the above

12.52 For newborn with RDS, aiming to keep pO2 between أسئلة سابقة

a. 40-70 mmHg
b. 50-80 mmHg
c. 55-90 mmHg
d. 45-75 mmHg

12.53 For an infant with RDS the proper position is: أسئلة سابقة

a. Side lying
b. Left lateral
c. Supine
d. Semi-Fowler's

12.54 The major main aim of surfactant therapy is: كويز 2 – 2022 (ملف Quiz2 pediatric 2022)

a. Increase air leaks of lung
b. Decreased oxygen requirement
c. Decrease surface tension of alveoli
d. Increase surface tension of alveoli

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

12.N1 The primary cause of RDS in preterm infants is: NCLEX

a. Infection
b. Surfactant deficiency
c. Meconium aspiration
d. Congenital heart disease

12.N2 Which is an EARLY sign of RDS? NCLEX

a. Grunting
b. Tachypnea
c. Cyanosis on room air
d. Apnea

12.N3 Select ALL risk factors for RDS. SELECT ALL

a. Prematurity
b. Maternal diabetes
c. Cold stress
d. Perinatal asphyxia
e. Female gender

12.N4 Antenatal corticosteroids are given to mothers at risk of preterm delivery between: NCLEX

a. 20–30 weeks
b. 24–34 weeks
c. 34–40 weeks
d. Any time

12.N5 Surfactant replacement is administered via: NCLEX

a. IV line
b. Endotracheal tube
c. Intramuscular injection
d. Oral syringe

12.N6 The expected ABG in an infant with RDS is: NCLEX

a. Respiratory alkalosis
b. Respiratory and metabolic acidosis
c. Metabolic alkalosis
d. Normal

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

RDS	Hyaline membrane disease; surfactant deficiency in preterm.	**Surfactant**	Decreases alveolar surface tension; prevents collapse.
Production	Begins ~24 wk; adequate ~32 wk; mature ~35 wk.	**Risk factors**	Prematurity, male, maternal DM, cold stress, asphyxia, C-section.
Grade I	Rapid respiration, nasal flaring.	**Grade II**	Intercostal/subcostal retractions.
Grade III	Expiratory grunting.	**Grade IV**	Cyanosis, altered consciousness.
Prevention	Antenatal corticosteroids 24–34 wk, tocolytics.	**Specific treatment**	Surfactant replacement via ET tube.
Support	CPAP, O2 keeping PaO2 50–80, NTE, minimal handling.	**Priority ND**	Impaired gas exchange.

Answer Key — All Lectures

الإجابات

1.1. a — Recognizes that the family is the constant in a child's life

1.2. c — Promoting disease prevention

1.3. a — Promoting disease prevention

1.4. a — Respect

1.5. a — Promoting disease prevention

1.6. d — Promoting disease prevention

1.7. c — Low level of father education

1.8. b — New

1.9. c — Enabling

1.10. b — Poverty

1.11. a — Recognizes that the family is the constant in a child's life

1.12. c — Promoting disease prevention

1.13. a — Accidents

1.14. b — Enabling

1.15. a — Diarrheal disease

1.16. a — Collaboration

1.17. d — Maintains the child's health and prevents diseases

1.18. c — Acute respiratory infections

1.19. a — Enabling

1.20. d — Promoting disease prevention

1.21. c — Recognizes that the family is the constant in a child's life

1.22. d — Acute lower respiratory infections

1.23. a — Birth weight

1.24. b — Social problems

1.25. a — Infections

1.26. c — Enabling

1.27. b — Birth weight

1.28. b — Respect

1.29. a — Promoting disease prevention

1.30. a — Acute lower respiratory tract infections

1.31. b — Promoting disease prevention

1.32. c — Promoting disease prevention

1.33. a — Last

1.34. a — Recognizes that the family is the constant in a child's life

1.35. a — Enabling

1.36. b — Birth weight

1.37. a — Birth weight

1.38. b — Social problems

1.39. a — Infections

1.40. c — Enabling

1.41. a — Respect.

1.42. a — Promoting disease prevention.

1.43. c — Low level of father education.

1.44. b — New

1.45. a — Acute lower respiratory tract infections

1.46. a — Last

1.47. a — Recognizes that the family is the constant in a child's life

1.48. b — Birth weight.

1.49. b — Respect

1.50. a — Accidents

1.51. b — Enabling

1.52. c — Enabling

1.53. c — Low level of father education

1.54. d — Promoting disease prevention

1.55. a — Birth weight

1.56. b — Social problems

1.57. a — Infections

1.58. a — Recognizes that the family is the constant in a child's life

1.59. a — Respect

1.60. a — Acute lower respiratory tract infections

1.N1. b — In the first year of life per 1000 live births
IMR = deaths of infants under 1 year per 1000 live births; neonatal mortality covers the first 28 days.

1.N2. b — Birth weight
Low birth weight is the strongest predictor of infant death.

1.N3. a, b, c, d — Poverty; Violence; School failure; Adjustment to divorce
The new morbidity refers to social, behavioral and educational problems; measles is a classic infectious disease.

1.N4. b — Enabling
Enabling gives families opportunities to display existing abilities and learn new ones to meet the child's needs.

1.N5. b — Immunization and health teaching
Primary prevention keeps disease from occurring (immunization, teaching); the others are secondary/tertiary.

1.N6. c — Injuries (accidents)
Unintentional injuries are the leading cause of death after the first year of life.

2.1. c — Conditions require in-patient care

2.2. a — Fidelity

2.3. b — Veracity

2.4. b — Autonomy

2.5. a — It is a right of hospitalized children

2.6. b — Autonomy

2.7. b — Autonomy

2.8. b — Veracity

2.9. a — Confidentiality

2.10. b — It is a right of hospitalized children

2.11. d — Ethical decision making

2.12. b — Autonomy

2.13. a — Do no harm

2.14. b — Veracity

2.15. d — Accommodate the mother to stay with the child

2.16. d — Autonomy

2.17. c — Justice

2.18. a — Administering the least invasive treatment to avoid unnecessary harm

2.19. d — Beneficence

2.20. d — Autonomy

2.21. c — Non-maleficence

2.22. d — Prioritizing care for a critically ill child based on their need
بالأحمر كتصحيح — العدل (d) واتعلّم على (b) الطالب في الصورة اختار توزيع الرعاية حسب الاحتياج، مش وقت متساوي للكل =.

2.23. b — Beneficence

2.24. b — Often referred to as fairness

2.25. b — Veracity

2.26. b — Conditions require in-patient care

2.27. c — Fidelity

2.28. a — Veracity

2.29. d — Conditions require in-patient care

2.30. a — Fidelity

2.31. a — Fidelity

2.32. c — Conditions require in-patient care

2.33. b — Veracity

2.34. b — Beneficence

2.35. a — It is a right of hospitalized children

2.36. b — Autonomy.

2.37. b — Veracity

2.38. a — Confidentiality

2.39. b — Conditions require in-patient care

2.40. c — Fidelity

2.41. a — Veracity

2.42. a — Fidelity

2.43. a — Do no harm

2.44. b — Often referred to as fairness.

2.45. d — Beneficence.

2.46. b — Veracity

2.47. d — Accommodate the mother to stay with child

2.48. b — Beneficence

2.49. a — Autonomy

2.50. b — Veracity

2.51. a — Confidentiality

2.52. b — It is a right of hospitalized children

2.53. b — Beneficence

2.54. a — Fidelity

2.55. b — Veracity

2.56. a — Do no harm

2.57. a — Fidelity

2.N1. b — Veracity
Veracity is the duty to tell the truth; children should be told honestly what to expect.

2.N2. b — Autonomy
Autonomy = the right of self-determination; children participate in decisions according to their developmental level.

2.N3. a, b, c, d — Autonomy; Beneficence; Non-maleficence; Justice
The four primary principles are autonomy, beneficence, non-maleficence and justice; veracity, fidelity and confidentiality are secondary.

2.N4. a — Non-maleficence
Non-maleficence = do no harm, including reducing pain and emotional trauma during procedures.

2.N5. b — Discussing a child's diagnosis with a nurse friend from another hospital
Sharing information with people not involved in care violates confidentiality; mandatory abuse reporting is a legal duty.

2.N6. b — Distributing care fairly according to the child's need
Justice means fairness — care allocated by need, not by social status or ability to pay.

3.1. c — Protest

3.2. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.3. c — Protest

3.4. c — Concerned with appearance

3.5. c — 6 months to 6 years

3.6. c — School-agers

3.7. a — Detached from both parents

3.8. c — Loss of control and privacy

3.9. a — Screams and hits the nurse

3.10. a — Separation anxiety

3.11. c — Establish a routine similar to the child's home

3.12. d — Encouraged and permitted freely

3.13. a — Histamine

3.14. c — Preschoolers

3.15. c — Separation from parents

3.16. a — Stops and starts again

3.17. b — Loss of control and privacy

3.18. d — Inactive, withdrawn, depressed and sad

3.19. a — Cries, screams and clings to parents

3.20. d — Separation from parents and siblings

3.21. b — Regresses to earlier behavior (e.g. thumb sucking, bed wetting)

3.22. c — 6 months to 6 years

3.23. c — Protest

3.24. c — Protest

3.25. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.26. d — Depressed, regressive behavior, withdrawal

3.27. d — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.28. c — Allow mother's rooming-in

3.29. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.30. c — Protest

3.31. c — Allow mother's rooming in.

3.32. a — It is a right of hospitalized children

3.33. c — Concerned with appearance.

3.34. c — 6 months to 6 years

3.35. b — Loss of control and privacy
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.36. a — Separation anxiety

3.37. c — Preschoolers

3.38. c — Separation from parents

3.39. d — Accommodate the mother to stay with child

3.40. c — Protest.

3.41. c — Establish a routine similar to child's home.

3.42. c — Conditions require in-patient care

3.43. d — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3.44. d — Detached from both parents.

3.N1. c — Protest
Protest: crying, screaming, searching for the parent and rejecting strangers.

3.N2. b — Altered body image and loss of peer contact
Adolescents worry most about appearance/body image, privacy and separation from peers.

3.N3. a, b, c — Encourage rooming-in of the parent; Keep the home routine (bedtime rituals); Bring a favorite toy or blanket from home
Parental presence, consistent routines and transitional objects reduce stress; consistent caregivers are preferred.

3.N4. c — Detachment (denial)
Detachment is the last stage — superficial adjustment with loss of interest in the parent; it is the most harmful stage.

3.N5. b — Preschoolers
Preschoolers often regress to earlier behaviors (bed-wetting, thumb sucking) as a coping mechanism.

3.N6. b — They fear losing control of their bodies and roles and may react with depression or hostility
School-agers strive for independence and productivity; loss of control and privacy is very stressful for them.

4.1. a — Evaluate hydration status

4.2. c — Imperforate anus
و imperforate anus: «في صورة الامتحان مكتوب «الإجابتين صح الاتنين بيسببوا عدم نزول العقي aganglionic megacolon (Hirschsprung).

4.3. c — Diaphragmatic hernia

4.4. c — Aspiration

4.5. c — Prone

4.6. b — Increase in saliva

4.7. a — Protect the operative site from infection and injury

4.8. b — Water-soluble vitamin containing

4.9. a — Burp the baby frequently

4.10. b — Prone

4.11. c — Prevent aspiration

4.12. b — Scaphoid abdomen and diminished breath sounds

4.13. b — Projectile vomiting

4.14. a — Currant jelly stool

4.15. c — Constipation and passage of ribbon-like stool

4.16. c — Water-soluble vitamins

4.17. a — Protect the operative site from infection or injury

4.18. b — Diminished breathing sounds

4.19. d — Measure abdominal girth periodically

4.20. a — Constipation and passage of ribbon-like stool

4.21. b — Evaluates hydration status

4.22. d — Protects operative sites from infection and injury

4.23. b — Supine

4.24. c — Failure to pass meconium immediately after birth

4.25. c — Evaluate hydration status

4.26. d — Aspiration

4.27. d — Diaphragmatic hernia

4.28. d — Aspiration

4.29. a — Ribbon-like, foul-smelling stools

4.30. c — Aspiration

4.31. a — Evaluate hydration status

4.32. c — Imperforate anus
و imperforate anus: «في صورة الامتحان مكتوب «الإجابتين صح الاتنين بيسببوا عدم نزول العقي aganglionic megacolon (Hirschsprung).

4.33. b — Imperforate anus
و imperforate anus: «في صورة الامتحان مكتوب «الإجابتين صح الاتنين بيسببوا عدم نزول العقي aganglionic megacolon (Hirschsprung).

4.34. c — Diaphragmatic hernia

4.35. c — Hypertrophic pyloric stenosis

4.36. a — Protect operative site from infection and injury.

4.37. b — Water soluble vitamin containing.

4.38. b — Prone

4.39. c — Prevent aspiration

4.40. c — Evaluate hydration status

4.41. a — Aspiration

4.42. d — Diaphragmatic hernia

4.43. a — Rkibbon like, foul-smelling stools

4.44. c — Imperforated anus
و imperforate anus: «في صورة الامتحان مكتوب «الإجابتين صح الاتنين بيسببوا عدم نزول العقي aganglionic megacolon (Hirschsprung).

4.45. c — Hypertrophic pyloric stenosis

4.46. b — Increase in saliva.

4.47. b — Supine.

4.48. c — Failure to pass meconium immediately after birth.

4.49. a — Constipation and passage of ribbon- like stool.

4.50. a — Currant jelly stool

4.51. a — Burp the baby frequently.

4.N1. b — Esophageal atresia with TEF
The 3 C's (coughing, choking, cyanosis) and excessive drooling are classic for EA/TEF; feeding must stop immediately.

4.N2. b — Head elevated 30–40° (prone or supine) with continuous suction of the pouch
Elevating the head and continuous suction reduce reflux and aspiration of secretions.

4.N3. a, b, c, e — Projectile non-bilious vomiting; Olive-shaped mass in the right upper quadrant; Visible peristaltic waves; Hypochloremic metabolic alkalosis
Currant-jelly stool belongs to intussusception; vomiting HCl leads to hypochloremic metabolic alkalosis.

4.N4. b — Elbow restraints and a Logan bow
Elbow restraints keep hands away from the lip and a Logan bow reduces tension; position supine or on the side, never prone, after lip repair.

4.N5. b — Chronic constipation with ribbon-like, foul-smelling stools and abdominal distension
Absence of ganglion cells → no peristalsis in the affected segment → obstruction, distension and ribbon-like stools.

4.N6. b — With head elevated and on the affected side
Lying on the affected side with head elevated lets the unaffected lung expand; avoid bag-mask ventilation that inflates the bowel.

5.1. a — True

5.2. c — Moderately low

5.3. d — Excessive lanugo hair

5.4. d — Full term, small for gestational age

5.5. b — Stop if there is evidence of overstimulation

5.6. c — Lanugo

5.7. d — Meticulous hand washing of all persons in contact

5.8. a — Minimal handling during care

5.9. b — Minimal handling during care

5.10. d — Dry the infant in a warm towel

5.11. a — Respiratory distress syndrome

5.12. a — Minimize neonate handling

5.13. a — Low birth weight

5.14. b — Lack of sole creases on his feet
preterm وده غلط — الـ scant lanugo في صورة الكويز الطالب اختار lack of sole creases (زي بنك، والإجابة الصح lanugo كتير (excessive) عنده الأسئلة).

5.15. d — Less than 1000 gm

5.16. b — Promote developmental care

5.17. a — Moderately low

5.18. d — Excessive lanugo hair

5.19. b — Turning on the overhead radiant warmer
الفقد بالإشعاع بيتمنع بتدفئة الأسطح اللي حوالين الطفل (radiant warmer). في صورة واحدة الطالب اختار closing doors وده بيمنع الـ convection.

5.20. a — True

5.21. c — Moderately low

5.22. a — True

5.23. b — Turning on the overhead radiant warmer
الفقد بالإشعاع بيتمنع بتدفئة الأسطح اللي حوالين الطفل (radiant warmer). في صورة واحدة الطالب اختار closing doors وده بيمنع الـ convection.

5.24. d — Meticulous hand washing of all persons in contact.

5.25. a — Minimal handling during care.

5.26. a — Moderately low

5.27. d — Excessive lanugo hair

5.28. c — Drying the infant with a warm blanket,

5.29. a — True

5.30. b — Turning on the overhead radiant warmer

5.31. a — Respiratory distress syndrome

5.32. d — A full term a small for gestational age.

5.33. c — Lanugo.

5.34. d — Less than 1000 gm.

5.35. b — Promote developmental care.

5.36. a — Low birth weight.

5.37. b — Lack of sole creases on his feet.
preterm وده غلط — الـ scant lanugo في صورة الكويز الطالب اختار lack of sole creases (زي بنك، والإجابة الصح lanugo كتير (excessive) عنده الأسئلة).

5.38. d — Dry the infant in a warm towel

5.39. a — Respiratory distress syndrome

5.40. b — Low birth weight

5.N1. b — Term, SGA (low birth weight)
Term = 37–42 weeks; weight below the 10th percentile/under 2500 g makes the baby SGA and LBW.

5.N2. b — Abundant lanugo and few sole creases
Preterm infants have abundant lanugo, thin skin, few sole creases, weak reflexes and soft ear cartilage.

5.N3. a, b, c, e — Poor temperature regulation; Respiratory difficulties and apnea; Susceptibility to infection; Immature liver (jaundice, bleeding)
Premature babies have poor heat regulation, immature lungs, weak immunity and immature liver/kidneys.

5.N4. b — Keep the baby away from drafts and keep incubator portholes closed
Convection = loss to moving air; evaporation is prevented by drying, conduction by warming surfaces, radiation by warming surroundings.

5.N5. b — Less than 1500 g
LBW <2500 g, VLBW <1500 g, ELBW <1000 g.

5.N6. b — Clustering care, minimal handling and reducing noise
Developmental care reduces stress and conserves energy: cluster care, dim lights, low noise, nesting and parental involvement.

6.1. c — Pretend play with other children

6.2. a — It helps to improve their physical development

6.3. a — Parallel
سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. (وصف الـ meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

6.4. d — Day dreaming
سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. (وصف الـ meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

6.5. b — Solitary
سؤال توصيل في الامتحان الأصلي: solitary (لعب إجابة الطالب. الرضيع بجسمه = sense-pleasure play، وبيكون لوحده).

6.6. b — False

6.7. a — Beside other children, but not with them

6.8. b — Suitable to age and health

6.9. b — Parallel play

6.10. a — Suitable for age and health

6.11. c — Sociodramatic

6.12. d — It helps to improve their physical growth

6.13. c — Helps to improve their physical growth

6.14. d — Release tension

6.15. a — Provides diversion and relaxation

6.16. d — Preschool

6.17. b — Release tension

6.18. d — Playing alongside other children
ملاحظة: الصفحة الأولى من الكويز ده كانت مقصوصة — سؤال 6 اختياراته مش ظاهرة كاملة فاتشال (play & cognitive development).

6.19. c — Parallel

6.20. a — Dramatic play with other children

6.21. c — Therapeutic

6.22. b — Socio-dramatic play

6.23. d — Parallel

6.24. c — Sociodramatic

6.25. a — True

6.26. a — Pretend play with other children

6.27. a — Therapeutic

6.28. b — Ahmed plays with a car beside Heba, who plays with a doll

6.29. c — Pretend play with other children

6.30. d — Ahmed plays with a car beside Heba, who plays with a doll

6.31. b — Toy is suitable to developmental age

6.32. d — Therapeutic

6.33. a — True

6.34. b — Parallel

6.35. b — False
حسب الملخص: الـ constructive play خاص بسن المدرسة (early school years) مش preschool → False.

6.36. a — Provides diversion and relaxation

6.37. a — Suitable for age and health

6.N1. b — Parallel play
Parallel play — independent play beside, not with, others — is typical of toddlers.

6.N2. b — Dramatic/pretend play such as playing 'doctor'
Preschoolers enjoy sociodramatic/imaginative play and imitating adult roles.

6.N3. a, b, c, d — Provides diversion and relaxation; Helps the child feel more secure; Lessens stress of separation; Allows expression of feelings
Hospital play reduces stress, gives a sense of control and helps the child express and master feelings.

6.N4. b — Therapeutic
Therapeutic play is used to achieve treatment goals, e.g., deep breathing.

6.N5. b — Large soft blocks with no detachable parts
Toys must suit the age and be free of small/removable parts that can be aspirated.

6.N6. c — School-age children
Constructive play (building, solving problems, creating products) is characteristic of the early school years.

7.1. a — Curiosity

7.2. d — Active preventive strategy

7.3. a — Home

7.4. c — Suffocation

7.5. a — Boys higher than girls

7.6. c — Drowning

7.7. d — Falling
أشيع (falls) الملف مفيهوش إجابة معلّمة — إجابة مقترحة: الوقوع حوادث البيت عند الرضّع اللي بيبدأوا يتقلّبوا ويزحفوا.

7.8. d — Furniture is to be away from the window

7.9. d — Active preventive strategy

7.10. b, c, e — A 2-month-old infant who sleeps in the bed with his mother; A 1-month-old infant who prefers to sleep on his stomach; A newborn whose mother smokes 2 packs/day

7.11. c — Unintentional injuries

7.12. c, d, e — Know the number of the nearest poison control center; Caution the child against eating nonedible items such as plants; Discard empty poison containers

7.13. a, b, e — Male gender; Low Apgar scores; Low birth weight

7.14. a — Aspiration/poisoning

7.15. b — Injury prevention and parent education should be handled on a growth and developmental basis.

7.16. d — Falling
أشيع (falls) الملف مفيهوش إجابة معلّمة — إجابة مقترحة: الوقوع حوادث البيت عند الرضّع اللي بيبدأوا يتقلّبوا ويزحفوا.

7.17. d — Furnitures are to be away from the window
إجابة البنك (furniture away from window) لاحظ إن الـ plugs صحيحة؛ لازم تتغطى بس مش 'at low level'.

7.18. c — Suffocation

7.19. a — Home

7.20. a — Boys higher than girls

7.N1. b — Aspiration of small objects and poisoning
Crawling infants explore with their mouths; aspiration of small objects and ingestion of household products are the classic hazards of this stage.

7.N2. c — Parent stores medicines and cleaners in a locked high cabinet
Active strategies need a repeated action by the person (parent locking cabinets); passive strategies protect automatically without individual effort.

7.N3. a, b, d — Place the infant on the back to sleep; Use a firm mattress without pillows or soft toys; Avoid smoking around the infant
Back-to-sleep, firm bare crib and a smoke-free environment lower SIDS risk; bed-sharing and overheating increase it.

7.N4. b — Curiosity and increasing mobility with poor awareness of danger
Toddlers are curious, mobile and impulsive but cannot yet recognize danger — the key developmental risk factor.

7.N5. b — Agent
The agent (vector) is the energy source that causes the injury — here thermal energy from the hot liquid; the child is the host.

7.N6. b — Drowning and animal bites occur mainly among rural children
Egyptian data: boys aged 2–6 years have more injuries; drowning, animal bites and sunstroke occur mostly in rural children.

8.1. a — Breastfeed the neonate frequently

8.2. b — High in antibodies, protein, vitamins, and minerals

8.3. d — Antibodies that the neonate lacks

8.4. c — Give only breast milk for the first 6 months of age

8.5. a — Colostrum is high in antibodies, protein, vitamins, and minerals

8.6. c — Breastfeed the neonate frequently

8.7. b — Excessive sweating

8.8. a — Early initiation and for six months

8.9. c — Latch on well to the breast

8.N1. b — Rich in antibodies (IgA) and acts as the baby's first immunization
Colostrum is rich in protein and secretory IgA, gives passive immunity and has a laxative effect that helps pass meconium.

8.N2. b — Oxytocin
Oxytocin from the posterior pituitary causes milk ejection; it is affected by emotions (worry, pain and stress reduce it).

8.N3. a, b, c — 6 or more wet diapers per day; Steady weight gain; Calm sleeping after feeds
Adequate urine output, weight gain and contentment after feeds indicate successful breastfeeding.

8.N4. b — Vitamin D
Breast milk contains all vitamins in adequate amounts except vitamin D.

8.N5. a — Within 1 hour of birth
Step 4 of the 10 steps: help mothers start breastfeeding within one hour of birth.

8.N6. c — 6 months while breastfeeding continues
Breast milk alone covers needs for the first 6 months; then complementary foods are added without replacing breast milk.

9.1. d — Cover the sac with a moist and sterile dressing

9.2. a — Prone

9.3. c — Absence of the cerebral hemispheres
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

9.4. b — A protrusion of nerve tissue and meninges through a skull defect
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

9.5. a — Incomplete fusion of the vertebrae that may be signaled by an overlying tuft of hair
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

9.6. d — A sac of meninges and CSF protruding through a vertebral defect without nerve tissue
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد. meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

9.7. b — Measuring head circumference

9.8. c — Infection in CSF is indicated

9.9. b — False

9.10. d — Aqueduct stenosis

9.11. a — Indicative of CSF infection

9.12. c — Apply a sterile, moist, non-adherent dressing
الإجابة الصح حسب بنك الأسئلة — (a) ورقة الطالب معلّم عليها والملخص: الغيار المعقّم المبلول غير اللاصق.

9.13. d — Measure head circumference

9.14. d — Cover the defect with a saline-soaked dressing

9.15. c — Monitor for increased temperature

9.16. b — Measure head circumference

9.17. b — On the non-operative site

9.18. c — Prone

9.19. d — Cover the sac with a moist and sterile dressing

9.20. a — Prone

9.21. c — Prone

9.22. d — Obstruction

9.23. b — Myelomeningocele

9.24. b — Babies with a myelomeningocele are at risk for hydrocephalus, which shows up as an increase in head size

9.25. b — UTI

9.26. b, c — Daily head circumference measurements; MRI scan

9.27. d — Place the child in the prone position with a sterile moist dressing over the defect. begin IV fluids to prevent dehydrations

9.28. c — Assist the child to change positions to avoid skin breakdown

9.29. a — Take a multivtaim with folic acid daily

9.30. c — Risk of infection

9.31. c — Place a wet dressing on the sac

9.32. b — Measure head circumference

9.33. b — Potential for infection r/t the physical defect

9.34. b — Tell the parent this might mean the baby has increased icp

9.35. d — Need lifelong management of urinary, orthopedic and neurological problems

9.36. c — Bulging fontanel and downwardly rotated eyes

9.37. a — Axillary temp of 102.2 (fever)

9.38. b — Catheterize the infants bladder

9.39. b — Its done to decrease the possibility of infection and further cord damage

9.40. c — Apply sterile, moist, non-adherent dressing

9.41. b — Flat in a crib

9.42. d — Full fontanels, poor feeding, rapid head growth

9.43. d — Measure head circumference

9.44. a — Indicted of CSF infection

9.45. d — Cover sac with a moist and sterile dressing

9.46. a — Prone

9.47. d — Aqueduct stenosis

9.N1. b — Covering the sac with a sterile, moist, non-adherent dressing
A moist sterile dressing prevents drying, rupture and infection of the sac; position prone.

9.N2. b — Rapidly increasing head circumference with bulging fontanel
Open sutures allow the head to enlarge; bulging fontanel, sunset eyes and irritability follow.

9.N3. a, b, c, e — Vomiting and irritability; Bulging fontanel; Fever; High-pitched cry
Shunt obstruction raises ICP (vomiting, bulging fontanel, high-pitched cry); fever and redness suggest infection.

9.N4. b — Folic acid 0.4 mg daily before conception and in early pregnancy
Periconceptional folic acid markedly reduces neural tube defects.

9.N5. b — A tuft of hair or dimple over the lumbosacral spine
The defect is hidden; skin signs such as a hair tuft, dimple or hemangioma may be present.

9.N6. b — Flat, on the non-operative side
Lying flat on the non-operative side avoids pressure on the valve and prevents too-rapid CSF drainage.

10.1. a — High risk for injury

10.2. c — Give penicillin until 21 years old

10.3. d — Seek immediate treatment for throat infection
ورقة الطالب معلّم عليها corticosteroid الوقاية؛ وده علاج مش وقاية الأولية = علاج التهاب الحلق البكتيري بسرعة.

10.4. c — Increased sedimentation rate

10.5. c — 2–3 weeks
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.6. b — Knees, ankles, elbows and wrists
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.7. b — Pain that moves from joint to joint (migratory)
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.8. d — Headache
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.9. b — Hot or tender joints
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.10. b — Musculoskeletal, cardiovascular and nervous
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.11. b — Fever and joint pain
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.12. b — Erythema marginatum
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.13. c — Penicillin
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.14. d — Nausea
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

10.15. c — Increased sedimentation rate

10.16. c — Give penicillin until 21 years old

10.17. a — High risk for injury

10.18. b — False

10.19. c — Heart

10.20. c — Joint pain and fever

10.21. b — False

10.22. a — Acetaminophen

10.23. b — Oral salicylates

10.24. c — Changes in vision

10.25. c — Streptococcus

10.26. c — The mother's history of type-2 diabetes

10.27. a, c — Elevated C-reactive protein; Elevated WBC count

10.28. b, c — Positive strep throat culture, arthritis, chorea; Recent scarlet fever, carditis, fever, arthralgia

10.29. d — Streptococcus pyogenes (Group A Streptococcus)

10.30. c — The cross-reactivity between bacterial antigens and host tissues.

10.31. c — Elevated C-reactive protein

10.32. c — Corticosteroid

10.33. d — Increasing fluid and salt intake

10.34. c — Chorea

10.35. d — Genetic markers associated with the immune response

10.36. c — Carditis

10.37. c — Meningitis

10.38. b — Arthritis
إجابة البنك (Arthritis) الـ غير دقيقة: polyarthritis معيار major، والمعيار minor هو الـ arthralgia. الأقرب هي لو الاختيارات دي جت، b.

10.39. c — Throat culture

10.40. d — Antibiotics targeting Gram-negative bacteria

10.41. b — Musculoskeletal system
إجابة البنك: musculoskeletal (المفاصل أشيع عرض). لكن أخطر وأدوم تأثير على القلب.

10.42. d — All of the above

10.43. b — Children ages 5-15

10.44. c — Early diagnosis and treatment of streptococcal infections

10.45. a — Heart

10.46. b — Rheumatic heart disease

10.N1. b — Group A beta-hemolytic streptococcus
Molecular mimicry between streptococcal antigens and host tissues (heart, joints, brain, skin) causes RF.

10.N2. a, b, c, d, e — Carditis; Polyarthritis; Chorea; Erythema marginatum; Subcutaneous nodules
The 5 major criteria are carditis, polyarthritis, chorea, erythema marginatum and subcutaneous nodules; elevated ESR/CRP is minor.

10.N3. b — Benzathine penicillin IM every 3–4 weeks for years
Long-acting penicillin prevents recurrence; duration depends on carditis (often until age 21 or longer).

10.N4. b — Risk for injury
Involuntary jerky movements put the child at risk of falls and injury — provide a safe, quiet environment.

10.N5. a — New heart murmur and tachycardia out of proportion to fever
Carditis presents with new murmurs, tachycardia, cardiomegaly or pericardial rub; it is the only manifestation that can cause permanent damage.

10.N6. b — Migratory polyarthritis of large joints, non-deforming
RF arthritis migrates between large joints (knees, ankles, elbows, wrists) and heals without deformity.

11.1. d — Tea-colored urine

11.2. b — Weight

11.3. b — There is blood in the child's urine that causes it to be tea-colored

11.4. a — Body image

11.5. a — Keep the child away from sick playmates

11.6. d — Poor appetite, edema, proteinuria

11.7. c — Keep the child away from sick playmates
واتصحّح غلط ✗ — الإجابة (decrease sodium) في الصورة الطالب اختار الصح: keep the child away from sick playmates (العدوى أشهر سبب للانتكاس).

11.8. a — Increased urine output

11.9. c — Massive proteinuria
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.10. c — Minimal change disease
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.11. d — Inflammation is a main feature of nephritis
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.12. b — Weak immune system (infection)
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.13. d — Kidney biopsy
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.14. b — Autoimmune diseases
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.15. b — Atorvastatin
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.16. c — Hypertension
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.17. c — Low-protein, low-sodium diet
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.18. d — Reduce inflammation
إجابة مقترحة (المصدر مفيهوش إجابة رسمية).

11.19. d — Tea colored urine

11.20. a — Increased urine output

11.21. a — True

11.22. b — False

11.23. a — 2:1

11.24. e — All of the above

11.25. e — Antihistaminic

11.26. b — There is blood in the child's urine that causes it to be tea colored

11.27. a — Restrict protein intake
الأدق حسب الملخص: تقييد الصوديوم والبروتين يتقيّد بس لو في oliguria ، والسوائل والبوتاسيوم في حالة الـ azotemia. إجابة البنك: restrict protein.

11.28. b, d, e — Monitor intake and output hourly; Assess color of urine after every void; Weigh the patient daily on a standing scale

11.29. b — Potassium-rich foods

11.30. c — Morning

11.31. a — A 3 year old male who has a positive ASO titer.

11.32. a, b, d — Hypotension; Increased glomerular filtration rate; Massive proteinuria

11.33. b — False
APSGN is a NEPHRITIC syndrome (RBCs + protein leak); nephrotic syndrome = protein only.

11.34. d — "This condition is not caused by the streptococcal bacteria attacking the glomerulus, but by the immune system's response to the bacteria by creating an antigen-antibody complex which inflames the glomerulus."

11.35. a — Red blood cells in urine

11.36. c — A streptococcal infection

11.37. d — Acute post-streptococcal glomerulonephritis

11.38. a — Smoky colored urine

11.39. a — Recent illness such as strep throat

11.40. d — Encourage the child to take all the antibiotics if diagnosed with strep throat.

11.41. b — Dark brown urine

11.42. b — Take the child's blood pressure and report the findings to the nurse while the nurse is still on the phone.

11.43. b — Eyes

11.44. d — Streptococcal infection

11.45. a — Blood

11.46. c — Dark and foamy

11.47. e — All the above

11.48. e — Both C and D

11.49. d — All the above statements are true

11.50. e — All of the above

11.51. a — Corticosteroids

11.52. d — Hypertension

11.53. d — Fowler's

11.54. d — Weighing on the same scale each day

11.55. c — Eyes and ankles

11.56. d — Measure the abdominal girth daily.

11.57. c — Selective albuminuria

11.58. c — Blood in the urine

11.59. c — Edema

11.60. b — Checking his urine for protein

11.61. a — Risk for infection

11.62. d — A 7 year old male recently diagnosed with Goodpasture's Syndrome.

11.63. f — B and D

11.64. a, e, f — Slight proteinuria; Tea-colored urine; Hypertension

11.65. b — Redden area on the patient's left leg that is swollen and warm

11.66. b — Low-fat

11.67. a — Keep the child away from sick playmates

11.68. b — Weight

11.69. d — Por appetite, edema, proteinuria

11.70. c — Keep the child away from sick playmates

11.71. a — Body image

11.72. e — All of the above

11.73. b — False

11.74. d — Analgesic

11.75. a — True

11.N1. b — Massive proteinuria, hypoalbuminemia, edema
Nephrotic syndrome = massive proteinuria, hypoalbuminemia, generalized edema (± hyperlipidemia).

11.N2. b — Hematuria
Immune-complex glomerular inflammation lets RBCs leak into the urine, giving a cola/tea color.

11.N3. a, b, c, e — Weigh daily on the same scale; Measure abdominal girth daily; Protect from infection; Provide meticulous skin care
Monitor fluid status, protect edematous skin and prevent infection (steroids + protein loss); sodium is restricted during edema.

11.N4. b — Group A beta-hemolytic streptococcal throat or skin infection
Post-streptococcal GN appears 1–3 weeks after a streptococcal pharyngitis or impetigo.

11.N5. b — Increased urine output (diuresis)
Diuresis is the first sign of recovery, followed by reduction in edema and blood pressure.

11.N6. b — Prednisone (corticosteroids)
Most children with minimal change disease respond to corticosteroids.

12.1. a — Rapid respiration and nasal flaring
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

12.2. b — Intercostal and subcostal retractions
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

12.3. c — Expiratory grunting
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

12.4. d — Cyanosis and altered conscious level
وصف الـ) سؤال توصيل في الامتحان الأصلي، اتحوّل لاختيار من متعدد meningocele مكتوب من الملخص لأنه كان مقصوص في الصورة).

12.5. d — Sternum retraction

12.6. d — Surfactant replacement

12.7. c — Frequent suctioning

12.8. b — Prevent lung collapse

12.9. b — Tachypnea and retractions

12.10. c — Impaired gas exchange

12.11. c — Impaired gas exchange

12.12. b — False
الـ grunting علامة متأخرة (Grade III) — المبكرة tachypnea.

12.13. b — False
grunting علامة متأخرة (Grade III) — المبكرة tachypnea.

12.14. b — Prevent the lung collapse

12.15. c — Frequently suctioning

12.16. b — Tachypnea and retractions

12.17. c — Decrease surface tension of alveoli

12.18. b — False
grunting علامة متأخرة (Grade III) — المبكرة tachypnea.

12.19. c — 24 – 34

12.20. c — Impaired gas exchange

12.21. d — Sternum retraction

12.22. c — Surfactant

12.23. d — Surfactant replacement

12.24. a — True

12.25. c — Limited birth

12.26. b — Decreased alveolar ventilation

12.27. b — 2

12.28. a — True

12.29. c — Infasurf
Infasurf = calfactant (من رئة العجل)؛ Survanta من رئة البقر، Curosurf من رئة الخنزير.

12.30. d — All of the above

12.31. b — False

12.32. b — False

12.33. d — Respiratory distress syndrome

12.34. c — 24 weeks

12.35. a — True

12.36. d — 32 weeks

12.37. c — Female gender

12.38. a — True

12.39. c — 35 weeks

12.40. b — Increased PH

12.41. c — Hypertension

12.42. a — True

12.43. d — Metabolic acidosis

12.44. b — Grade II

12.45. a — Occurs in ~50% of premature births with a weight of <1500 g

12.46. a — The baby whose mother has diabetes mellitus.

12.47. c — 4 & respiratory distress

12.48. d — Give beta blockers

12.49. b — Assist the neonatologist during the intubation procedure.

12.50. b — Oxygen saturation 96%

12.51. d — All of the above

12.52. b — 50-80 mmHg

12.53. d — Semi-Fowler's

12.54. c — Decrease surface tension of alveoli

12.N1. b — Surfactant deficiency
Lack of surfactant increases alveolar surface tension, causing collapse (atelectasis).

12.N2. b — Tachypnea
Tachypnea and nasal flaring appear early; grunting and cyanosis are later grades.

12.N3. a, b, c, d — Prematurity; Maternal diabetes; Cold stress; Perinatal asphyxia
Male gender is the risk factor.

12.N4. b — 24–34 weeks
Steroids accelerate fetal lung maturity and surfactant production.

12.N5. b — Endotracheal tube
The nurse assists intubation; avoid suctioning immediately after dosing.

12.N6. b — Respiratory and metabolic acidosis
Hypoxia and CO2 retention cause respiratory and metabolic (lactic) acidosis.