PEDIATRIC NURSING (THEORY) · LECTURE 3 · QUESTION BANK

Hospitalization of Children

دخول الأطفال المستشفى — بنك الأسئلة

44	6	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

ميدترم 2020 · ميدترم 2021 · ميدترم 2022 · ميدترم 2024 (ترم أول 2023-2024) · كويز 1 – 2021 · كويز 1 – 2022 · كويز 1 – 2022 (نسخة بلاك بورد) · كويز 1 – 2023/2024 (شعبة 4) · كويز 1 إعادة – ترم تاني 2023/2024 · أسئلة سابقة · تجميعة صور ميدترمات سابقة · كويز 1 أطفال نظري (Microsoft Forms) · كويز 1 – ترم تاني 2025

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 ميدترم 2020

A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر

a. Despair
b. Detachment
c. Protest
d. Regression

2 ميدترم 2020

Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر

a. Concerned with appearance
b. Separation from peer group
c. Stranger anxiety
d. Fear of pain

3 ميدترم 2021

A hospitalized toddler, for two days, cries because he was left alone. The nurse realizes this behavior is typical of:

a. Regression
b. Despair
c. Protest
d. Stranger anxiety

4 ميدترم 2022

Which of the following represents the major stressor of hospitalization for adolescents? تكرر في 2 مصادر

a. Separation anxiety
b. Fear of pain
c. Concerned with appearance
d. Loss of control

5 ميدترم 2024 (ترم أول 2023-2024)

Separation anxiety is more drastic among children aged: تكرر في 2 مصادر

a. 2 months to 6 months
b. 6 months to 12 months
c. 6 months to 6 years
d. 6 months to 12 years

6 The pediatric nurse should realize that guided imagery is successfully used to decrease pain of: ميدترم 2024 (ترم أول 2023-2024)

a. Toddlers
b. Preschoolers
c. School-agers
d. Adolescents

7 A child hospitalized for a chronic illness showed symptoms of depression but later started interacting with others. What does the nurse infer? The child is ..... كويز 1 – 2021

a. Detached from both parents
b. Getting used to the surroundings
c. Showing improved social skills
d. Content with the care provided

8 Which of the following stressors would the nurse recognize as having the greatest impact on the school-age hospitalized child? تكرر في 2 مصادر كويز 1 – 2022

a. Separation from family
b. Fear of death
c. Loss of control and privacy
d. Concerned with appearance

9 Which of the following child's behaviors leads the nurse to believe that the child is exhibiting the stage of protest? كويز 1 – 2022

a. Screams and hits the nurse
b. Withdrawn from others
c. Returns to the habit of bed-wetting
d. Sits in a corner with a toy

10 Which of the following represents the major stressor of hospitalization for children from middle infancy throughout the preschool years? تكرر في 2 مصادر كويز 1 – 2022 (نسخة بلاك بورد)

a. Separation anxiety
b. Fear of bodily injury
c. Fear of pain
d. Loss of control

11 Which of the following should the nurse do to prevent separation anxiety in a hospitalized toddler? تكرر في 2 مصادر كويز 1 – 2022 (نسخة بلاك بورد)

a. Assume the parental role when parents are not present
b. Limit visits by siblings and relatives
c. Establish a routine similar to the child's home
d. Rotate nursing staff so the child becomes comfortable

12 Attendance of parents during painful procedures on their toddlers should be: كويز 1 – 2023/2024 (شعبة 4)

a. Based on individual assessment of parents
b. Based on the type of procedure to be performed
c. Discouraged for the benefit of parents and child
d. Encouraged and permitted freely

13 Which of the following substances increases pain sensation? كويز 1 – 2023/2024 (شعبة 4)

a. Histamine
b. Endorphins
c. Enkephalins
d. Dynorphins

14 Regression is a reaction to hospitalization; ..... are at high risk for it. كويز 1 – 2023/2024 (شعبة 4)

a. Infants
b. Toddlers
c. Preschoolers
d. School-agers

15 Which one of the following stressors most affects a 2-year-old toddler? كويز 1 – 2023/2024 (شعبة 4)

a. Separation from peers
b. Fear of the dark
c. Separation from parents
d. Fear of death

16 The nurse best describes intermittent pain as pain that: كويز 1 إعادة – ترم تاني 2023/2024

a. Stops and starts again
b. Moves from one area to another
c. Covers a large area
d. Is sticking and intense

17 Which of the following stressors would the nurse recognize as having the greatest impact on the school-age hospitalized child? تكرر في 2 مصادر كويز 1 إعادة – ترم تاني 2023/2024

a. Fear of death
b. Loss of control and privacy
c. Concerned with appearance
d. Separation from family

18 In the despair phase of separation anxiety the child behaves as follows: كويز 1 إعادة – ترم تاني 2023/2024

a. Shows signs of interest in the surroundings
b. Avoids and rejects contact with strangers
c. Loud cry, scream and search for parent with eyes
d. Inactive, withdrawn, depressed and sad

19 Which of the following child's behaviors exhibits the stage of protest in separation anxiety? كويز 1 – ترم تاني 2025

a. Cries, screams and clings to parents
b. Withdrawn from others, sad and depressed
c. Interested in surroundings, active and communicative
d. Interacts with strangers and any other children

20 Which of the following is a primary stressor for a child during hospitalization? كويز 1 – ترم تاني 2025

a. Boredom from lack of activities
b. Unfamiliar hospital food
c. The noise of medical equipment
d. Separation from parents and siblings

21 A behavior observed in the despair phase of separation anxiety as a hospitalization reaction: كويز 1 أطفال نظري (Microsoft Forms)

a. Forms new but superficial relationships and appears happy
b. Regresses to earlier behavior (e.g. thumb sucking, bed wetting)
c. Shows increased interest in surroundings, interacts with strangers
d. Physically attacks strangers (kicks, bites, hits, pinches)

22 Which of the following is the period in which hospitalization reactions are obvious and risky? كويز 1 أطفال نظري (Microsoft Forms)

a. 6 months to 18 months
b. 3 months to 3 years
c. 6 months to 6 years
d. 3 months to 12 months

23 A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Detachment
b. Regression
c. Protest
d. Despair

24 A hospitalized toddler newly admitted screams because he was left alone. The nurse realizes this behavior is typical of: تكرر في 2 مصادر تجميعة صور ميدترمات سابقة

a. Despair
b. Detachment
c. Protest
d. Regression

25 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر تجميعة صور ميدترمات سابقة

a. Concerned with appearance
b. Separation from peer group
c. Stranger anxiety
d. Fear of pain

26 The characteristics of the despair phase of separation anxiety are: تجميعة صور ميدترمات سابقة

a. Rejects contact with strangers
b. The child shows signs of interest in the surroundings
c. Loud cry, scream and searches for parent with eyes
d. Depressed, regressive behavior, withdrawal

27 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر تجميعة صور ميدترمات سابقة

a. Fear of pain
b. Concerned with appearance
c. Stranger anxiety
d. Separation from peer group

28 What is the appropriate nursing intervention to minimize separation anxiety in a hospitalized child? تجميعة صور ميدترمات سابقة

a. Encourage contact with children the same age
b. Provide for privacy
c. Allow mother's rooming-in
d. Explain procedures and routines

29 Which of the following is considered more stressful for hospitalized school-age children? تكرر في 3 مصادر تجميعة صور ميدترمات سابقة

a. Stranger anxiety
b. Separation from peer group
c. Concerned with appearance
d. Fear of pain

30 A hospitalized toddler newly admitted and screams because he was left alone. The nurse realizes this behavior as typical of: أسئلة سابقة

a. Despair
b. Detachment
c. Protest
d. Regression

31 What is the appropriate nursing intervention minimize separation anxiety in a hospitalized child ? أسئلة سابقة

a. Encourage contact with children the same age
b. Provide for privacy.
c. Allow mother's rooming in.
d. Explain procedures and routines.

32 The pediatric hospitals must offer suitable accommodations for the parents because: أسئلة سابقة

a. It is a right of hospitalized children
b. Parents likes to spend time side them
c. Rooms are plenty in the hospitals.
d. Parents pay for the accommodations.

33 Which of the following represents the major stressor of hospitalization for adolescents ? أسئلة سابقة
تكرر في 2 مصادر

a. Separation anxiety.
b. Fear of pain.
c. Concerned with appearance.
d. Loss of control.

34 Separation anxiety is more drastic among children aged..... تكرر في 2 مصادر أسئلة سابقة

a. 2 months to 6 months
b. 6 months to 12 months
c. 6 months to 6 years
d. 6 months to 12 years

35 Which of the following is considered more stressful for hospitalized school age children? أسئلة سابقة
تكرر في 3 مصادر

a. Stranger anxiety
b. Loss of control and privacy
c. Considered with appearance
d. Fear of pain .

36 Which of the following represents the major stressor of hospitalization for children from middle infancy throughout the preschool years? تكرر في 2 مصادر أسئلة سابقة

a. Separation anxiety
b. Fear of bodily injury
c. Fear of pain
d. Loss of control

37 Regression is a reaction to hospitalization that is ..... are at high risk for it. أسئلة سابقة

a. Infants
b. Toddlers
c. Preschoolers
d. School agers

38 Which one of the following stressors most affect a 2-year-old toddler child? أسئلة سابقة

a. Separation from peers
b. Fear of the dark
c. Separation from parents
d. Fear of death

39 Which of the following reflects one of hospitalized child's rights? أسئلة سابقة

a. Withhold information from the child's mother
b. Ask mother to change her clothes into white
c. Discuss child condition with her friends
d. Accommodate the mother to stay with child

40 A hospitalized toddler, for two days, cries because he left alone. The nurse realizes this behavior is typical of: أسئلة سابقة

a. Regression.
b. Despair.
c. Protest.
d. Stranger anxiety.

41 Which of the following should the nurse do to prevent separation anxiety in a hospitalized toddler? تكرر في 2 مصادر أسئلة سابقة

a. Assume the parental role when parents are not present.
b. Limit visits by siblings and relatives.
c. Establish a routine similar to child's home.
d. Rotate nursing staff so the child becomes comfortable

42 Hospitalized children have the right to be admitted to hospital only when their: أسئلة سابقة

a. Diseases are not contagious
b. Socio-economic state is low
c. Conditions require in-patient care
d. Parents are tiered of caring for them

43 Which of the following is considered more stressful for hospitalized school age children? تكرر في 3 مصادر أسئلة سابقة

a. Fear of pain
b. Considered with appearance
c. Stranger anxiety
d. Separation from peer group

44 A child is hospitalized for a chronic illness. Initially, the child showed symptoms of depression but later started interacting with others. What does the nurse infer from the patient's behavior? The child is: أسئلة سابقة

a. Content with the care provided.
b. Showing improved social skills.
c. Getting used to the surroundings.
d. Detached from both parents.

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A 2-year-old cries loudly, clings to the mother and refuses the nurse. This is the stage of: NCLEX

a. Despair
b. Detachment
c. Protest
d. Denial

N2 Which is the major stressor for a hospitalized adolescent? NCLEX

a. Separation from parents
b. Altered body image and loss of peer contact
c. Fear of the dark
d. Stranger anxiety

N3 **Select ALL interventions that reduce separation anxiety in a hospitalized toddler.** SELECT ALL

a. Encourage rooming-in of the parent
b. Keep the home routine (bedtime rituals)
c. Bring a favorite toy or blanket from home
d. Rotate different nurses every shift
e. Limit parental visits

N4 **A child who previously cried now appears happy and interacts superficially with everyone, ignoring the parent on visits. This is:** NCLEX

a. Protest
b. Despair
c. Detachment (denial)
d. Regression

N5 **Regression during hospitalization is most common among:** NCLEX

a. Infants
b. Preschoolers
c. School-age children
d. Adolescents

N6 **Which statement best describes the effect of loss of control on a school-age child?** NCLEX

a. It has no effect
b. They fear losing control of their bodies and roles and may react with depression or hostility
c. They develop stranger anxiety
d. They return to parallel play

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Separation anxiety age	Most marked from 6 months to about 30 months (bank: 6 months–6 years).	**Protest**	Cries, screams, searches for parent, rejects strangers.
Despair	Inactive, withdrawn, sad, depressed; may regress.	**Detachment (denial)**	Superficial happiness, interacts with everyone, ignores parent — most harmful.
Toddler major stressor	Separation from parents.	**Preschooler reaction**	Regression; fear of bodily injury/mutilation.
School-age stressor	Loss of control & privacy; separation from peers.	**Adolescent stressor**	Body image/appearance; separation from peers.
Reducing separation anxiety	Rooming-in, home routines, favorite objects, consistent nurse.	**Pain substances**	Histamine, bradykinin, prostaglandins increase pain; endorphins decrease it.
Guided imagery	Pain technique effective for school-age children.	**Parents during painful procedures**	Encouraged to be present (supportive role).

Answer Key

الإجابات

1. c — Protest

2. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

3. c — Protest

4. c — Concerned with appearance

5. c — 6 months to 6 years

6. c — School-agers

7. a — Detached from both parents

8. c — Loss of control and privacy

9. a — Screams and hits the nurse

10. a — Separation anxiety

11. c — Establish a routine similar to the child's home

12. d — Encouraged and permitted freely

13. a — Histamine

14. c — Preschoolers

15. c — Separation from parents

16. a — Stops and starts again

17. b — Loss of control and privacy

18. d — Inactive, withdrawn, depressed and sad

19. a — Cries, screams and clings to parents

20. d — Separation from parents and siblings

21. b — Regresses to earlier behavior (e.g. thumb sucking, bed wetting)

22. c — 6 months to 6 years

23. c — Protest

24. c — Protest

25. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

26. d — Depressed, regressive behavior, withdrawal

27. d — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

28. c — Allow mother's rooming-in

29. b — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

30. c — Protest

31. c — Allow mother's rooming in.

32. a — It is a right of hospitalized children

33. c — Concerned with appearance.

34. c — 6 months to 6 years

35. b — Loss of control and privacy
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

36. a — Separation anxiety

37. c — Preschoolers

38. c — Separation from parents

39. d — Accommodate the mother to stay with child

40. c — Protest.

41. c — Establish a routine similar to child's home.

42. c — Conditions require in-patient care

43. d — Separation from peer group
البنك فيه نسختين من السؤال ده: مرة الإجابة loss of control and privacy ومرة separation from peer group — الاتنين من ضغوط سن المدرسة؛ اختار اللي موجود في الاختيارات.

44. d — Detached from both parents.

N1. c — Protest
Protest: crying, screaming, searching for the parent and rejecting strangers.

N2. b — Altered body image and loss of peer contact
Adolescents worry most about appearance/body image, privacy and separation from peers.

N3. a, b, c — Encourage rooming-in of the parent; Keep the home routine (bedtime rituals); Bring a favorite toy or blanket from home
Parental presence, consistent routines and transitional objects reduce stress; consistent caregivers are preferred.

N4. c — Detachment (denial)
Detachment is the last stage — superficial adjustment with loss of interest in the parent; it is the most harmful stage.

N5. b — Preschoolers
Preschoolers often regress to earlier behaviors (bed-wetting, thumb sucking) as a coping mechanism.

N6. b — They fear losing control of their bodies and roles and may react with depression or hostility
School-agers strive for independence and productivity; loss of control and privacy is very stressful for them.