PEDIATRIC NURSING (THEORY) · LECTURE 5 · QUESTION BANK

High-Risk Neonate

حديث الولادة عالي الخطورة — بنك الأسئلة

40	6	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

ميدترم 2020 · ميدترم 2021 · ميدترم 2022 · ميدترم 2024 (ترم أول 2023-2024) · كويز 1 – 2022 (نسخة بلاك بورد) · تجميعة صور ميدترمات سابقة · كويز (ملف DOC-2022) · كويز 1 (ملف Quiz 1 and 2) · كويز 1 إعادة – ترم تاني 2023/2024 · أسئلة سابقة

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 ميدترم 2020

An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر

a. True
b. False

2 ميدترم 2020

The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

a. Extremely low
b. Appropriate
c. Moderately low
d. Very low

3 ميدترم 2020

Which of the following is a characteristic of prematurity? تكرر في 2 مصادر

a. Well-developed genitalia
b. Sole full of creases
c. Strong grasp reflex
d. Excessive lanugo hair

4 ميدترم 2021

The nurse is caring for an infant born at 40 weeks and weighing 1800 g. What would the nurse expect in the chart?

a. Preterm, small for gestational age
b. Full term, large for gestational age
c. Preterm, large for gestational age
d. Full term, small for gestational age

5 ميدترم 2021

The nurse is teaching a mother of a newborn about developmental care. The nurse should instruct the mother to:

a. Provide more than one stimulus at a time
b. Stop if there is evidence of overstimulation
c. Offer it during a period of sleeping
d. Provide it for a long period

6 The nurse observes a preterm infant with fine hair covering the forehead, back and shoulders. This finding correctly describes: — ميدترم 2021

a. Vernix
b. Milia
c. Lanugo
d. Mottling

7 The best intervention to prevent infection in a high-risk neonate begins with: — ميدترم 2022

a. Not wearing a fresh scrub on care
b. Monitoring the mortality rate of neonates
c. Observing continually for signs of infection
d. Meticulous hand washing of all persons in contact

8 The best nursing intervention to deal with a preterm neonate in the NICU: تكرر في 2 مصادر — ميدترم 2022

a. Minimal handling during care
b. Use pen in length measurement
c. Cover the neonate with linen
d. Leave the window of the incubator open

9 The best nursing intervention to deal with a preterm neonate in the NICU: تكرر في 2 مصادر — ميدترم 2024 (ترم أول 2023-2024)

a. Cover the neonate with linen
b. Minimal handling during care
c. Leave the window of the incubator open
d. Use pen in length measurement

10 To prevent heat loss in high-risk neonates resulting from evaporation, the nurse should: — ميدترم 2024 (ترم أول 2023-2024)

a. Warm the crib pad
b. Turn on the radiant warmer
c. Close the room's doors
d. Dry the infant in a warm towel

11 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر — كويز 1 – 2022 (نسخة بلاك بورد)

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

12 The best nursing intervention to conserve energy in a high-risk neonate is to: — كويز 1 إعادة – ترم ثاني 2023/2024

a. Minimize neonate handling
b. Increase environmental noise
c. Encourage oral feeding
d. Maintain patent airway

13 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر — كويز 1 (ملف Quiz 1 and 2)

a. Low birth weight
b. Very very low birth weight
c. Very low birth weight
d. Appropriate birth weight

14 Which one of the following is considered a characteristic of being a preterm infant? تكرر في 2 مصادر — كويز 1 (ملف Quiz 1 and 2)

a. Low-set ears that will rise later
b. Lack of sole creases on his feet
c. Continuous irritability or crying
d. Scant lanugo on her back and arms

15 Extremely low birth weight infants — their weight is: تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. 1500 to 2499 gm
b. 1000 to 1499 gm
c. More than 2500 gm
d. Less than 1000 gm

16 Which of the following interventions should be included in the preterm infant's care plan? تكرر في 2 مصادر

كويز (ملف DOC-2022)

a. Use scissors to remove tape
b. Promote developmental care
c. Feed dextrose water immediately after birth
d. Excess handling

17 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Moderately low
b. Very low
c. Appropriate
d. Extremely low

18 Which of the following is a characteristic of prematurity? تكرر في 2 مصادر

تجميعة صور ميدترمات سابقة

a. Well-developed genitalia
b. Sole full of creases
c. Strong grasp reflex
d. Excessive lanugo hair

19 The nurse should prevent heat loss in the newborn infant resulting from radiation by:

تجميعة صور ميدترمات سابقة

a. Drying the infant with a warm blanket
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Closing the doors to the room

20 An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

21 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر

تجميعة صور ميدترمات سابقة

a. Extremely low
b. Appropriate
c. Moderately low
d. Very low

22 An infant whose weight falls above the 90th percentile on the intrauterine growth charts is considered large for gestational age. (True / False) تكرر في 3 مصادر

تجميعة صور ميدترمات سابقة

a. True
b. False

23 The nurse should prevent heat loss in the newborn infant resulting from radiation by:

تجميعة صور ميدترمات سابقة

a. Closing the doors to the room
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Drying the infant with a warm blanket

24 The best intervention to prevent infection in high-risk neonate begins with: أسئلة سابقة

a. Not wearing a fresh scrub on care.
b. Monitor mortality rate of neonate.
c. Observing continually for signs of infection.
d. Meticulous hand washing of all persons in contact.

25 The best nursing intervention to deal with preterm neonate in NICU: أسئلة سابقة

a. Minimal handling during care.
b. Use pen in length measurement
c. Cover neonate with linen
d. Leave window of incubator open

26 The neonate whose birth weight is 2000 gm. is considered: تكرر في 4 مصادر أسئلة سابقة

a. Moderately low
b. Very low
c. Appropriate
d. Extremely low

27 Which of the following is characteristics of prematurity? أسئلة سابقة

a. Well developed genitalias
b. Sole full with creases
c. Strong grasp reflex
d. Excessive lanugo hair

28 The nurse assisted with the delivery of a newborn. Which nursing action is most effective in preventing heat loss by evaporation? أسئلة سابقة

a. Warming the crib pad
b. Closing the doors to the room
c. Drying the infant with a warm blanket,
d. Turning on the overhead radiant warmer

29 An infant whose weight falls above the 90th. percentile on the intrauterine growth charts considered large for gestational age تكرر في 3 مصادر أسئلة سابقة

a. True
b. False

30 The nurse should prevent the heat loss in the new born infant resulting from radlation by: أسئلة سابقة

a. Closing the doors to the room
b. Turning on the overhead radiant warmer
c. Warming the crib pad
d. Drying the infant with a warm blanket

31 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر أسئلة سابقة

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

32 The nurse caring for an infant born at 40 weeks and weighing 1800g. What statement would nurse expect in chart أسئلة سابقة

a. Preterm a small for gestational age.
b. A full term a large for gestational age.
c. A preterm a large for gestational age.
d. A full term a small for gestational age.

33 The nurse observes a preterm infants with fine hair covered the forehead, back and shoulders. This finding correctly describes: أسئلة سابقة

a. Vernix.
b. Millia.
c. Lanugo.
d. Mottling.

34 Extremely low birth weight infants, their weight is: تكرر في 2 مصادر أسئلة سابقة

a. 1500 to 2499 gm.
b. 1000 to 1499 gm.
c. More than 2500 gm.
d. Less than 1000 gm.

35 Which of the following interventions should be included in the preterm ?infant's care plan أسئلة سابقة
تكرر في 2 مصادر

a. Use scissors to remove tape
b. Promote developmental care.
c. Feed dextrose water immediately after birth.
d. Excess handling.

36 The neonate infant whose birth weight is 2000 gm is considered: أسئلة سابقة

a. Low birth weight.
b. Very very low birth weight.
c. Very low birth weight.
d. Appropriate birth weight.

37 Which one of the following is considered a characteristic of being a preterm infant? أسئلة سابقة
تكرر في 2 مصادر

a. Low-set ears that will rise later.
b. Lack of sole creases on his feet.
c. Continuous irritability or crying
d. Scant lanugo on her back and arms.

38 To prevent heat loss in the high risk neonates resulting from evaporation the nurse should... أسئلة سابقة

a. Warm the crib pad
b. Turn on the radiant warmer
c. Close the room' doors
d. Dry the infant in a warm towel

39 Which of the following is a possible complication of prematurity? تكرر في 2 مصادر أسئلة سابقة

a. Respiratory distress syndrome
b. Increased blood pressure
c. High blood sugar
d. Hypotension

40 The neonate whose birth weight is 2000 gm is considered: تكرر في 4 مصادر أسئلة سابقة

a. Appropriate birth weight
b. Low birth weight
c. Very low birth weight
d. Very very low birth weight

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A neonate born at 39 weeks weighs 2100 g. The correct classification is: NCLEX

a. Preterm, AGA
b. Term, SGA (low birth weight)
c. Post-term, LGA
d. Term, AGA

N2 **Which finding is characteristic of a preterm infant?** NCLEX

a. Deep sole creases
b. Abundant lanugo and few sole creases
c. Strong suck and cry
d. Well-formed ear cartilage

N3 **Select ALL physiological handicaps of the premature infant.** SELECT ALL

a. Poor temperature regulation
b. Respiratory difficulties and apnea
c. Susceptibility to infection
d. Strong immunity from maternal antibodies
e. Immature liver (jaundice, bleeding)

N4 **To prevent heat loss by CONVECTION the nurse should:** NCLEX

a. Dry the baby immediately
b. Keep the baby away from drafts and keep incubator portholes closed
c. Warm the stethoscope
d. Place the baby under a radiant warmer

N5 **Very low birth weight (VLBW) refers to a birth weight of:** NCLEX

a. Less than 2500 g
b. Less than 1500 g
c. Less than 1000 g
d. 2500–4000 g

N6 **Developmental care in the NICU includes:** NCLEX

a. Bright continuous lighting
b. Clustering care, minimal handling and reducing noise
c. Frequent unnecessary stimulation
d. Separating parents from the baby

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Preterm	Born before completion of 37 weeks.	**LBW / VLBW / ELBW**	<2500 g / <1500 g / <1000 g.
SGA / LGA	Below 10th / above 90th percentile for gestational age.	**Preterm features**	Excess lanugo, thin skin, few sole creases, weak reflexes, soft ear cartilage.
Heat loss - evaporation	Prevent by drying immediately with warm towel.	**Heat loss - radiation**	Prevent with radiant warmer/warming surroundings.
Heat loss - convection	Avoid drafts; close incubator portholes.	**Heat loss - conduction**	Warm surfaces (crib pad, scale, stethoscope).
Developmental care	Minimal handling, clustered care, dim light, low noise, nesting.	**Infection prevention**	Meticulous hand washing of everyone in contact.

Premature handicaps	Poor thermoregulation, respiratory problems, infection, immature liver/kidney, poor feeding.	**Complication of prematurity**	Respiratory distress syndrome (surfactant deficiency).

Answer Key

الإجابات

1. a — True

2. c — Moderately low

3. d — Excessive lanugo hair

4. d — Full term, small for gestational age

5. b — Stop if there is evidence of overstimulation

6. c — Lanugo

7. d — Meticulous hand washing of all persons in contact

8. a — Minimal handling during care

9. b — Minimal handling during care

10. d — Dry the infant in a warm towel

11. a — Respiratory distress syndrome

12. a — Minimize neonate handling

13. a — Low birth weight

14. b — Lack of sole creases on his feet
preterm وده غلط — الـ scant lanugo في صورة الكويز الطالب اختار lack of sole creases (زي بنك، والإجابة الصح lanugo كتير (excessive) عنده الأسئلة).

15. d — Less than 1000 gm

16. b — Promote developmental care

17. a — Moderately low

18. d — Excessive lanugo hair

19. b — Turning on the overhead radiant warmer
الفقد بالإشعاع بيتمنع بتدفئة الأسطح اللي حوالين الطفل (radiant warmer). وده بيمنع الـ closing doors في صورة واحدة الطالب اختار convection.

20. a — True

21. c — Moderately low

22. a — True

23. b — Turning on the overhead radiant warmer
الفقد بالإشعاع بيتمنع بتدفئة الأسطح اللي حوالين الطفل (radiant warmer). وده بيمنع الـ closing doors في صورة واحدة الطالب اختار convection.

24. d — Meticulous hand washing of all persons in contact.

25. a — Minimal handling during care.

26. a — Moderately low

27. d — Excessive lanugo hair

28. c — Drying the infant with a warm blanket,

29. a — True

30. b — Turning on the overhead radiant warmer

31. a — Respiratory distress syndrome

32. d — A full term a small for gestational age.

33. c — Lanugo.

34. d — Less than 1000 gm.

35. b — Promote developmental care.

36. a — Low birth weight.

37. b — Lack of sole creases on his feet.
preterm وده غلط — الـ scant lanugo في صورة الكويز الطالب اختار lack of sole creases (زي بنك، والإجابة الصح lanugo كتير (excessive) عنده الأسئلة).

38. d — Dry the infant in a warm towel

39. a — Respiratory distress syndrome

40. b — Low birth weight

N1. b — Term, SGA (low birth weight)
Term = 37–42 weeks; weight below the 10th percentile/under 2500 g makes the baby SGA and LBW.

N2. b — Abundant lanugo and few sole creases
Preterm infants have abundant lanugo, thin skin, few sole creases, weak reflexes and soft ear cartilage.

N3. a, b, c, e — Poor temperature regulation; Respiratory difficulties and apnea; Susceptibility to infection; Immature liver (jaundice, bleeding)
Premature babies have poor heat regulation, immature lungs, weak immunity and immature liver/kidneys.

N4. b — Keep the baby away from drafts and keep incubator portholes closed
Convection = loss to moving air; evaporation is prevented by drying, conduction by warming surfaces, radiation by warming surroundings.

N5. b — Less than 1500 g
LBW <2500 g, VLBW <1500 g, ELBW <1000 g.

N6. b — Clustering care, minimal handling and reducing noise
Developmental care reduces stress and conserves energy: cluster care, dim lights, low noise, nesting and parental involvement.