PEDIATRIC NURSING (THEORY) · LECTURE 7 · QUESTION BANK

Accident Prevention

الوقاية من الحوادث — بنك الأسئلة

20	6	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة · ملف (Quiz 1 and 2) كويز 2 شعبة 3 · كويز 2 – 2022 · كويز 2 – 2021 · ميدترم 2024 (ترم أول 2023-2024)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 Toddlers are more prone to accidents because of their: — ميدترم 2024 (ترم أول 2023-2024)

a. Curiosity
b. Rank
c. Race
d. Growth

2 When the mother keeps all medicines and household products out of the reach of children, this represents: — ميدترم 2024 (ترم أول 2023-2024)

a. Passive preventive strategy
b. Posttraumatic care role
c. Prevent complication
d. Active preventive strategy

3 The common site of accidental injuries in children under the age of 15 years is: — كويز 2 – 2021

تكرر في 2 مصادر

a. Home
b. School
c. Street
d. Club

4 Which of the following is the most common injury in infants? تكرر في 2 مصادر — كويز 2 – 2021

a. Motor vehicle injuries
b. Burns
c. Suffocation
d. Drowning

5 The incidence of injuries among 2–6-year-old children in Egypt is as follows: — كويز 2 – 2022

a. Boys higher than girls
b. Girls higher than boys
c. Boys are equal to girls
d. Boys are twice girls

6 Which one of the following is considered the most common cause of injuries among rural children? — كويز 2 – 2022

a. Falls
b. Poisons
c. Drowning
d. Fractures

7 The most common indoor accident among 6-month-old infants is: كويز 2 شعبة 3 (ملف Quiz 1 and 2)

تكرر في 2 مصادر

a. Drowning
b. Poisoning
c. Burning
d. Falling

8 Which of the following measures should be followed at home to ensure child safety? كويز 2 شعبة 3 (ملف Quiz 1 and 2)

a. Medicines are to be kept on the dinner table
b. Electrical plugs must be covered and at low level
c. Kitchen curtains are better to be used than doors
d. Furniture is to be away from the window

9 When the mother keeps all medicines and household products out of the reach from children, this represents أسئلة سابقة

a. Massive preventive strategy
b. Posttraumatic care role
c. Prevent complication
d. Active preventive strategy

10 Which patients are most at risk for sudden infant death syndrome? (Select all that apply) أسئلة سابقة

a. A newborn whose mother had 1 drink of alcohol early in pregnancy
b. A 2-month-old infant who sleeps in the bed with his mother
c. A 1-month-old infant who prefers to sleep on his stomach
d. A 1-month-old infant who prefers to sleep on her back
e. A newborn whose mother smokes 2 packs/day

11 What is the leading cause of death in children older than 1 year? أسئلة سابقة

a. Suicide
b. Chronic respiratory illness
c. Unintentional injuries
d. Congenital anomalies

12 What should the nurse teach parents of a toddler about prevention of accidental poisoning? (Select all that apply) أسئلة سابقة

a. Keep toxic substances in the garage
b. Remove colorful labels from containers of toxic materials
c. Know the number of the nearest poison control center
d. Caution the child against eating nonedible items such as plants
e. Discard empty poison containers

13 Which groups of infants have a higher risk for SIDS? (Select all that apply) أسئلة سابقة

a. Male gender
b. Low Apgar scores
c. First-born infants
d. Full-term infants
e. Low birth weight

14 For what injuries are crawling infants most at risk? أسئلة سابقة

a. Aspiration/poisoning
b. Falls
c. Motor vehicle collisions
d. Burns

15 Which statement is true regarding injury prevention in infants and toddlers? أسئلة سابقة

a. Repeated "No" is needed to be effective with this age group.
b. Injury prevention and parent education should be handled on a growth and developmental basis.
c. Falls are the most common injuries for these age groups.
d. All injuries can be prevented with proper education regarding safety measures.

16 The most common indoor accident among 6-month-old infants is: تكرر في 2 مصادر أسئلة سابقة

a. Drowning
b. Poisoning
c. Burning
d. Falling

17 Which of the following measures should at home to ensure child safety? أسئلة سابقة

a. Medicines are to be kept on dinner table
b. Electrical plugs must be covered and at low level
c. Kitchen curtains are better to be used than doors
d. Furnitures are to be away from the window

18 Which of the following is the most common injury in infants? تكرر في 2 مصادر أسئلة سابقة

a. Motor vehicle injuries
b. Burns
c. Suffocation
d. Drowning

19 The common site of accidental injuries in children under the age of 15 years is: أسئلة سابقة
تكرر في 2 مصادر

a. Home
b. School
c. Street
d. Club

20 The incidence of injuries among 2 – 6 years old children at Egypt is as following أسئلة سابقة

a. Boys higher than girls
b. Girls higher than boys
c. Boys are equal girls
d. Boys are twice girls

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A mother of a 9-month-old asks which injury her crawling infant is most at risk for. The nurse answers: NCLEX

a. Motor vehicle injury as a pedestrian
b. Aspiration of small objects and poisoning
c. Sports injuries
d. Firearm injuries

N2 Which is an example of an ACTIVE (behavioral) preventive strategy? NCLEX

a. Child-resistant medicine caps made by the factory
b. Building codes for window guards
c. Parent stores medicines and cleaners in a locked high cabinet
d. Automatic airbags in cars

N3 Select ALL teaching points that reduce the risk of SIDS. SELECT ALL

a. Place the infant on the back to sleep
b. Use a firm mattress without pillows or soft toys
c. Bed-share with parents every night
d. Avoid smoking around the infant
e. Overdress the infant during sleep

N4 Why are toddlers especially prone to accidents? NCLEX

a. Fully developed judgment
b. Curiosity and increasing mobility with poor awareness of danger
c. Peer pressure
d. Risk-taking to impress friends

N5 In the epidemiological model of injury, a hot cup of tea that scalds a toddler represents the: NCLEX

a. Host
b. Agent
c. Environment
d. Vector-free factor

N6 Which statement about injuries among Egyptian children is correct? NCLEX

a. Girls have more injuries than boys
b. Drowning and animal bites occur mainly among rural children
c. Injuries are rare under 6 years
d. Burns are the least common injury

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Accident	Unintentional event causing injury from external forces or environmental hazards, without deliberate intent.	**Injury**	Damage to the body from acute exposure to thermal, mechanical, electrical or chemical energy, or absence of heat/oxygen.
Impairment	Losing part/all of a limb or having a defective organ or body mechanism.	**Disability**	Loss or reduction of functional ability.
Handicap	Permanent disadvantage or restriction of activity caused by disability.	**Epidemiological model**	Host (child) + Agent (energy/vector) + Environment.
Active prevention	Requires repeated action by the person (e.g., parent locks medicines).	**Passive prevention**	Protects automatically without individual action (e.g., child-resistant caps, airbags).
Most common non-fatal injury in young children	Falls.	**Leading cause of death after age 1**	Unintentional injuries.

Egypt: injuries 2–6 years	Higher in boys than girls; drowning & animal bites mostly in rural children.
Crawling infant hazard	Aspiration of small objects and poisoning.

Answer Key

الإجابات

1. a — Curiosity

2. d — Active preventive strategy

3. a — Home

4. c — Suffocation

5. a — Boys higher than girls

6. c — Drowning

7. d — Falling
أشيع (falls) الملف مفيهوش إجابة معلّمة — إجابة مقترحة: الوقوع حوادث البيت عند الرضّع اللي بيبدأوا يتقلّبوا ويزحفوا.

8. d — Furniture is to be away from the window

9. d — Active preventive strategy

10. b, c, e — A 2-month-old infant who sleeps in the bed with his mother; A 1-month-old infant who prefers to sleep on his stomach; A newborn whose mother smokes 2 packs/day

11. c — Unintentional injuries

12. c, d, e — Know the number of the nearest poison control center; Caution the child against eating nonedible items such as plants; Discard empty poison containers

13. a, b, e — Male gender; Low Apgar scores; Low birth weight

14. a — Aspiration/poisoning

15. b — Injury prevention and parent education should be handled on a growth and developmental basis.

16. d — Falling
أشيع (falls) الملف مفيهوش إجابة معلّمة — إجابة مقترحة: الوقوع حوادث البيت عند الرضّع اللي بيبدأوا يتقلّبوا ويزحفوا.

17. d — Furnitures are to be away from the window
إجابة البنك (furniture away from window) صحيحة؛ لاحظ إن الـ plugs لازم تتغطى بس مش 'at low level'.

18. c — Suffocation

19. a — Home

20. a — Boys higher than girls

N1. b — Aspiration of small objects and poisoning
Crawling infants explore with their mouths; aspiration of small objects and ingestion of household products are the classic hazards of this stage.

N2. c — Parent stores medicines and cleaners in a locked high cabinet
Active strategies need a repeated action by the person (parent locking cabinets); passive strategies protect automatically without individual effort.

N3. a, b, d — Place the infant on the back to sleep; Use a firm mattress without pillows or soft toys; Avoid smoking around the infant
Back-to-sleep, firm bare crib and a smoke-free environment lower SIDS risk; bed-sharing and overheating increase it.

N4. b — Curiosity and increasing mobility with poor awareness of danger
Toddlers are curious, mobile and impulsive but cannot yet recognize danger — the key developmental risk factor.

N5. b — Agent
The agent (vector) is the energy source that causes the injury — here thermal energy from the hot liquid; the child is the host.

N6. b — Drowning and animal bites occur mainly among rural children
Egyptian data: boys aged 2–6 years have more injuries; drowning, animal bites and sunstroke occur mostly in rural children.