PEDIATRIC SURGERY · LECTURE 2 · QUESTION BANK

Common Pediatric Surgical Conditions (Hernia, Hydrocele & UDT)

الحالات الجراحية الشائعة (الفتق والقيلة المائية والخصية المعلقة) — بنك الأسئلة

21	6	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة · فاينال (مراجعة أسئلة) · ميدترم 2025 · ميدترم السنة اللي فاتت (Microsoft Forms)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 **Strangulation in inguinal hernia:** — ميدترم السنة اللي فاتت (Microsoft Forms)

a. May cause death
b. Is due to delay in diagnosis
c. Is due to delay in management
d. All of the above

2 **Glans injury is a dangerous injury during ..... surgery.** — ميدترم السنة اللي فاتت (Microsoft Forms)

a. Circumcision
b. Cleft palate
c. Tongue tie
d. Cleft lip

3 **In cases of neonatal inguinal hernia, we wait for the age of ..... to do surgery.** — ميدترم السنة اللي فاتت (Microsoft Forms)

a. 1 year
b. 2 years
c. 3 years
d. None of the above — once diagnosed, do surgery as soon as possible

4 **At what age is surgical correction recommended for undescended testes?** — ميدترم 2025

a. 1 month
b. 6 months to 1 year
c. 5 years
d. Adolescence

5 **Most common site for undescended testes?** — ميدترم 2025

a. Abdominal cavity
b. Inguinal canal
c. Scrotal neck
d. Perineum

6 Congenital hydrocele is caused by: ميدترم 2025

a. Trauma
b. Infection
c. Incomplete closure of processus vaginalis
d. Tumor

7 What is the main clinical feature of hydrocele? ميدترم 2025

a. Hard, tender scrotal swelling
b. Soft, non-tender scrotal swelling that transilluminates
c. Scrotal redness
d. Fever

8 The treatment for the most common congenital hydrocele is: ميدترم 2025

a. Immediate surgery
b. Observation until the age of 2 years
c. Antibiotics
d. Diuretics

9 What is the most common type of hernia in children? ميدترم 2025

a. Umbilical hernia
b. Inguinal hernia
c. Femoral hernia
d. Epigastric hernia

10 Which of the following is a sign of an incarcerated inguinal hernia? ميدترم 2025

a. Reducible, painless swelling
b. Non-reducible, tender swelling
c. Swelling only during crying
d. No swelling

11 Definitive treatment of inguinal hernia in a child: ميدترم 2025

a. Manual reduction
b. Herniotomy/herniorrhaphy (surgical repair)
c. Observation
d. Use of truss

12 Which of the following is NOT associated with untreated cryptorchidism (undescended testis)? ميدترم 2025

a. Infertility
b. Testicular torsion
c. Appendicitis
d. Testicular cancer

13 Complete: Indications of circumcision are ..... فاينال (مراجعة أسئلة)

.....

14 Complete: Common risks after circumcision are ..... فاينال (مراجعة أسئلة)

.....

15 In case of inguinal hernia, wait for surgery until the child reaches 3 years. (True / False) فاينال (مراجعة أسئلة)

a. True
b. False

16 Umbilical hernia is an emergency condition. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

17 In case of unpalpable undescended testis, perform MRI. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

18 Glans injury is most common in circumcision. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

19 Inguinal hernia in females does not need surgery. (True / False) أسئلة سابقة

a. True **b.** False

20 Inguinal hernia may be treated by antibiotics or needs MRI. (True / False) أسئلة سابقة

a. True **b.** False

21 Complete: Inguinal hernia surgery should be done early to prevent ..... أسئلة سابقة

.....

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A 3-month-old has a groin swelling that increases with crying and reduces at rest. The nurse anticipates: NCLEX

a. Observation until 3 years
b. Elective herniotomy soon after diagnosis
c. Antibiotics
d. Truss application

N2 Which finding indicates an incarcerated inguinal hernia? NCLEX

a. Soft, reducible swelling
b. Tender, irreducible swelling with vomiting
c. Swelling that transilluminates
d. Swelling only when crying

N3 Select ALL features of a congenital communicating hydrocele. SELECT ALL

a. Due to patent processus vaginalis **b.** Soft, painless scrotal swelling
c. Transilluminates **d.** Usually needs emergency surgery
e. Many resolve by 1–2 years

N4 Orchiopexy for undescended testis is ideally performed at: NCLEX

a. Birth **b.** 6–12 months (before 1–2 years)
c. 5 years **d.** Puberty

N5 **Complications of untreated cryptorchidism include all EXCEPT:** NCLEX

a. Infertility
b. Testicular torsion
c. Testicular cancer
d. Appendicitis

N6 **After circumcision, the nurse teaches parents to watch mainly for:** NCLEX

a. Bleeding and signs of infection
b. Constipation
c. Hearing problems
d. Rash on the face

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Inguinal hernia	Most common hernia in children; patent processus vaginalis.	**Inguinal hernia timing**	Operate once diagnosed (no waiting).
Incarceration	Irreducible tender swelling — emergency.	**Strangulation**	Compromised blood supply; may cause death; due to delayed diagnosis/management.
Hernia in females	Needs surgery — may contain the ovary.	**Hydrocele**	Fluid in tunica vaginalis; soft, painless, transilluminates.
Hydrocele treatment	Observe until ~2 years; surgery if persists/communicating.	**Undescended testis**	Most common site: inguinal canal.
Orchiopexy age	6–12 months (before 2 years).	**UDT complications**	Infertility, torsion, malignancy, hernia, psychological.
Unpalpable testis	Laparoscopy (not MRI).	**Circumcision risks**	Bleeding, infection, pain; glans injury is serious.

Answer Key

الإجابات

1. d — All of the above

2. a — Circumcision

3. d — None of the above — once diagnosed, do surgery as soon as possible

4. b — 6 months to 1 year

5. b — Inguinal canal

6. c — Incomplete closure of processus vaginalis

7. b — Soft, non-tender scrotal swelling that transilluminates

8. b — Observation until the age of 2 years

9. b — Inguinal hernia

10. b — Non-reducible, tender swelling

11. b — Herniotomy/herniorrhaphy (surgical repair)

12. c — Appendicitis

13. ✎ — Phimosis, paraphimosis, recurrent UTI

14. ✎ — Bleeding, infection, pain

15. b — False

16. b — False

17. b — False

18. b — False
الملف كاتب False (مع إن ميدترم، الأشيع بعد الطهارة النزيف والعدوى) glans injury — إصابة خطيرة في الطهارة السنة اللي فاتت قال إن خطيرة ≠ الأشيع.

19. b — False
False — لازم جراحة لأن الفتق ممكن يكون فيه المبيض.

20. b — False

21. ✎ — Strangulation

N1. b — Elective herniotomy soon after diagnosis
Inguinal hernia does not resolve spontaneously; surgery is done once diagnosed to prevent incarceration.

N2. b — Tender, irreducible swelling with vomiting
Irreducible tender swelling with vomiting/irritability means incarceration — an emergency.

N3. a, b, c, e — Due to patent processus vaginalis; Soft, painless scrotal swelling; Transilluminates; Many resolve by 1–2 years
Most congenital hydroceles resolve spontaneously; surgery if persistent after ~2 years.

N4. b — 6–12 months (before 1–2 years)
Early orchiopexy protects fertility and allows malignancy surveillance.

N5. d — Appendicitis
Appendicitis is unrelated.

N6. a — Bleeding and signs of infection
Bleeding, infection and pain are the common risks after circumcision.