PEDIATRIC SURGERY · LECTURE 4 · QUESTION BANK

Intussusception

الانغلاف المعوي — بنك الأسئلة

5	6	7
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة · سؤال فاينال (صورة من الجروب) · ميدترم السنة اللي فاتت (Microsoft Forms) · كويز 1 – 2023/2024 (شعبة 4) (أطفال نظري)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 ميدترم السنة اللي فاتت (Microsoft Forms)

Intussusception:

a. Always needs surgery
b. Pneumatic reduction may solve the problem
c. You can wait for 8 hours
d. The treatment is medical

2 سؤال فاينال (صورة من الجروب)

Air enema is both diagnostic and therapeutic in intussusception. (True / False)

a. True **b.** False

3 أسئلة سابقة

In intussusception, surgery is always needed. (True / False)

a. True **b.** False

4 أسئلة سابقة

Which clinical manifestations would the nurse expect to observe while caring for the child diagnosed with intussusception? تكرر في 2 مصادر

a. Currant jelly stool **b.** Foul-smelling stool
c. Dark brown stool **d.** Watery stool

5 كويز 1 – 2023/2024 (شعبة 4) (أطفال نظري)

Which clinical manifestations would the nurse expect to observe while caring for the child diagnosed with intussusception?

تكرر في 2 مصادر

a. Currant jelly stool **b.** Foul-smelling stool
c. Dark brown stool **d.** Watery stool

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A 9-month-old has sudden colicky pain with drawing up of legs, vomiting and red currant-jelly stools. The likely diagnosis is: NCLEX

a. Pyloric stenosis
b. Intussusception
c. Hirschsprung disease
d. Appendicitis

N2 First-line management of uncomplicated intussusception is: NCLEX

a. Immediate laparotomy
b. Air (pneumatic) or hydrostatic enema reduction
c. Antibiotics only
d. Laxatives

N3 Select ALL signs that a reduced intussusception has resolved. SELECT ALL

a. Passage of normal brown stool
b. Disappearance of the abdominal mass
c. Child becomes calm and pain-free
d. Return of currant-jelly stool
e. Increasing distension

N4 The most common type of intussusception in children is: NCLEX

a. Ileocolic
b. Colocolic
c. Jejunojejunal
d. Gastroduodenal

N5 The diagnostic imaging of choice for intussusception is: NCLEX

a. Abdominal ultrasound (target sign)
b. Barium swallow
c. CT head
d. Chest X-ray

N6 Peak age for intussusception is: NCLEX

a. Neonatal period
b. 6 months to 2 years
c. School age
d. Adolescence

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Intussusception	Telescoping of bowel (ileocolic); 6 months–2 years.	**Intussusception triad**	Colicky pain, sausage mass, currant-jelly stool.
Intussusception diagnosis	Ultrasound (target sign).	**Intussusception treatment**	Air/hydrostatic enema (diagnostic + therapeutic); surgery if fails.
Mass	Sausage-shaped RUQ mass; empty RLQ (Dance sign).	**Red flags**	Peritonitis, perforation, shock → surgery.
After reduction	Watch for normal stool; recurrence risk.		

Answer Key

الإجابات

1. b — Pneumatic reduction may solve the problem

2. a — True
الإجابة الرسمية في الفورم True. (تعليق الصورة على الجروب قال إن كمان barium enema الـ diagnostic وtherapeutic.)

3. b — False

4. a — Currant jelly stool

5. a — Currant jelly stool

N1. b — Intussusception
Classic triad: colicky pain, sausage-shaped mass and currant-jelly stool.

N2. b — Air (pneumatic) or hydrostatic enema reduction
Enema reduction is both diagnostic and therapeutic; surgery if it fails or peritonitis is present.

N3. a, b, c — Passage of normal brown stool; Disappearance of the abdominal mass; Child becomes calm and pain-free
Passage of normal stool and relief of pain indicate successful reduction.

N4. a — Ileocolic
Terminal ileum telescopes into the colon.

N5. a — Abdominal ultrasound (target sign)
Ultrasound shows target/doughnut sign.

N6. b — 6 months to 2 years
Most cases occur between 3 months and 3 years.