PEDIATRIC SURGERY · LECTURE 5 · QUESTION BANK

Anorectal Malformations

تشوهات الشرج والمستقيم — بنك الأسئلة

26	6	11
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة · فاينال (مراجعة أسئلة) · ميدترم 2025 · ميدترم السنة اللي فاتت (Microsoft Forms)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 ميدترم السنة اللي فاتت (Microsoft Forms)

All of the following are preoperative preparations of neonatal intestinal obstruction EXCEPT:

a. Intravenous fluids
b. CBC, serum electrolytes and coagulation profile
c. Ryle/NGT
d. Oral feeding

2 ميدترم السنة اللي فاتت (Microsoft Forms)

Meconium ileus is a cause of bowel obstruction in:

a. Neonates
b. Toddlers
c. Preschool
d. School age

3 ميدترم السنة اللي فاتت (Microsoft Forms)

In cases of high imperforate anus we do:

a. Anoplasty
b. Colostomy, anoplasty
c. Colostomy, anoplasty, closure of colostomy
d. Conservative management

4 ميدترم السنة اللي فاتت (Microsoft Forms)

Which of the following is an essential nursing care in all babies with intestinal obstruction?

a. Endotracheal tube
b. Intravenous line
c. Clear oral fluids
d. Enema

5 ميدترم السنة اللي فاتت (Microsoft Forms)

Presentation of intestinal obstruction includes the following EXCEPT:

a. Dehydration
b. Low urine output
c. Hyperactive baby
d. Vomiting

6 Causes of neonatal intestinal obstruction include all of the following EXCEPT: — ميدترم السنة اللي فاتت (Microsoft Forms)

a. Jejunal atresia
b. Meconium ileus
c. Tracheo-esophageal fistula
d. Hirschsprung's disease

7 In intestinal obstruction, what do you insert to protect from aspiration? — ميدترم السنة اللي فاتت (Microsoft Forms)

a. NGT/Ryle
b. Urine catheter
c. Umbilical catheter
d. Central line

8 Which of the following is a common sign of anorectal malformation in a newborn? — ميدترم 2025

a. Excessive urination
b. No meconium passage and abdominal distention
c. Jaundice
d. Cyanosis

9 What is the imaging study special for evaluation of anorectal malformation? — ميدترم 2025

a. MRI
b. Cross-table lateral X-ray
c. CT scan
d. Ultrasound

10 What is the most common anorectal malformation in males? — ميدترم 2025

a. Rectovaginal fistula
b. Rectourethral fistula
c. Rectoperineal fistula
d. Rectovesical fistula

11 Which sign is the most consistent with duodenal atresia? — ميدترم 2025

a. Triple bubble sign
b. Double bubble sign
c. Air-fluid levels
d. Pneumoperitoneum

12 What is the first management of neonatal intestinal atresia? — ميدترم 2025

a. Immediate surgical intervention
b. Antibiotic therapy
c. Nasogastric decompression
d. Fluid restriction

13 Complete: Causes of intestinal obstruction in the neonate ..... — فاينال (مراجعة أسئلة)

.....

14 Complete: Rectal biopsy is used to diagnose ..... — فاينال (مراجعة أسئلة)

.....

15 Complete: Double bubble sign on X-ray is seen in ..... — فاينال (مراجعة أسئلة)

.....

16 In all cases of imperforate anus we perform colostomy. (True / False) — فاينال (مراجعة أسئلة)

a. True
b. False

17 Hirschsprung's disease affects the colon of the neonate and results in midgut volvulus. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

18 Surgical treatment of Hirschsprung's disease is removal of the aganglionic segment. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

19 Meconium ileus causes bowel obstruction in neonates. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

20 In imperforate anus, wait until the infant reaches 30 days to perform surgery. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

21 Vomiting and failure to pass stool are signs of intestinal obstruction. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

22 Dehydration can be present in neonatal intestinal obstruction. (True / False) فاينال (مراجعة أسئلة)

a. True **b.** False

23 Complete: The operation for low imperforate anus is ..... أسئلة سابقة

.....

24 Complete: The operation for high imperforate anus is ..... أسئلة سابقة

.....

25 Complete: If a newborn does not pass meconium, suspect ..... أسئلة سابقة

.....

26 Complete: A child has intestinal obstruction; to prevent aspiration you ..... أسئلة سابقة

.....

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A newborn has no anal opening. The surgical plan for a HIGH anomaly is: NCLEX

a. Single anoplasty
b. Colostomy, then anoplasty (PSARP), then closure of colostomy
c. Observation
d. Dilatation only

N2 The nurse should first suspect imperforate anus when: NCLEX

a. The baby has jaundice
b. The first rectal temperature cannot be taken or no meconium passes in 24 h
c. The baby vomits once
d. The baby is cyanosed

N3 Colostomy care teaching for parents includes: NCLEX

a. Apply tight dressing over the stoma
b. Clean skin around stoma, use a well-fitting bag and watch stoma color
c. Avoid emptying the bag
d. Use alcohol on the stoma

N4 A cross-table lateral X-ray for a newborn with ARM is best done after: NCLEX

a. 1 hour
b. 24 hours
c. 1 week
d. 1 month

N5 The most common anorectal malformation in females is: NCLEX

a. Rectovestibular fistula
b. Rectourethral fistula
c. Cloaca only
d. Rectovesical fistula

N6 ARM is part of which association? NCLEX

a. VACTERL
b. TORCH
c. Down syndrome only
d. Turner

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Anorectal malformation	Spectrum of imperforate anus; low vs high.	**ARM imaging**	Cross-table lateral X-ray (invertogram) after 24 h.
Commonest ARM in males	Rectourethral fistula.	**Low ARM surgery**	Anoplasty.
High ARM surgery	Colostomy → PSARP/anoplasty → closure of colostomy.	**Neonatal obstruction causes**	Duodenal/jejunal atresia, ARM, Hirschsprung, meconium ileus, malrotation.
Double bubble	Duodenal atresia.	**Obstruction nursing**	NPO, NGT to prevent aspiration, IV line, fluids & electrolytes.
Females	Rectovestibular fistula commonest.	**Wangensteen/invert**	Lateral X-ray after 24 h.
Post-PSARP	Anal dilatation program.		

Answer Key

الإجابات

1. d — Oral feeding

2. a — Neonates

3. c — Colostomy, anoplasty, closure of colostomy

4. b — Intravenous line

5. c — Hyperactive baby

6. c — Tracheo-esophageal fistula

7. a — NGT/Ryle

8. b — No meconium passage and abdominal distention

9. b — Cross-table lateral X-ray

10. b — Rectourethral fistula

11. b — Double bubble sign

12. c — Nasogastric decompression

13. ✎ — Hirschsprung's disease, meconium ileus, duodenal atresia

14. ✎ — Hirschsprung's disease

15. ✎ — Duodenal atresia

16. b — False

17. b — False

18. a — True
الملف كاتب «removal of ganglionic» — المقصود إزالة الجزء الـ aganglionic.

19. a — True

20. b — False

21. a — True

22. a — True

23. ✎ — Anoplasty

24. ✎ — Colostomy then anoplasty (then closure of colostomy)

25. ✎ — Imperforate anus

26. ✎ — Insert NGT (Ryle)

N1. b — Colostomy, then anoplasty (PSARP), then closure of colostomy
High lesions need a staged approach; low lesions are treated by anoplasty.

N2. b — The first rectal temperature cannot be taken or no meconium passes in 24 h
Inspection of the perineum and failure to pass meconium are the earliest clues.

N3. b — Clean skin around stoma, use a well-fitting bag and watch stoma color
The stoma should be pink and moist; protect peristomal skin and empty the bag when 1/3 full.

N4. b — 24 hours
Allows bowel gas to reach the rectal pouch.

N5. a — Rectovestibular fistula
Males: rectourethral (bulbar) fistula.

N6. a — VACTERL
Screen for spinal, cardiac, renal anomalies.