MATERNAL & NEWBORN HEALTH NURSING · LECTURE 3 · QUESTION BANK

Preconception Care

رعاية ما قبل الحمل — بنك الأسئلة

14	8	12
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

ميدترم 2022–2023 · كويز 1 – 2023 · كويز 1 – 2022 · كويز 1 – 2021 · كويز 1 – 2020

فاينال 2019–2020 (بنك الأسئلة) · أسئلة سابقة · كويز تجريبي (فورم)

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 Lack of preconception services is considered a barrier. (True / False) — ميدترم 2022–2023

a. True **b.** False

2 What is the optimum time for preconception care? — كويز 1 – 2023

a. 1–2 weeks before the conceiving attempt
b. 1–2 months before the conceiving attempt
c. 3–6 weeks before the conceiving attempt
d. 3–6 months before the conceiving attempt

3 Which statement reflects one purpose of preconception care? — كويز 1 – 2022

a. Identify women who should not become pregnant
b. Encourage healthy lifestyles for couples desiring pregnancy
c. Ensure that women know about prenatal care
d. Ensure that pregnancy complications do not occur

4 Interconception is described as the period before the first pregnancy. (True / False) — كويز 1 – 2021

تكرر في 2 مصادر

a. True **b.** False

5 "Fit for pregnancy" is one of the preconception health concepts. (True / False) — كويز 1 – 2021

a. True **b.** False

6 Preconception care started one month before pregnancy is the best timing. (True / False) — كويز 1 – 2021

a. True **b.** False

7 Unwanted pregnancies are at greater risk for both maternal and fetal outcomes. (True / False) — كويز 1 – 2020

a. True **b.** False

8 Women with a previous history of fetal abnormality need to take folic acid before becoming pregnant. (True / False) — كويز 1 – 2020

a. True **b.** False

9 The minimum amount of folic acid supplementation a woman should use while planning pregnancy is 4 mg. (True / False) — فاينال 2019–2020 (بنك الأسئلة) — تكرر في 2 مصادر

a. True **b.** False

10 Preconception health only matters if you have health problems. (True / False) — فاينال 2019–2020 (بنك الأسئلة) — تكرر في 2 مصادر

a. True **b.** False

11 Interconception is described as the period before the first pregnancy. (True / False) — أسئلة سابقة — تكرر في 2 مصادر

a. True **b.** False

12 Preconception health only matters if you have health problems. (True / False) — أسئلة سابقة — تكرر في 2 مصادر

a. True **b.** False

13 The minimum amount of folic acid supplementation a woman should use while planning pregnancy is 4 mg. (True / False) — أسئلة سابقة — تكرر في 2 مصادر

a. True **b.** False

14 Which statement reflects the purpose of preconception care? — كويز تجريبي (فورم)

a. Identify women who should not become pregnant
b. Encourage healthy lifestyles for couples desiring pregnancy
c. Ensure that pregnancy complications do not occur
d. Ensure that women know about prenatal care

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 What is the optimal time to begin preconception care? NCLEX

a. 1 month before conception **b.** 3–6 months before a conception attempt
c. After the first missed period **d.** In the second trimester

N2 A woman planning pregnancy asks why she should start folic acid now. The nurse explains it mainly prevents: NCLEX

a. Gestational diabetes
b. Neural tube defects
c. Preterm labor
d. Anemia of pregnancy only

N3 Select ALL components of preconception care. SELECT ALL

a. Risk assessment
b. Health promotion
c. Immunization (e.g., rubella)
d. Management of chronic disease
e. Fetal non-stress testing

N4 A woman with poorly controlled diabetes wants to conceive. The priority preconception advice is: NCLEX

a. Conceive now and control glucose later
b. Achieve good glycemic control before conception
c. Stop all medications
d. Avoid all exercise

N5 Interconception health stresses waiting at least how long between pregnancies? NCLEX

a. 6 months
b. 12 months
c. 18 months
d. 36 months

N6 Which vaccine must be given at least 1 month BEFORE pregnancy, not during it? NCLEX

a. Tetanus
b. Influenza (inactivated)
c. Rubella (MMR)
d. Hepatitis B

N7 Which statement shows the client understood preconception care? NCLEX

a. It only matters if I have a health problem
b. It helps me become as healthy as possible before pregnancy
c. It starts after a positive pregnancy test
d. It replaces antenatal care

N8 Lack of preconception services is best described as: NCLEX

a. A risk factor
b. A barrier to preconception care
c. An intervention
d. A goal

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Preconception care	Interventions before pregnancy to optimize maternal and fetal outcomes	**Optimum timing**	3–6 months before conception attempt
Interconception	Period between pregnancies — wait ≥ 18 months	**Folic acid**	Taken ~3 months before pregnancy; prevents 50–70% of NTDs

Folic acid dose (course)	4 mg — especially with previous NTD/fetal anomaly
Three life-course pathways	Before first pregnancy · Managing emerging risks · Next baby and beyond
Barrier example	Lack of preconception services
Live vaccines	MMR / varicella before pregnancy — avoid during it

Two practical ideas	Planning pregnancy & Fit for pregnancy
Purpose	Encourage healthy lifestyles for couples desiring pregnancy
Unwanted pregnancy	Higher maternal and fetal risks
Slogan	Healthy women, healthy mothers, healthy babies

Answer Key

الإجابات

1. a — True

2. d — 3–6 months before the conceiving attempt

3. b — Encourage healthy lifestyles for couples desiring pregnancy

4. b — False

5. a — True

6. b — False

7. a — True

8. a — True

9. a — True
The course answer key marks this True. Note: the usual dose for low-risk women is 0.4 mg/day; 4 mg is for women with a previous neural-tube-defect pregnancy.

10. b — False

11. b — False

12. b — False

13. a — True
The course answer key marks this True. Note: the usual dose for low-risk women is 0.4 mg/day; 4 mg is for women with a previous neural-tube-defect pregnancy.

14. b — Encourage healthy lifestyles for couples desiring pregnancy

N1. b — 3–6 months before a conception attempt
3–6 months allows risks to be reduced and conditions controlled before conception.

N2. b — Neural tube defects
Folic acid taken about 3 months before pregnancy prevents 50–70% of neural tube defects.

N3. a, b, c, d — Risk assessment; Health promotion; Immunization (e.g., rubella); Management of chronic disease
NST is an antenatal fetal test, not preconception care.

N4. b — Achieve good glycemic control before conception
Hyperglycemia at conception increases miscarriage and congenital anomaly risk.

N5. c — 18 months
An interval of at least 18 months helps optimize birth outcomes.

N6. c — Rubella (MMR)
Live vaccines such as MMR are contraindicated in pregnancy; avoid conception for about 1 month after.

N7. b — It helps me become as healthy as possible before pregnancy
Preconception care is for all women of reproductive age, not only those with problems.

N8. b — A barrier to preconception care
Lack of services/providers is one of the listed barriers.