MATERNAL & NEWBORN HEALTH NURSING · LECTURE 9 · QUESTION BANK

Physiology & Nursing Management of Normal Labor

فسيولوجيا الولادة الطبيعية ورعايتها — بنك الأسئلة

47	8	16
Past exam questions	NCLEX-style	Flashcards

المصادر اللي اتجمعت منها أسئلة المحاضرة دي:

أسئلة سابقة · فاينال 2019–2020 (بنك الأسئلة) · كويز 2 – 2021 (نسخة أ) · كويز 2 – 2021 (نسخة ب) · كويز 2 – 2022 · كويز طب نسا

جنب كل سؤال مصدره. لو السؤال اتكرر في كذا امتحان مكتوب كل مرة بمصدرها، وعليه علامة "تكرر". الإجابات كلها في آخر الملف.

اللَّهُمَّ لا سَهْلَ إِلَّا ما جَعَلْتَهُ سَهْلًا، وأنتَ تَجْعَلُ الحَزْنَ إذا شِئْتَ سَهْلًا — صلِّ على النبي ﷺ

Part A · Past Exam Questions

أسئلة الدفعات السابقة

1 In a multipara, the first stage of labor usually lasts: تكرر في 2 مصادر — كويز 2 – 2022

a. 6 to 8 hours
b. 3 to 6 hours
c. 10 to 15 hours
d. 24 hours

2 Ripening of the cervix occurs during the: تكرر في 2 مصادر — كويز 2 – 2022

a. First stage
b. Second stage
c. Third stage
d. Fourth stage

3 Which of the following statements is unrelated to the signs of placental separation? تكرر في 2 مصادر — كويز 2 – 2022

a. Gush of blood
b. Uterus rises in the lower abdomen
c. Umbilical cord protrudes farther out of the vagina
d. Uterus becomes flaccid

4 When the fetus is in the right occiput anterior position and at +1 station, the presenting part is: — كويز 2 – 2022

a. 1 cm below the ischial spines
b. Directly in line with the ischial spines
c. 1 cm above the ischial spines
d. In no relationship to the ischial spines

5 The nurse is monitoring the progress of a mother in labor. Which finding should be reported to the physician immediately? كويز 2 – 2022

a. Scant bloody discharge
b. Frequent urination
c. Green-tinged amniotic fluid
d. Moderate uterine contractions

6 A patient is in the active deceleration phase. How frequently should the nurse assess her uterine contractions? كويز 2 – 2022

a. Every 5 minutes
b. Every 15 minutes
c. Every 30 minutes
d. Every 60 minutes

7 In the second stage of labor, expulsion of the fetus from the birth canal depends on which important contributor to the involuntary expulsive force? تكرر في 2 مصادر كويز 2 – 2022

a. Maternal bearing down
b. Cervical dilatation
c. Uterine contractions
d. Adequate pelvic size

8 A nullipara is admitted with cervix 6 cm, 70% effaced, at −1 station. What phase of labor is she in? تكرر في 2 مصادر كويز 2 – 2022

a. Active acceleration phase
b. Active deceleration phase
c. Latent phase
d. Expulsion phase

9 At which stage of labor is the mother advised to avoid bearing down? تكرر في 2 مصادر كويز 2 – 2022

a. Second stage
b. First stage
c. Fourth stage
d. Third stage

10 A sudden gush of blood or lengthening of the cord after delivery of the infant should warn the nurse of: كويز 2 – 2021 (نسخة ب)

a. Placenta accreta
b. Placental separation
c. Placental retention
d. Abruptio placentae

11 Ripening of the cervix occurs during the: تكرر في 2 مصادر كويز 2 – 2021 (نسخة ب)

a. First stage
b. Second stage
c. Third stage
d. Fourth stage

12 In the second stage of labor, expulsion of the fetus from the birth canal depends on which important factor? تكرر في 2 مصادر كويز 2 – 2021 (نسخة ب)

a. Maternal bearing down
b. Cervical dilatation and effacement
c. Uterine contractions only
d. Adequate pelvic size

13 A mother is in the active deceleration phase. How frequently should the nurse assess her uterine contractions? كويز 2 – 2021 (نسخة ب)

a. Every 5 minutes
b. Every 15 minutes
c. Every 30 minutes
d. Every 60 minutes

14 In a multipara, the first stage of labor usually lasts: تكرر في 2 مصادر — كويز 2 – 2021 (نسخة ب)

a. 3 to 6 hours
b. 6 to 8 hours
c. 10 to 15 hours
d. 24 hours

15 A nullipara is admitted with cervix 6 cm, 70% effaced, at −1 station. What phase of labor is she in? تكرر في 2 مصادر — كويز 2 – 2021 (نسخة أ)

a. Active acceleration phase
b. Active deceleration phase
c. Latent phase
d. Expulsion phase

16 Which of the following statements is unrelated to the signs of placental separation? تكرر في 2 مصادر — كويز 2 – 2021 (نسخة أ)

a. Gush of blood
b. Uterus rises in the lower abdomen
c. Umbilical cord protrudes farther out of the vagina
d. Uterus becomes flaccid

17 To determine the duration of uterine contractions, the nurse notes the time from: تكرر في 2 مصادر — فاينال 2019–2020 (بنك الأسئلة)

a. The onset of the contraction until it ends
b. The beginning of one contraction until the end of the next
c. The onset of one contraction until the onset of the next
d. The end of one contraction until the end of the next

18 Bloody show is: تكرر في 2 مصادر — فاينال 2019–2020 (بنك الأسئلة)

a. A result of placenta previa
b. Not seen in breech presentations
c. A consequence of effacement and dilatation of the cervix
d. A sign of impending obstetric hemorrhage

19 Secondary forces acting to aid delivery include: تكرر في 2 مصادر — فاينال 2019–2020 (بنك الأسئلة)

a. Birth canal friction
b. Maternal bearing down
c. Cervical resistance
d. Uterine contractions

20 Mrs. A, a multigravida, has a cervix 7 cm dilated and 3 contractions per 10 minutes. She is in which phase? تكرر في 2 مصادر — فاينال 2019–2020 (بنك الأسئلة)

a. Latent
b. Active
c. Transition
d. Prodromal

21 Which assessment indicates the client is beginning the second stage of labor? تكرر في 2 مصادر — فاينال 2019–2020 (بنك الأسئلة)

a. The contractions are irregular
b. The cervix is 8 cm dilated
c. The spontaneous urge to push is initiated from perineal pressure
d. The membranes have ruptured

22 Show is considered a sure sign of normal labor. (True / False) أسئلة سابقة

a. True **b.** False

23 Labor is defined as a series of events by which the fetus and placenta are expelled from the woman's uterus. (True / False) أسئلة سابقة

a. True **b.** False

24 Which assessment finding indicates that the woman is beginning the second stage of labor? أسئلة سابقة

a. The spontaneous urge to push is initiated
b. The cervix is 8 cm dilated
c. The contractions are irregular
d. The membranes have ruptured

25 A nurse concludes that a primigravida is in the active deceleration (transition) phase without doing a vaginal examination. Which sign led to that conclusion? أسئلة سابقة

a. The woman vomited
b. The woman laughed at something
c. The woman was cooperative with the nurse
d. The woman fell asleep during a contraction

26 A laboring woman, G4 P3, who was 6 cm dilated 1 hour ago, cries to go to the bathroom. Which action is appropriate? أسئلة سابقة

a. Assess cervical dilation **b.** Help the woman to the bathroom
c. Check the fetal heart rate **d.** Ask if she needs pain medicine

27 At which stage of labor is the mother advised to avoid bearing down? تكرر في 2 مصادر أسئلة سابقة

a. First stage **b.** Third stage
c. Fourth stage **d.** Second stage

28 The nurse assesses the frequency of a laboring client's contractions by timing from the beginning of one contraction: أسئلة سابقة

a. Until it is completely over
b. To the end of a second contraction
c. To the beginning of the next contraction
d. Until the uterus becomes very firm

29 A client is admitted in early active labor. The priority nursing intervention on admission is: أسئلة سابقة

a. Auscultating the fetal heart
b. Taking an obstetric history
c. Asking when she last ate
d. Ascertaining whether membranes ruptured

30 **A G1 P0 client is 100% effaced, 3 cm dilated, and the fetus is at +1 station. The fetal head is:** أسئلة سابقة

a. Not yet engaged
b. Entering the pelvic inlet
c. Below the ischial spines
d. Visible at the vaginal opening

31 **Which observation indicates that the placenta has separated from the uterine wall and is ready for delivery?** أسئلة سابقة

a. The umbilical cord shortens and changes color
b. A soft and boggy uterus
c. Maternal complaints of severe cramping
d. Change in the shape of the uterus

32 **The nurse tells a client in early labor that effleurage is:** أسئلة سابقة

a. A form of biofeedback to enhance bearing down
b. Light stroking of the abdomen to facilitate relaxation
c. Pressure on the sacrum to relieve backache
d. Contracting a specific muscle group while others rest

33 **A client in the second stage of labor with contractions every 2 minutes cries out in pain with each contraction. The nurse recognizes this behavior as:** أسئلة سابقة

a. Exhaustion
b. Fear of losing control
c. Involuntary grunting
d. Valsalva's maneuver

34 **The fetus is documented at −1 station. The presenting part is:** أسئلة سابقة

a. 1 cm above the ischial spines
b. 1 fingerbreadth below the symphysis pubis
c. 1 inch below the coccyx
d. 1 inch below the iliac crest

35 **After delivery of the newborn, the nurse observes the umbilical cord lengthen and a spurt of blood from the vagina. The nurse documents these as signs of:** أسئلة سابقة

a. Hematoma
b. Placenta previa
c. Uterine atony
d. Placental separation

36 **A client in active labor with intact membranes is prepared for amniotomy. After this procedure she will most likely have:** أسئلة سابقة

a. Less pressure on her cervix
b. Increased efficiency of contractions
c. Decreased number of contractions
d. The need for increased BP monitoring

37 **The nurse determines that the client is beginning the second stage of labor when:** أسئلة سابقة

a. She begins to expel clear vaginal fluid
b. The contractions are regular
c. The membranes have ruptured
d. The cervix is completely dilated

38 Before initiating a prescribed IV oxytocin (Pitocin) infusion for a client in labor, the nurse ensures that which is implemented? أسئلة سابقة

a. Complete bed rest
b. Continuous electronic fetal monitoring
c. IV infusion of antibiotics
d. A code cart at the bedside

39 To determine the duration of uterine contractions, the nurse notes the time from: تكرر في 2 مصادر أسئلة سابقة

a. The onset of the contraction until it ends
b. The beginning of one contraction until the end of the next
c. The onset of one contraction until the onset of the next
d. The end of one contraction until the end of the next

40 Bloody show is: تكرر في 2 مصادر أسئلة سابقة

a. A result of placenta previa
b. Not seen in breech presentations
c. A consequence of effacement and dilatation of the cervix
d. A sign of impending obstetric hemorrhage

41 Secondary forces acting to aid delivery include: تكرر في 2 مصادر أسئلة سابقة

a. Birth canal friction
b. Maternal bearing down
c. Cervical resistance
d. Uterine contractions

42 Mrs. A, a multigravida, has a cervix 7 cm dilated and 3 contractions per 10 minutes. She is in which phase? تكرر في 2 مصادر أسئلة سابقة

a. Latent
b. Active
c. Transition
d. Prodromal

43 Which assessment indicates the client is beginning the second stage of labor? تكرر في 2 مصادر أسئلة سابقة

a. The contractions are irregular
b. The cervix is 8 cm dilated
c. The spontaneous urge to push is initiated from perineal pressure
d. The membranes have ruptured

44 Which of the following presentations is the most favorable for delivery? كويز طب نسا

a. Breech
b. Transverse
c. Cephalic
d. Oblique

45 Which of the following changes occurs in the cervix during labor? كويز طب نسا

a. Thickening
b. Effacement
c. Elongation
d. Firming

46 During which stage of labor does the cervix dilate? كويز طب نسا

a. First stage
b. Second stage
c. Third stage
d. Fourth stage

47 Which hormone is primarily responsible for initiating labor (stimulating uterine contractions)? كويز طب نسا

a. Estrogen
b. Progesterone
c. Oxytocin
d. Prolactin

Part B · NCLEX-Style Practice

أسئلة على طريقة NCLEX (إضافة)

N1 A primigravida is 5 cm dilated with regular contractions every 3–5 minutes. She is in: NCLEX

a. Latent phase
b. Active phase
c. Transition
d. Second stage

N2 The second stage of labor begins with: NCLEX

a. Rupture of membranes
b. Full cervical dilatation (10 cm)
c. Crowning
d. Placental separation

N3 Select ALL signs of placental separation. SELECT ALL

a. Sudden gush of blood
b. Lengthening of the cord
c. Uterus becomes globular and rises
d. Uterus becomes soft and boggy
e. Fundus descends into pelvis

N4 Membranes rupture and the fluid is green-tinged. The nurse should: NCLEX

a. Document as normal
b. Report to the provider and assess FHR
c. Encourage pushing
d. Give oxytocin

N5 Contraction frequency is measured from: NCLEX

a. Beginning to end of one contraction
b. Beginning of one contraction to beginning of the next
c. End of one to beginning of the next
d. Peak to peak only

N6 A woman feels the urge to push at 7 cm. The nurse instructs her to: NCLEX

a. Push hard with each contraction
b. Pant/blow during contractions to avoid bearing down
c. Hold her breath
d. Lie supine

N7 Station +2 means the presenting part is: NCLEX

a. 2 cm above the ischial spines
b. At the ischial spines
c. 2 cm below the ischial spines
d. Crowning

N8 In the fourth stage of labor, the fundus is boggy. The FIRST action is: NCLEX

a. Notify the provider
b. Massage the fundus
c. Check vital signs
d. Give fluids

Part C · Quizlet Flashcards

كروت مراجعة سريعة

غطّي العمود الأبيض بورقة واختبر نفسك: شوف المصطلح وقول التعريف.

Labor	Series of events expelling fetus, placenta and membranes	**First stage**	Onset of true labor → full dilatation
Latent phase	0–3 cm	**Active phase**	4–7 cm
Transition	8–10 cm; vomiting, irritability, urge to push	**Second stage**	Full dilatation → birth of baby
Third stage	Birth of baby → delivery of placenta	**Fourth stage**	First 1–2 hours after placenta
First stage duration	Primi 10–12 h; multi 6–8 h	**Primary power**	Uterine contractions
Secondary power	Maternal bearing down	**Station**	Relation of presenting part to ischial spines (− above, + below)
Signs of placental separation	Gush of blood, cord lengthens, uterus rises & becomes globular	**Show**	Blood-tinged mucus from effacement/dilatation
Avoid bearing down	First stage (before full dilatation)	**Frequency of contractions**	Beginning of one to beginning of next

Answer Key

الإجابات

1. a — 6 to 8 hours

2. a — First stage

3. d — Uterus becomes flaccid

4. a — 1 cm below the ischial spines

5. c — Green-tinged amniotic fluid

6. b — Every 15 minutes

7. a — Maternal bearing down

8. a — Active acceleration phase

9. b — First stage

10. b — Placental separation

11. a — First stage

12. a — Maternal bearing down

13. b — Every 15 minutes

14. b — 6 to 8 hours

15. a — Active acceleration phase

16. d — Uterus becomes flaccid

17. a — The onset of the contraction until it ends

18. c — A consequence of effacement and dilatation of the cervix

19. b — Maternal bearing down

20. b — Active

21. c — The spontaneous urge to push is initiated from perineal pressure

22. a — True

23. a — True

24. a — The spontaneous urge to push is initiated

25. a — The woman vomited

26. a — Assess cervical dilation

27. a — First stage

28. c — To the beginning of the next contraction

29. a — Auscultating the fetal heart

30. c — Below the ischial spines

31. d — Change in the shape of the uterus

32. b — Light stroking of the abdomen to facilitate relaxation

33. b — Fear of losing control

34. a — 1 cm above the ischial spines

35. d — Placental separation

36. b — Increased efficiency of contractions

37. d — The cervix is completely dilated

38. b — Continuous electronic fetal monitoring

39. a — The onset of the contraction until it ends

40. c — A consequence of effacement and dilatation of the cervix

41. b — Maternal bearing down

42. b — Active

43. c — The spontaneous urge to push is initiated from perineal pressure

44. c — Cephalic

45. b — Effacement

46. a — First stage

47. c — Oxytocin

N1. b — Active phase
Active phase 4–7 cm; transition 8–10 cm.

N2. b — Full cervical dilatation (10 cm)
2nd stage = full dilatation to birth of the baby.

N3. a, b, c — Sudden gush of blood; Lengthening of the cord; Uterus becomes globular and rises
A boggy uterus indicates atony, not separation.

N4. b — Report to the provider and assess FHR
Meconium-stained fluid may indicate fetal distress.

N5. b — Beginning of one contraction to beginning of the next
Duration = beginning to end of one contraction; frequency = beginning to beginning.

N6. b — Pant/blow during contractions to avoid bearing down
Pushing before full dilatation causes cervical edema and lacerations.

N7. c — 2 cm below the ischial spines
Minus = above spines, 0 = engaged at spines, plus = below.

N8. b — Massage the fundus
Fundal massage stimulates contraction to control bleeding.