

Day 7: Calculus Fundamentals, Derivatives II

Jean Clipperton

Math Camp 2022

Day 7: Concept Agenda

- ▶ More Derivatives!
 - ▶ Higher order derivatives
 - ▶ Inflection points– minima and maxima
 - ▶ Partial derivatives– what to do when there are 2+ variables
 - ▶ Advanced rules of derivatives: product, quotient, chain rules and more

Fundamentals of Derivatives: Review

Yesterday we went over the basics of derivatives, what they are conceptually, and how to solve for them.

Let's rehash these problems from yesterday to warm up:

- ▶ $f(x) = 5$
- ▶ $f(x) = 3x - 7$
- ▶ $f(x) = 3x^2$
- ▶ $f(x) = \frac{x^2}{x}$
- ▶ $f(s) = s^{-2}$
- ▶ $f(y) = y(y + 7)(y - 3)$
- ▶ $f(z) = \frac{z^2 - 5z - 6}{z + 1}$

Fundamentals of Derivatives: Review

Yesterday we went over the basics of derivatives, what they are conceptually, and how to solve for them.

Let's rehash these problems from yesterday to warm up:

- ▶ $f(x) = 5, f'(x) = 0$
- ▶ $f(x) = 3x - 7, f'(x) = 3$
- ▶ $f(x) = 3x^2, f'(x) = 6x$
- ▶ $f(x) = \frac{x^2}{x}, f'(x) = 1$
- ▶ $f(s) = s^{-2}, f'(s) = -2s^{-3}$ (not continuous at $s = 0$)
- ▶ $f(y) = y(y + 7)(y - 3), f'(y) = 3y^2 + 8y - 21$
- ▶ $f(z) = \frac{z^2 - 5z - 6}{z + 1}, f'(z) = 1$ (not continuous at $z = -1$)

Fundamentals of Derivatives: Review

Moving on from these basic rules, we will now cover higher order derivatives, partial derivatives, and more advanced (but still basic) rules of derivatives, such as the product rule, quotient rule, and chain rule.

Higher Order Derivatives

Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

► $f(x) = x^5 + 3x^3 + 2x + 8$

Higher Order Derivatives

Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

- ▶ $f(x) = x^5 + 3x^3 + 2x + 8$

- ▶ $f'(x) = 5x^4 + 9x^2 + 2$

Higher Order Derivatives


Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

- ▶ $f(x) = x^5 + 3x^3 + 2x + 8$
- ▶ $f'(x) = 5x^4 + 9x^2 + 2$
- ▶ $f''(x) = 20x^3 + 18x$

Critical Points

Critical points occur where the derivative is zero. We can find them by graphing (ocular method) or plugging in values after calculating the derivative.



Critical Points

When the derivative is zero: can be local max or min

Try:

- ▶ $x^2 + 4x$: max or min? Where is the critical point?

Critical Points

When the derivative is zero: can be local max or min

Try:

- ▶ $x^2 + 4x$: max or min? Where is the critical point? Min, critical point at $x = -2$

Critical Points

When the derivative is zero: can be local max or min

Try:

- ▶ $x^2 + 4x$: max or min? Where is the critical point? Min, critical point at $x = -2$
- ▶ $-x^2 + 4x$: max or min? Where is the critical point?

Critical Points

When the derivative is zero: can be local max or min

Try:

- ▶ $x^2 + 4x$: max or min? Where is the critical point? Min, critical point at $x = -2$
- ▶ $-x^2 + 4x$: max or min? Where is the critical point? Max, critical point at $x = 2$

Sometimes there are multiple critical points

- ▶ How many?
- ▶ How to find?

Sometimes there are multiple critical points

- ▶ How many? (max) number of critical points is the highest degree of the derivative (same as finding zeroes)
- ▶ How to find?

Sometimes there are multiple critical points

- ▶ How many? (max) number of critical points is the highest degree of the derivative (same as finding zeroes)
- ▶ How to find? take the derivative and set to zero

Sometimes there are multiple critical points

- ▶ How many? (max) number of critical points is the highest degree of the derivative (same as finding zeroes)
- ▶ How to find? take the derivative and set to zero

Try: $f(x) = x^4 - 16x^2$

Critical Points, pt. 2

Sometimes there are multiple critical points

- ▶ How many? (max) number of critical points is the highest degree of the derivative (same as finding zeroes)
- ▶ How to find? take the derivative and set to zero

Try: $f(x) = x^4 - 16x^2$

(up to) Three critical points. $f'(x) = 4x^3 - 16x = 4x(x^2 - 4)$.

Critical Points, pt. 2

Sometimes there are multiple critical points

- ▶ How many? (max) number of critical points is the highest degree of the derivative (same as finding zeroes)
- ▶ How to find? take the derivative and set to zero

Try: $f(x) = x^4 - 16x^2$

(up to) Three critical points. $f'(x) = 4x^3 - 16x = 4x(x^2 - 4)$.

Zeroes at -2, 0, 2

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

THIS IS IMPORTANT FOR INTERACTION TERMS!

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

THIS IS IMPORTANT FOR INTERACTION TERMS!

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z$$

$$f(x, y) = x + 4y$$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

THIS IS IMPORTANT FOR INTERACTION TERMS!

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z$$

$$f(x, y) = x + 4y$$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

THIS IS IMPORTANT FOR INTERACTION TERMS!

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z \quad \partial_x = 7z + 8x$$

$$f(x, y) = x + 4y \quad \partial_x = 1$$

Partial derivatives show how rate of change moves with another variable. What is expected change of Y in relation to X ?

$$Y = \beta_0 + \beta_1 X + \beta_2 Z + \beta_3 XZ + \epsilon$$

Partial derivatives show how rate of change moves with another variable. What is expected change of Y in relation to X ?

$$Y = \beta_0 + \beta_1 X + \beta_2 Z + \beta_3 XZ + \epsilon \quad \partial_X = \beta_1 + \beta_3 Z$$

Partial derivatives show how rate of change moves with another variable. What is expected change of Y in relation to X ?

$$Y = \beta_0 + \beta_1 X + \beta_2 Z + \beta_3 XZ + \epsilon \quad \partial_X = \beta_1 + \beta_3 Z$$

How is that different from just β_1 or just β_3 ?

The effect is $\partial_X = \beta_1 + \beta_3 Z$. Now, suppose Z can be 0 or 1.

The effect is $\partial_X = \beta_1 + \beta_3 Z$. Now, suppose Z can be 0 or 1.

Understanding Interactions

$$Y = \beta_0 + \beta_1 X + \beta_2 Z + \beta_3 XZ + \varepsilon$$

Hypothesis H_1 : An increase in X is associated with an increase in Y when condition Z is met, but not when condition Z is absent.




Fig. 1 A graphical illustration of an interaction model consistent with hypothesis H_1 .

(Additional) Rules for derivatives

So far, we've just had simple functions but what if we are multiplying, dividing, or have an otherwise-more-advanced function?

(Additional) Rules for derivatives

So far, we've just had simple functions but what if we are multiplying, dividing, or have an otherwise-more-advanced function?

- ▶ Product Rule: $f(x) * g(x)$
- ▶ Quotient Rule: $\frac{f(x)}{g(x)}$
- ▶ Chain Rule: $f(g(x))$
- ▶ Other: eg, exponentials: e^x , $\ln(x)$

(Additional) Rules for derivatives

So far, we've just had simple functions but what if we are multiplying, dividing, or have an otherwise-more-advanced function?

- ▶ Product Rule: $f(x) * g(x)$, $x^3 * x^2$
- ▶ Quotient Rule: $\frac{f(x)}{g(x)}$, $\frac{x^4+3x}{x^2}$
- ▶ Chain Rule: $f(g(x))$, $(x^2 + 1)^3$ (composition!)
- ▶ Other: eg, exponentials: e^x , $\ln(x)$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example:** $(3x+4)(x+2)$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example: $(3x+4)(x+2)$** To take the derivative using the product rule, we do the following:
$$f'(x) * g(x) + g'(x)f(x).$$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example: $(3x+4)(x+2)$** To

take the derivative using the product rule, we do the following:

$f'(x) * g(x) + g'(x)f(x)$. For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example: $(3x+4)(x+2)$** To

take the derivative using the product rule, we do the following:

$f'(x) * g(x) + g'(x)f(x)$. For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$. Then, we substitute these in:

$$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example: $(3x+4)(x+2)$** To

take the derivative using the product rule, we do the following:

$f'(x) * g(x) + g'(x)f(x)$. For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$. Then, we substitute these in:

$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$ Simplify to get:
 $3x + 6 + 3x + 4 = 6x + 10$.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule. **Example: $(3x+4)(x+2)$** To

take the derivative using the product rule, we do the following:

$f'(x) * g(x) + g'(x)f(x)$. For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$. Then, we substitute these in:

$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$ Simplify to get:
 $3x + 6 + 3x + 4 = 6x + 10$.

To take the derivative using our previous approach, we first multiply:
 $3x^2 + 4x + 6x + 8 = 3x^2 + 10x + 8$. Then, just take the derivative of each term: $f'(x) = 6x + 10$.

WHY DO WITH THE PRODUCT RULE??

Product Rule

Suppose that instead you had $(3x^2 + 3x + 4)(x^3 + 2x^2 + x + 2)$.
Now the product rule is looking a little nicer!

Product Rule

Suppose that instead you had $(3x^2 + 3x + 4)(x^3 + 2x^2 + x + 2)$.
Now the product rule is looking a little nicer!

$$f(x) = (3x^2 + 3x + 4) \text{ and } g(x) = (x^3 + 2x^2 + x + 2)$$

Product Rule

Suppose that instead you had $(3x^2 + 3x + 4)(x^3 + 2x^2 + x + 2)$.
Now the product rule is looking a little nicer!

$$f(x) = (3x^2 + 3x + 4) \text{ and } g(x) = (x^3 + 2x^2 + x + 2)$$
$$f'(x) = 6x + 3 \text{ and } g'(x) = 3x^2 + 4x + 1.$$

We can substitute this into the formula: $f'(x) * g(x) + g'(x)f(x)$

Product Rule

Suppose that instead you had $(3x^2 + 3x + 4)(x^3 + 2x^2 + x + 2)$.
Now the product rule is looking a little nicer!

$$f(x) = (3x^2 + 3x + 4) \text{ and } g(x) = (x^3 + 2x^2 + x + 2)$$
$$f'(x) = 6x + 3 \text{ and } g'(x) = 3x^2 + 4x + 1.$$

We can substitute this into the formula: $f'(x) * g(x) + g'(x)f(x)$
 $(6x + 3)(x^3 + 2x^2 + x + 2) + (3x^2 + 4x + 1)(3x^2 + 3x + 4)$. This is
a mess – but you have your answer at least (and much easier than
doing it the long way)!

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Example: $\frac{3x^2}{x+2}$. Formula is $\frac{f'(x)*g(x)-g'(x)f(x)}{(g(x))^2}$

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Example: $\frac{3x^2}{x+2}$. Formula is $\frac{f'(x) \cdot g(x) - g'(x) \cdot f(x)}{(g(x))^2}$ So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Example: $\frac{3x^2}{x+2}$. Formula is $\frac{f'(x) \cdot g(x) - g'(x) \cdot f(x)}{(g(x))^2}$ So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$ $f'(x) = 6x$ and $g'(x) = 1$. Plug in to get

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Example: $\frac{3x^2}{x+2}$. Formula is $\frac{f'(x) \cdot g(x) - g'(x) \cdot f(x)}{(g(x))^2}$ So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$ $f'(x) = 6x$ and $g'(x) = 1$. Plug in to get $\frac{6x(x+2) - 1(3x^2)}{(x+2)^2} = \frac{6x^2 + 12x - 3x^2}{(x+2)^2} = \frac{3x^2 + 12x}{(x+2)^2}$

Practice: $\frac{x-4}{x+5} \cdot \frac{3x^3}{x+2}$

Quotient Rule

There are two ways to think about the quotient rule: a) you have something divided by something else or b) you have something multiplied by something to a negative power (chain rule, next up!)

Example: $\frac{3x^2}{x+2}$. Formula is $\frac{f'(x) \cdot g(x) - g'(x) \cdot f(x)}{(g(x))^2}$ So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$ $f'(x) = 6x$ and $g'(x) = 1$. Plug in to get $\frac{6x(x+2) - 1(3x^2)}{(x+2)^2} = \frac{6x^2 + 12x - 3x^2}{(x+2)^2} = \frac{3x^2 + 12x}{(x+2)^2}$

Practice: $\frac{x-4}{x+5} \cdot \frac{3x^3}{x+2}$ **Ans:** $\frac{9}{(x+5)^2} \cdot \frac{6x^3 + 18x^2}{(x+2)^2}$

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$.

We can use the chain rule to evaluate this.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$.

We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$. We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$. So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$. We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$. So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$. We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$. So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$. We substitute in to get: $3(x + 3)^2 * 1$.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$. We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$. So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$. We substitute in to get: $3(x + 3)^2 * 1$.

Try: $f(x) = (2x^2 + 8x)^4$ $f(x) = (9x - x^2)^6$

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$. We can use the chain rule to evaluate this. What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$. So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$. We substitute in to get: $3(x + 3)^2 * 1$.

Try: $f(x) = (2x^2 + 8x)^4$ $f(x) = (9x - x^2)^6$

Ans $4(4x + 8)(2x^2 + 8x)^3$

$6(9 - 2x)(9x - x^2)^5 = (54 - 12x)(9x - 2x^2)^5$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

You can make these more complicated by including a function of x . How would we take the derivative in that case?

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

EX: $\ln(3x)$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{g(x)} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

$$\text{EX: } \ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

$$\text{EX: } \ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$$

$$\text{TRY: } \ln(x^2), e^{2x}$$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \quad (\text{a favorite of mine})$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \quad (\text{used rarely, if ever})$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \quad (\text{rarely used})$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

$$\text{EX: } \ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$$

$$\text{TRY: } \ln(x^2), e^{2x} \quad \text{ANS: } \frac{2}{x} \text{ and } 2e^{2x}$$

Additional Resources

- ▶ Daniel Kleitman's Calculus for Beginners and Artists:
www-math.mit.edu/~djk/calculus_beginners
- ▶ Dan Sloughter (online textbook): math.furman.edu/~dcs/book
- ▶ Calc refresher (Harvey Mudd Calc Tutorial):
www.math.hmc.edu/calculus/tutorials/

Derivatives in Review

We can use derivatives to calculate rates of change and we're concerned how functions behave. We have a series of rules that can help us get there, even if we have multiple variables.

Derivatives in Review

We can use derivatives to calculate rates of change and we're concerned how functions behave. We have a series of rules that can help us get there, even if we have multiple variables.

Table 6.1: List of Rules of Differentiation

Sum rule	$(f(x) + g(x))' = f'(x) + g'(x)$
Difference rule	$(f(x) - g(x))' = f'(x) - g'(x)$
Multiply by constant rule	$f'(ax) = af'(x)$
Product rule	$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$
Quotient rule	$\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$
Chain rule	$(g(f(x)))' = g'(f(x))f'(x)$
Inverse function rule	$(f^{-1}(x))' = \frac{1}{f'(f^{-1}(x))}$
Constant rule	$(a)' = 0$
Power rule	$(x^n)' = nx^{n-1}$
Exponential rule 1	$(e^x)' = e^x$
Exponential rule 2	$(a^x)' = a^x(\ln(a))$
Logarithm rule 1	$(\ln(x))' = \frac{1}{x}$
Logarithm rule 2	$(\log_a(x))' = \frac{1}{x(\ln(a))}$
Trigonometric rules	$(\sin(x))' = \cos(x)$ $(\cos(x))' = -\sin(x)$ $(\tan(x))' = 1 + \tan^2(x)$
Piecewise rules	Treat each piece separately

Derivatives in Review

We can use derivatives to calculate rates of change and we're concerned how functions behave. We have a series of rules that can help us get there, even if we have multiple variables.

Table 6.1: List of Rules of Differentiation

Sum rule	$(f(x) + g(x))' = f'(x) + g'(x)$
Difference rule	$(f(x) - g(x))' = f'(x) - g'(x)$
Multiply by constant rule	$f'(ax) = af'(x)$
Product rule	$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$
Quotient rule	$\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$
Chain rule	$(g(f(x)))' = g'(f(x))f'(x)$
Inverse function rule	$(f^{-1}(x))' = \frac{1}{f'(f^{-1}(x))}$
Constant rule	$(a)' = 0$
Power rule	$(x^n)' = nx^{n-1}$
Exponential rule 1	$(e^x)' = e^x$
Exponential rule 2	$(a^x)' = a^x(\ln(a))$
Logarithm rule 1	$(\ln(x))' = \frac{1}{x}$
Logarithm rule 2	$(\log_a(x))' = \frac{1}{x(\ln(a))}$
Trigonometric rules	$(\sin(x))' = \cos(x)$ $(\cos(x))' = -\sin(x)$ $(\tan(x))' = 1 + \tan^2(x)$
Piecewise rules	Treat each piece separately

However, there comes a time where we have the rate of change but we need to know what the original is.

Derivatives in Review

We can use derivatives to calculate rates of change and we're concerned how functions behave. We have a series of rules that can help us get there, even if we have multiple variables.

Table 6.1: List of Rules of Differentiation

Sum rule	$(f(x) + g(x))' = f'(x) + g'(x)$
Difference rule	$(f(x) - g(x))' = f'(x) - g'(x)$
Multiply by constant rule	$f'(ax) = af'(x)$
Product rule	$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$
Quotient rule	$\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$
Chain rule	$(g(f(x)))' = g'(f(x))f'(x)$
Inverse function rule	$(f^{-1}(x))' = \frac{1}{f'(f^{-1}(x))}$
Constant rule	$(a)' = 0$
Power rule	$(x^n)' = nx^{n-1}$
Exponential rule 1	$(e^x)' = e^x$
Exponential rule 2	$(a^x)' = a^x(\ln(a))$
Logarithm rule 1	$(\ln(x))' = \frac{1}{x}$
Logarithm rule 2	$(\log_a(x))' = \frac{1}{x(\ln(a))}$
Trigonometric rules	$(\sin(x))' = \cos(x)$ $(\cos(x))' = -\sin(x)$ $(\tan(x))' = 1 + \tan^2(x)$
Piecewise rules	Treat each piece separately

However, there comes a time where we have the rate of change but we need to know what the original is. Next, we'll cover integrals – undoing derivatives!