

Day 4 AM: Algebra Review

Jean Clipperton

Math Camp 2021

Questions?

Concluding CH 1 of Moore & Siegel – moving to Ch 2

Algebra Review

Properties

- ▶ **Associative property:** $(a + b) + c = a + (b + c)$ and $(a * b) * c = a * (b * c)$
- ▶ **Commutative property** $a + b = b + a$ and $a * b = b * a$
- ▶ **Distributive property** $a(b + c) = ab + ac$
- ▶ **Identity property** $x + 0 = x$ and $x * 1 = x$
- ▶ **Inverse property** $-x + x = 0$. Multiplicative inverse exists, but not for all numbers $x^{-1} * x = 1$

Factoring

We may need to break down different functions.

1. $x^2 + 3x - 4 =$

2. $m^2 + 3m + 2 =$

3. $x^2 + 5x + 6 =$

4. $25 - x^2 =$

Factoring

We may need to break down different functions.

1. $x^2 + 3x - 4 = (x - 1)(x - 4)$

2. $m^2 + 3m + 2 =$

3. $x^2 + 5x + 6 =$

4. $25 - x^2 =$

Factoring

We may need to break down different functions.

1. $x^2 + 3x - 4 = (x - 1)(x - 4)$

2. $m^2 + 3m + 2 = (m + 1)(m + 2)$

3. $x^2 + 5x + 6 =$

4. $25 - x^2 =$

Factoring

We may need to break down different functions.

1. $x^2 + 3x - 4 = (x - 1)(x - 4)$

2. $m^2 + 3m + 2 = (m + 1)(m + 2)$

3. $x^2 + 5x + 6 = (x + 3)(x + 2)$

4. $25 - x^2 =$

Factoring

We may need to break down different functions.

$$1. \ x^2 + 3x - 4 = (x - 1)(x - 4)$$

$$2. \ m^2 + 3m + 2 = (m + 1)(m + 2)$$

$$3. \ x^2 + 5x + 6 = (x + 3)(x + 2)$$

$$4. \ 25 - x^2 = (5 - x)(5 + x)$$

Factoring

We may need to break down different functions.

$$1. \ x^2 + 3x - 4 = (x - 1)(x - 4)$$

$$2. \ m^2 + 3m + 2 = (m + 1)(m + 2)$$

$$3. \ x^2 + 5x + 6 = (x + 3)(x + 2)$$

$$4. \ 25 - x^2 = (5 - x)(5 + x)$$

Try:

$$1. \ -b^2 + 9a^4$$

$$2. \ 2x^2 + 7x + 3$$

$$3. \ 6y^2 + y - 2$$

Factoring

We may need to break down different functions.

$$1. \ x^2 + 3x - 4 = (x - 1)(x - 4)$$

$$2. \ m^2 + 3m + 2 = (m + 1)(m + 2)$$

$$3. \ x^2 + 5x + 6 = (x + 3)(x + 2)$$

$$4. \ 25 - x^2 = (5 - x)(5 + x)$$

Try:

$$1. \ -b^2 + 9a^4 \ (-b + 3a^2)(b + 3a^2)$$

$$2. \ 2x^2 + 7x + 3$$

$$3. \ 6y^2 + y - 2$$

Factoring

We may need to break down different functions.

$$1. \ x^2 + 3x - 4 = (x - 1)(x - 4)$$

$$2. \ m^2 + 3m + 2 = (m + 1)(m + 2)$$

$$3. \ x^2 + 5x + 6 = (x + 3)(x + 2)$$

$$4. \ 25 - x^2 = (5 - x)(5 + x)$$

Try:

$$1. \ -b^2 + 9a^4 \ (-b + 3a^2)(b + 3a^2)$$

$$2. \ 2x^2 + 7x + 3 \ (2x + 1)(x + 3)$$

$$3. \ 6y^2 + y - 2$$

Factoring

We may need to break down different functions.

$$1. \ x^2 + 3x - 4 = (x - 1)(x - 4)$$

$$2. \ m^2 + 3m + 2 = (m + 1)(m + 2)$$

$$3. \ x^2 + 5x + 6 = (x + 3)(x + 2)$$

$$4. \ 25 - x^2 = (5 - x)(5 + x)$$

Try:

$$1. \ -b^2 + 9a^4 = (-b + 3a^2)(b + 3a^2)$$

$$2. \ 2x^2 + 7x + 3 = (2x + 1)(x + 3)$$

$$3. \ 6y^2 + y - 2 = (2y - 1)(3y + 2)$$

Inequalities

Relations: Intervals & Inequalities

Interval notation can be used to express ranges of numbers :

$[a, b]$ $a \leq x \leq b$ Square brackets include end points (closed interval)

(a, b) $a < x < b$ Parenthesis mean exclude end points (open interval)

$\{a, b\}$ Typically used for sets – not inequalities/intervals

Relations

Graph the following:

▶ $4 < x$

▶ $y > 12$

▶ $3 < z < 7$

▶ $(3, 9)$

▶ $[-7, 2)$

Solving Inequalities

Solve like regular equations but FLIP inequality when multiplying by negative values.

EX:

$$\blacktriangleright 3x < 4x + 2$$

Solving Inequalities

Solve like regular equations but FLIP inequality when multiplying by negative values.

EX:

- ▶ $3x < 4x + 2$

- ▶ $60x > -10(x + 7)$

Solving Inequalities

Solve like regular equations but FLIP inequality when multiplying by negative values.

EX:

- ▶ $3x < 4x + 2$

- ▶ $60x > -10(x + 7)$

Try:

- ▶ $-6(x + 8) < 12$

Absolute Value

Solve for TWO possibilities: quantity is positive or negative. EX:

$$|x - 3| > 4$$

- ▶ Quantity is positive: drop bars, solve like usual:

$$x - 3 > 4, x > 7$$

- ▶ Quantity is negative: then, really have $x - 3 < -4$ Solve to find $x < -1$

Absolute Value

Solve for TWO possibilities: quantity is positive or negative. EX:

$$|x - 3| > 4$$

- ▶ Quantity is positive: drop bars, solve like usual:

$$x - 3 > 4, x > 7$$

- ▶ Quantity is negative: then, really have $x - 3 < -4$ Solve to find $x < -1$

Solution: $x < -1$ OR $x > 7$

Absolute Value

Solve for TWO possibilities: quantity is positive or negative. EX:

$$|x - 3| > 4$$

- ▶ Quantity is positive: drop bars, solve like usual:

$$x - 3 > 4, x > 7$$

- ▶ Quantity is negative: then, really have $x - 3 < -4$ Solve to find $x < -1$

Solution: $x < -1$ OR $x > 7$ Try: $|2x - 3| > 9$

Factorial

The factorial, $!$, multiplies a number by each subsequent number down to one.

For example, $4! = 4 * 3 * 2 * 1$

You can also divide and multiply factorials: $\frac{3!}{4!} = \frac{3*2*1}{4*3*2*1} = \frac{1}{4}$

CANNOT ADD THE NUMBERS!! (e.g. $6!3! \neq 9!$)

Combinatorics: Combining elements

We can use factorials to help us understand ways of combining elements: e.g. suppose you are forming a committee of 3 people from a group of 5. How many ways can we do that?

Combinatorics: Combining elements

We can use factorials to help us understand ways of combining elements: e.g. suppose you are forming a committee of 3 people from a group of 5. How many ways can we do that?

Say these people are named A, B, C, D, and E.

Then we can have the following committees: ABC, ABD, ABE, ACD, ACE, ADE, BCD, BCE, BDE, CDE, for a total of ten configurations.

Combinatorics: Combining elements

We can use factorials to help us understand ways of combining elements: e.g. suppose you are forming a committee of 3 people from a group of 5. How many ways can we do that?

Say these people are named A, B, C, D, and E.

Then we can have the following committees: ABC, ABD, ABE, ACD, ACE, ADE, BCD, BCE, BDE, CDE, for a total of ten configurations.

We could also calculate them using the binomial function:

$$\binom{5}{3} = \frac{5!}{3!2!} = \frac{5*4}{2} = 10$$

Try $\binom{4}{2}$ with population A,B,C,D. Using the general formula:

$$\binom{N}{k} = \frac{N!}{k!(N-k)!}$$

Combinatorics: Combining elements

We can use factorials to help us understand ways of combining elements: e.g. suppose you are forming a committee of 3 people from a group of 5. How many ways can we do that?

Say these people are named A, B, C, D, and E.

Then we can have the following committees: ABC, ABD, ABE, ACD, ACE, ADE, BCD, BCE, BDE, CDE, for a total of ten configurations.

We could also calculate them using the binomial function:

$$\binom{5}{3} = \frac{5!}{3!2!} = \frac{5*4}{2} = 10$$

Try $\binom{4}{2}$ with population A,B,C,D. Using the general formula:

$$\binom{N}{k} = \frac{N!}{k!(N-k)!} \text{ (Answer: 6)}$$

WHY DOES THIS MATTER AT ALL????

WHY DOES THIS MATTER AT ALL????

You may have different populations and different treatments – how many different groups would you need to test the possible combinations?

Functions and Relations

Relations allow comparison of variables and expressions – some may be more or less specific in how they assign or specify these relationships between the *range* (y) and *domain* (x). Suppose we have a domain as follows { apple, kiwi, lime } and a range of { 4, 5 }.

Functions and Relations

Suppose we have a domain as follows { apple, kiwi, lime } and a range of { 4, 5}.

Functions

Functions assign *one* element of the range to each element of the domain.

Functions and Relations

Suppose we have a domain as follows { apple, kiwi, lime } and a range of { 4, 5}.

Functions

Functions assign *one* element of the range to each element of the domain. EX: apple \rightarrow 5, kiwi \rightarrow 4, lime \rightarrow 4.
(Exactly one y per x, but y can be assigned to multiple x)

Correspondences

Correspondences assign a subset of the range to each element of the domain.

Functions and Relations

Suppose we have a domain as follows $\{ \text{apple, kiwi, lime} \}$ and a range of $\{ 4, 5 \}$.

Functions

Functions assign *one* element of the range to each element of the domain. EX: $\text{apple} \rightarrow 5$, $\text{kiwi} \rightarrow 4$, $\text{lime} \rightarrow 4$.
(Exactly one y per x , but y can be assigned to multiple x)

Correspondences

Correspondences assign a subset of the range to each element of the domain. EX: $\text{apple} \rightarrow \{4, 5\}$. (Can have multiple y assigned to each x)

Functions: different mappings

We can visualize mappings either on a plane, or, even more simply, as connections between two sets:

Functions: different mappings

We can visualize mappings either on a plane, or, even more simply, as connections between two sets:




Figure 1:

Functions: the vertical line test

Functions can only assign one x to one y (and not one x to multiple y values) – otherwise it is a relation.

Functions: the vertical line test

Functions can only assign one x to one y (and not one x to multiple y values) – otherwise it is a relation. Can use the vertical line test to check:




Figure 2: "Vertical line test"

Functions

Functions specify the relationship between x and y , e.g. $y = a + bx$:
we often talk about *mapping* something onto something else.

Functions

Functions specify the relationship between x and y , e.g. $y = a + bx$: we often talk about *mapping* something onto something else.

Functions map from the domain to the codomain. Note that there is a distinction between the codomain and the range: codomain are possible values (e.g. natural numbers) and range are only the values reached/obtained. (sometimes called the image).

Functions

Functions specify the relationship between x and y , e.g. $y = a + bx$: we often talk about *mapping* something onto something else.

Functions map from the domain to the codomain. Note that there is a distinction between the codomain and the range: codomain are possible values (e.g. natural numbers) and range are only the values reached/obtained. (sometimes called the image).

You can also combine functions, e.g. $f(x) = x$ and $g(x) = x^3$ can be combined to produce $f \circ g(x) = f(g(x))$.

Function Terms

Table 3.1 from book (pg 49)

| Term | Meaning |
|-------------------------|--|
| Identity function | Elements in domain are mapped to identical elements in codomain |
| Inverse function | Function that when composed with original function returns identity function |
| Surjective (onto) | Every value in codomain produced by value in domain |
| Injective (one-to-one) | Each value in range comes from only one value in domain |
| Bijjective (invertible) | Both surjective and injective; function has an inverse |

Function Terms

Table 3.1 from book (pg 49)

| Term | Meaning |
|-------------------------|--|
| Identity function | Elements in domain are mapped to identical elements in codomain |
| Inverse function | Function that when composed with original function returns identity function |
| Surjective (onto) | Every value in codomain produced by value in domain |
| Injective (one-to-one) | Each value in range comes from only one value in domain |
| Bijjective (invertible) | Both surjective and injective; function has an inverse |

Why do we care?

Function Terms

We care about whether a function is surjective, injective, or bijective because we will know if we can trace back what produced what we have.

<https://www.mathsisfun.com/sets/injective-surjective-bijective.html>

Function Terms

- ▶ We can see that both functions are surjective – each ‘covers’ the codomains
- ▶ In the case of $g(x)$, we can see that it is injective, while $f(x)$ is not
- ▶ Thus, $f(x)$ is not bijective but $g(x)$ is (both surjective and injective) – thus, it is invertible (to be defined)

Composite Functions

Inner/Outer Functions: We begin at these two functions $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. You essentially substitute the second function as 'x'.

Composite Functions

Inner/Outer Functions: We begin at these two functions $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. You essentially substitute the second function as 'x'.

- Consider $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. Combine them to create $f \circ g(x) = f(g(x)) = (x^2 + 5x + 1)^5$

Composite Functions

Inner/Outer Functions: We begin at these two functions $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. You essentially substitute the second function as 'x'.

- ▶ Consider $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. Combine them to create $f \circ g(x) = f(g(x)) = (x^2 + 5x + 1)^5$
- ▶ Similarly, $f(x) = x^{0.5}$ and $g(x) = x(5x - 1)$ produce $f \circ g(x) = f(g(x)) = \sqrt{5x^2 - x}$

Composite functions, $f \circ g(x) = f(g(x))$ and $g \circ f(x) = g(f(x))$

Try: $f(x) = x^2 + 1$, $g(x) = 2x$

Composite Functions

Inner/Outer Functions: We begin at these two functions $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. You essentially substitute the second function as 'x'.

- ▶ Consider $f(x) = x^5$ and $g(x) = x^2 + 5x + 1$. Combine them to create $f \circ g(x) = f(g(x)) = (x^2 + 5x + 1)^5$
- ▶ Similarly, $f(x) = x^{0.5}$ and $g(x) = x(5x - 1)$ produce $f \circ g(x) = f(g(x)) = \sqrt{5x^2 - x}$

Composite functions, $f \circ g(x) = f(g(x))$ and $g \circ f(x) = g(f(x))$

Try: $f(x) = x^2 + 1$, $g(x) = 2x$

- ▶ $f \circ g(x) = 4x^2 + 1$
- ▶ $g \circ f(x) = 2x^2 + 2$

Function Terms: Invertible functions

When you compose a function with its inverse, you get the identity function.

Function Terms: Invertible functions

When you compose a function with its inverse, you get the identity function.

Example: suppose $f(x)=3x$ and $g(x)=x/3$. Find $f \circ g(x) = f(g(x))$

Function Terms: Invertible functions

When you compose a function with its inverse, you get the identity function.

Example: suppose $f(x)=3x$ and $g(x)=x/3$. Find $f \circ g(x) = f(g(x))$

In this case, we get $f(g(x)) = 3 * (x/3) = x$, where x is the identity function.

Function Terms: Invertible functions

When you compose a function with its inverse, you get the identity function.

Example: suppose $f(x)=3x$ and $g(x)=x/3$. Find $f \circ g(x) = f(g(x))$

In this case, we get $f(g(x)) = 3 * (x/3) = x$, where x is the identity function.

We can see that we only get one y value for each x and that, depending on the domain/codomain, the function is surjective as well, making this function bijective and have a legitimate inverse.

Monotonic Function Terms

| Term | Meaning |
|-----------------------|--|
| Increasing | Function increases on subset of domain |
| Decreasing | Function decreases on subset of domain |
| Strictly increasing | Function always increases on subset of domain |
| Strictly decreasing | Function always decreases on subset of domain |
| Weakly increasing | Function does not decrease on subset of domain |
| Weakly decreasing | Function does not increase on subset of domain |
| (Strict) monotonicity | Order preservation; function (strictly) increasing or decreasing over domain |

Table 3.2 from book (pg 51)

Monotonic Function Terms

| Term | Meaning |
|-----------------------|--|
| Increasing | Function increases on subset of domain |
| Decreasing | Function decreases on subset of domain |
| Strictly increasing | Function always increases on subset of domain |
| Strictly decreasing | Function always decreases on subset of domain |
| Weakly increasing | Function does not decrease on subset of domain |
| Weakly decreasing | Function does not increase on subset of domain |
| (Strict) monotonicity | Order preservation; function (strictly) increasing or decreasing over domain |

Table 3.2 from book (pg 51)

This is useful for math land but also for theory building: how are x and y related? Does more x ALWAYS mean more y (function is monotonically increasing, etc)?

Linear Equation vs Linear Function

Your beloved friend: $y = a + bx$ (AKA $y = mx + b$ AKA affine function).

Linear Equation vs Linear Function

Your beloved friend: $y = a + bx$ (AKA $y = mx + b$ AKA affine function).

We have the **intercept** (where the line crosses the y axis) and the **slope** (unit increase in iv related to dv). This is the 'plain vanilla' version.

The **linear function** is much more expansive: includes multiple variables, exponents, and logs (logarithms).

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$
- ▶ Simplify bases when same exponent: $x^4 * y^4 = (xy)^4$ and
 $m^{-2}n^2 = (m^{-1}n)^2$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$
- ▶ Simplify bases when same exponent: $x^4 * y^4 = (xy)^4$ and
 $m^{-2}n^2 = (m^{-1}n)^2$
- ▶ Power of power: multiply, $(x^3)^3 = x^{3*3} = x^9$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$
- ▶ Simplify bases when same exponent: $x^4 * y^4 = (xy)^4$ and
 $m^{-2}n^2 = (m^{-1}n)^2$
- ▶ Power of power: multiply, $(x^3)^3 = x^{3*3} = x^9$

Try: $xz^2(x^3z^{-2})^3$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$
- ▶ Simplify bases when same exponent: $x^4 * y^4 = (xy)^4$ and
 $m^{-2}n^2 = (m^{-1}n)^2$
- ▶ Power of power: multiply, $(x^3)^3 = x^{3*3} = x^9$

Try: $xz^2(x^3z^{-2})^3 = (xz^2(x^9z^{-6}))$

Exponents

- ▶ Multiplying a number by itself: $x * x * x * x = x^4$
- ▶ Add or subtract exponents when multiplying same variable:
 $x^4 * x^{-3} = x$
- ▶ Simplify bases when same exponent: $x^4 * y^4 = (xy)^4$ and
 $m^{-2}n^2 = (m^{-1}n)^2$
- ▶ Power of power: multiply, $(x^3)^3 = x^{3*3} = x^9$

Try: $xz^2(x^3z^{-2})^3 = (xz^2(x^9z^{-6})) = x^{10}z^{-4}$

Exponents, Exponentials, Exponential functions

Exponents

Exponents are where you take a variable to some power – e.g. x^a where x is a variable and a is a constant. Typically, we focus on the numerical portion of the exponent—calling it ‘the exponent’.

Exponential

An exponential is the reverse of the exponent – here it is a number to the power of a variable, e.g. a^x . To get the x ‘down’, we need to use logarithms (aka logs).

Exponential Function

The exponential function has a particular base, e , (where e is Euler’s e and is approx 2.72.)

Functions: Quadratic Functions and Polynomials

Quadratic functions : highest order (largest exponent) is equal to 2.

Functions: Quadratic Functions and Polynomials

Quadratic functions : highest order (largest exponent) is equal to 2.

$$y = \alpha + \beta_1 x + \beta_2 x^2.$$

Functions: Quadratic Functions and Polynomials

Quadratic functions : highest order (largest exponent) is equal to 2.

$$y = \alpha + \beta_1 x + \beta_2 x^2.$$

Higher order polynomials: more of the same, but now the highest order can be anything.

Functions: Quadratic Functions and Polynomials

Quadratic functions : highest order (largest exponent) is equal to 2.

$$y = \alpha + \beta_1 x + \beta_2 x^2.$$

Higher order polynomials: more of the same, but now the highest order can be anything. Some examples include:

$$y = \alpha + \beta_1 x + \beta_2 x^3. \text{ and } y = \alpha + \beta_1 x + \beta_2 x^2 + \beta_3 x^3.$$

Functions: Quadratic Functions and Polynomials

Quadratic functions : highest order (largest exponent) is equal to 2.

$$y = \alpha + \beta_1x + \beta_2x^2.$$

Higher order polynomials: more of the same, but now the highest order can be anything. Some examples include:

$$y = \alpha + \beta_1x + \beta_2x^3. \text{ and } y = \alpha + \beta_1x + \beta_2x^2 + \beta_3x^3.$$

We care because we might see these as utility functions or use higher order values as a way to represent the relationship between two variables in a linear regression Know what these concepts are—will be relevant later.

Logs and other functions

Logs are the inverses of exponents: the power to which you raise the base, e.g. 10, to produce a given value, e.g. z

Logs and other functions

Logs are the inverses of exponents: the power to which you raise the base, e.g. 10, to produce a given value, e.g. z

You will see logs!

Logs and other functions

Logs are the inverses of exponents: the power to which you raise the base, e.g. 10, to produce a given value, e.g. z

You will see logs!

Logarithms (typically base 10 ($\log(x)$) or base e ($\ln(e)$), but any base is possible, e.g. $\log_{8675309}x$ (Bases aside from e and 10 will be specified).

- ▶ $y = \log(z) \leftrightarrow 10^y = z$
- ▶ $y = \ln(z) \leftrightarrow e^y = z$
- ▶ $\log(1) = 0$

Logs and other functions

Logs are the inverses of exponents: the power to which you raise the base, e.g. 10, to produce a given value, e.g. z

You will see logs!

Logarithms (typically base 10 ($\log(x)$) or base e ($\ln(e)$), but any base is possible, e.g. $\log_{8675309} x$ (Bases aside from e and 10 will be specified).

- ▶ $y = \log(z) \leftrightarrow 10^y = z$
- ▶ $y = \ln(z) \leftrightarrow e^y = z$
- ▶ $\log(1) = 0$

Exponents in log are different from what you might expect:

- ▶ $\log(x^2) = 2(\log(x))$
- ▶ $\log(x/y) = \log(x) - \log(y)$ provided $(x, y > 0)$

Logs help weigh smaller values more heavily; adding units not linear—less meaningful for larger values

($\log(100) = 2$ $\log(1000) = 3$)

Logs Practice

Simplify the following

▶ $\log(x^4)$

▶ $\log(xy)$

▶ $\ln(e^3)$

▶ $\ln(1)$

▶ $\log(3) + \log(7)$

Logs Practice

Simplify the following

▶ $\log(x^4) = 4\log(x)$

▶ $\log(xy)$

▶ $\ln(e^3)$

▶ $\ln(1)$

▶ $\log(3) + \log(7)$

Logs Practice

Simplify the following

- ▶ $\log(x^4) = 4\log(x)$
- ▶ $\log(xy) = \log(x) + \log(y)$
- ▶ $\ln(e^3)$
- ▶ $\ln(1)$
- ▶ $\log(3) + \log(7)$

Logs Practice

Simplify the following

- ▶ $\log(x^4) = 4\log(x)$
- ▶ $\log(xy) = \log(x) + \log(y)$
- ▶ $\ln(e^3) = 3$
- ▶ $\ln(1)$
- ▶ $\log(3) + \log(7)$

Logs Practice

Simplify the following

- ▶ $\log(x^4) = 4\log(x)$
- ▶ $\log(xy) = \log(x) + \log(y)$
- ▶ $\ln(e^3) = 3$
- ▶ $\ln(1) = 0$
- ▶ $\log(3) + \log(7)$

Simplify the following

- ▶ $\log(x^4) = 4\log(x)$
- ▶ $\log(xy) = \log(x) + \log(y)$
- ▶ $\ln(e^3) = 3$
- ▶ $\ln(1) = 0$
- ▶ $\log(3) + \log(7) = \log(21)$

More Practice

▶ $2e^{6x} = 18$

▶ $e^{x^2} = 1$

▶ $2^x = e^5$

▶ $2^{x-2} = 5$

▶ $\ln(x^2) = 5$

▶ $\ln(x^3) - \ln(x) = \ln(16)$

When would I ever actually see a log function??

When would I ever actually see a log function??

<https://thesocietypages.org/graphicsociology/2010/12/07/1247/>

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education, partisan id, income, age, closeness. *How can we categorize these variables?*

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education (discrete (count)), partisan id (categorical (0,1)), income (continuous), age (discrete (count)), closeness (interval-ratio, 0-1). *(categorization in parentheses)*

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education (discrete (count)), partisan id (categorical (0,1)), income (continuous), age (discrete (count)), closeness (interval-ratio, 0-1). *(categorization in parentheses)*

The probability is given by this function: how can we interpret the following? $p_v = f(ed, p, i, a, c)$

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education (discrete (count)), partisan id (categorical (0,1)), income (continuous), age (discrete (count)), closeness (interval-ratio, 0-1). *(categorization in parentheses)*

The probability is given by this function: how can we interpret the following? $p_v = f(ed, p, i, a, c)$ *The probability is a function of the variables given above.*

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education (discrete (count)), partisan id (categorical (0,1)), income (continuous), age (discrete (count)), closeness (interval-ratio, 0-1). (*categorization in parentheses*)

The probability is given by this function: how can we interpret the following? $p_v = f(ed, p, i, a, c)$ *The probability is a function of the variables given above.* Can further specify as a direct linear function: $p_v = \alpha + \beta_1 ed + \beta_2 p + \beta_3 i + \beta_4 a + \beta_5 c$

Tying it all together

Consider the example in Moore and Siegel (pg 72): predicting voting probability of an individual in a national education. Variables: education (discrete (count)), partisan id (categorical (0,1)), income (continuous), age (discrete (count)), closeness (interval-ratio, 0-1). (*categorization in parentheses*)

The probability is given by this function: how can we interpret the following? $p_v = f(ed, p, i, a, c)$ *The probability is a function of the variables given above.* Can further specify as a direct linear function: $p_v = \alpha + \beta_1 ed + \beta_2 p + \beta_3 i + \beta_4 a + \beta_5 c$ But this is one of many options (how would we apply the quadratic or linear functions?)