

Day 5: Calculus I, Derivatives

Sarah Moore and Jason Seawright

Math Camp 2022

Day 5 Agenda

- ▶ Derivatives: Concept, notation, and how-to
- ▶ Fundamental Derivative Rules
- ▶ Partial Derivatives
- ▶ Advanced Rules of Derivatives

- ▶ This is often where we get a lot of nervous faces in the room. But, I can assure you that we are going to stick together in this and everyone will come out knowing some minimal calculus stuff to get you through these quant methods courses.

What is calculus?

- ▶ So basically, calculus is the study of *instantaneous change of a function*.

Well, what does that mean?

What is calculus?

- ▶ So basically, calculus is the study of *instantaneous change of a function*.

Well, what does that mean?

- ▶ On the one hand, we could be interested in *discrete* change. Discrete change tells us the difference between two points on a graph, thus the difference between two observations modeled by a function.

What is calculus?

- ▶ So basically, calculus is the study of *instantaneous change of a function*.

Well, what does that mean?

- ▶ On the one hand, we could be interested in *discrete* change. Discrete change tells us the difference between two points on a graph, thus the difference between two observations modeled by a function.
- ▶ Finding discrete change means finding the slope of two points, or the *secant*. You all are probably familiar with $\frac{\text{rise}}{\text{run}}$ from algebra.

What is calculus?

- ▶ But, finding the secant has a limitation. Discrete change only tells about the functional behavior *over an interval*. Instead we might want to find the rate of change at a *very specific moment* in the function.

What is calculus?

- ▶ But, finding the secant has a limitation. Discrete change only tells about the functional behavior *over an interval*. Instead we might want to find the rate of change at a *very specific moment* in the function.
- ▶ Consequently, we might also want to find extrema (min and max) of a function, any other critical points, or understand its shape. As you may remember from other courses, these are difficult tasks with merely a function $f(x)$.

What is calculus?

- ▶ But, finding the secant has a limitation. Discrete change only tells about the functional behavior *over an interval*. Instead we might want to find the rate of change at a *very specific moment* in the function.
- ▶ Consequently, we might also want to find extrema (min and max) of a function, any other critical points, or understand its shape. As you may remember from other courses, these are difficult tasks with merely a function $f(x)$.
- ▶ Calculus gives us some tools to calculate instantaneous change and other downstream quantities.

Limits, Secants, and Capturing Change


- ▶ In our last session we went over limits of functions. As a reminder, the limit of a function $f(x)$ at a given point x is the value of the function as it approaches the given x .

Limits, Secants, and Capturing Change

- ▶ In our last session we went over limits of functions. As a reminder, the limit of a function $f(x)$ at a given point x is the value of the function as it approaches the given x .
- ▶ Therefore, to capture discrete change we calculate the secant between two points by taking the difference of the functional limits, i.e. $\frac{f(x_2)-f(x_1)}{x_2-x_1}$.


Limits, Secants, and Capturing Change

- To capture the instantaneous or continuous rate of change of a function, we *could* take the difference of the limits over iterations, making the interval between each x smaller and smaller.



Limits, Secants, and Capturing Change

- ▶ To capture the instantaneous or continuous rate of change of a function, we *could* take the difference of the limits over iterations, making the interval between each x smaller and smaller.



- ▶ However, the concept of **tangents**, and consequentially **derivatives**, makes this easier.

Tangents and Derivatives

- ▶ While a secant is a slope of a given line, a *tangent* is a line that touches the function at a given point. The tangent's slope tell us about the slope of the primary function, or the instantaneous rate of change, *at that particular point*.

Tangents and Derivatives

- ▶ While a secant is a slope of a given line, a *tangent* is a line that touches the function at a given point. The tangent's slope tell us about the slope of the primary function, or the instantaneous rate of change, *at that particular point*.
- ▶ Therefore, the problem becomes *how to find the tangent's slope*.

Notation

- ▶ There are a couple of ways to notate derivatives, all meaning the same thing.

$$f'(x)$$

$$\frac{dy}{dx}$$

Notation

- ▶ There are a couple of ways to notate derivatives, all meaning the same thing.

$f'(x)$ Read: "f prime of x"

$\frac{dy}{dx}$ Read: "dy-dx" or "dy over dx"

Notation

- ▶ There are a couple of ways to notate derivatives, all meaning the same thing.

$f'(x)$ Read: "f prime of x"

$\frac{dy}{dx}$ Read: "dy-dx" or "dy over dx"

- ▶ There also possibilities for **higher order derivatives**, i.e. derivatives of derivatives and so on. . .

Notation

- ▶ There are a couple of ways to notate derivatives, all meaning the same thing.

$f'(x)$ Read: "f prime of x"

$\frac{dy}{dx}$ Read: "dy-dx" or "dy over dx"

- ▶ There also possibilities for **higher order derivatives**, i.e. derivatives of derivatives and so on. . .
- ▶ Higher order derivatives will use the same types of notation, with details to indicate the order of the derivative

Notation

- ▶ There are a couple of ways to notate derivatives, all meaning the same thing.

$f'(x)$ Read: "f prime of x"

$\frac{dy}{dx}$ Read: "dy-dx" or "dy over dx"

- ▶ There also possibilities for **higher order derivatives**, i.e. derivatives of derivatives and so on. . .
- ▶ Higher order derivatives will use the same types of notation, with details to indicate the order of the derivative
- ▶ $f''(x)$ or $\frac{d^2y}{dx^2}$ for a second derivative; $f'''(x)$ or $\frac{d^3y}{dx^3}$ for a third derivative

Calculating a Derivative

To calculate the derivative, begin with the secant formula. Use this formula to reduce the difference to some arbitrarily small value, h .

As h goes to zero, we go from discrete to instantaneous change.

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

$$\lim_{h \rightarrow 0} \frac{3(x + h) - 3x}{h}$$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

$$\lim_{h \rightarrow 0} \frac{3(x + h) - 3x}{h} = \lim_{h \rightarrow 0} \frac{3x + 3h - 3x}{h}$$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

$$\lim_{h \rightarrow 0} \frac{3(x + h) - 3x}{h} = \lim_{h \rightarrow 0} \frac{3x + 3h - 3x}{h} = \lim_{h \rightarrow 0} \frac{3h}{h}$$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

$$\lim_{h \rightarrow 0} \frac{3(x + h) - 3x}{h} = \lim_{h \rightarrow 0} \frac{3x + 3h - 3x}{h} = \lim_{h \rightarrow 0} \frac{3h}{h} = \lim_{h \rightarrow 0} 3$$

Example: Calculating the Derivative of $f(x) = 3x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $3x$

$$\lim_{h \rightarrow 0} \frac{3(x + h) - 3x}{h} = \lim_{h \rightarrow 0} \frac{3x + 3h - 3x}{h} = \lim_{h \rightarrow 0} \frac{3h}{h} = \lim_{h \rightarrow 0} 3 = 3$$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

$$\lim_{h \rightarrow 0} \frac{2(x + h) - 2x}{h}$$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

$$\lim_{h \rightarrow 0} \frac{2(x + h) - 2x}{h} = \lim_{h \rightarrow 0} \frac{2x + 2h - 2x}{h}$$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

$$\lim_{h \rightarrow 0} \frac{2(x + h) - 2x}{h} = \lim_{h \rightarrow 0} \frac{2x + 2h - 2x}{h} = \lim_{h \rightarrow 0} \frac{2h}{h}$$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

$$\lim_{h \rightarrow 0} \frac{2(x + h) - 2x}{h} = \lim_{h \rightarrow 0} \frac{2x + 2h - 2x}{h} = \lim_{h \rightarrow 0} \frac{2h}{h} = \lim_{h \rightarrow 0} 2$$

Example: Calculating the Derivative of $f(x) = 2x$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: $2x$

$$\lim_{h \rightarrow 0} \frac{2(x + h) - 2x}{h} = \lim_{h \rightarrow 0} \frac{2x + 2h - 2x}{h} = \lim_{h \rightarrow 0} \frac{2h}{h} = \lim_{h \rightarrow 0} 2 = 2$$

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

$$\lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h}$$

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

$$\lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + 2h^2 - x^2}{h}$$

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

$$\lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + 2h^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{2xh + 2h^2}{h}$$

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h} &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + 2h^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{2xh + 2h^2}{h} \\ &= \lim_{h \rightarrow 0} 2x + 2h \end{aligned}$$

Example: Calculating the Derivative of $f(x) = x^2$

Secant Formula:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Example: x^2

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h} &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + 2h^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{2xh + 2h^2}{h} \\ &= \lim_{h \rightarrow 0} 2x + 2h = 2x + 0 = 2x \end{aligned}$$

So... what does it all mean?

- Derivatives give us information regarding the rate of change of given function at a value x , we can use this information to learn more about the base function at that particular point.

So... what does it all mean?

- ▶ Derivatives give us information regarding the rate of change of given function at a value x , we can use this information to learn more about the base function at that particular point.
- ▶ Think back to what the slope of a linear equation tells us from the formula $y = mx + b$.

So... what does it all mean?

- ▶ Derivatives give us information regarding the rate of change of given function at a value x , we can use this information to learn more about the base function at that particular point.
- ▶ Think back to what the slope of a linear equation tells us from the formula $y = mx + b$.
- ▶ If m is positive, the slope is increasing; if m is negative, the slope is decreasing.

So... what does it all mean?

- ▶ Derivatives give us information regarding the rate of change of given function at a value x , we can use this information to learn more about the base function at that particular point.
- ▶ Think back to what the slope of a linear equation tells us from the formula $y = mx + b$.
- ▶ If m is positive, the slope is increasing; if m is negative, the slope is decreasing.
- ▶ The derivative gives us information that we interpret similarly.

Derivative as information: Rate of change

- ▶ Positive Derivative

Derivative as information: Rate of change

- ▶ Positive Derivative Function is increasing
- ▶ Negative Derivative

Derivative as information: Rate of change


- ▶ Positive Derivative Function is increasing
- ▶ Negative Derivative Function is decreasing
- ▶ Zeroes?

Derivative as information: Rate of change


- ▶ Positive Derivative Function is increasing
- ▶ Negative Derivative Function is decreasing
- ▶ Zeroes? max or min

Derivatives and Extrema

Extrema: max or min of a function, i.e. where is the topmost or bottommost of the function?



Derivatives and Extrema



Worst Behavior


As we saw before: the function must be *continuous* on the *interval* to be differentiable.

Some functions are not differentiable *at all* or are not differentiable at *a certain point*. Need to determine the continuity of the function.


Continuity

Continuous function: draw without picking up a pencil

YES



NO



Derivative: Calculation

While we can calculate the derivative using the formula from before, it's a bit tedious. Can't there be another way?

Derivative: Calculation

While we can calculate the derivative using the formula from before, it's a bit tedious. Can't there be another way?

Think through the work it would take to differentiate $4x^3 + 3x - 2$ using $\frac{f(x_0+h)-f(x_0)}{h} \dots$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

2. $f(x) = k$ has derivative $f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

► $f(x) = 3x,$

2. $f(x) = k$ has derivative $f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

► $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4,$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3,$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

▶ $f(x) = 3x, g(x) = 7,$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

▶ $f(x) = 3x, g(x) = 7, 3 + 0 = 3$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

▶ $f(x) = 3x, g(x) = 7, 3 + 0 = 3$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

▶ $[3x - 7]',$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

▶ $f(x) = 3x, g(x) = 7, 3 + 0 = 3$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

▶ $[3x - 7]', 3 - 0 = 3$

Derivative Rules

Take the derivative $f(x)$ as $f'(x)$ below with constant k :

1. $f(k * x) = k * f(x), f'(k * x) = k * f'(x)$

▶ $f(x) = 3x, f'(x) = 3$

2. $f(x) = k$ has derivative $f'(x) = 0$

▶ $f(x) = 4, f'(x) = 0$

3. $f(x) = x^n, f'(x) = n * x^{n-1}$

▶ $f(x) = x^3, f'(x) = 3x^2$

4. $[f(x) + g(x)]' = f'(x) + g'(x)$

▶ $f(x) = 3x, g(x) = 7, 3 + 0 = 3$

5. $[f(x) - g(x)]' = f'(x) - g'(x)$

▶ $[3x - 7]', 3 - 0 = 3$

NOTE: $[f(x) * g(x)]' = f'(x) * g'(x)$ Ex: $(3x * 10x)' = 30$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.

Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$$

$$f'(x) = \frac{-1}{x(x+0)}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$$

$$f'(x) = \frac{-1}{x(x+0)}$$

$$f'(x) = \frac{-1}{x^2}$$

Derivatives Two Ways

We can check these handy formulas work as they should. Let's try.
Find the derivative of $f(x) = \frac{1}{x}$

► Formal Definition: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x}{x(x+h)} - \frac{(x+h)}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$$

$$f'(x) = \frac{-1}{x(x+0)}$$

$$f'(x) = \frac{-1}{x^2}$$

$$f'(x) = -x^{-2}$$

Using the Simpler Rules

Relevant Rules: $f(x) = x^n, f'(x) = n * x^{n-1}$

Using the Simpler Rules

Relevant Rules: $f(x) = x^n, f'(x) = n * x^{n-1}$

$$f(x) = x^{-1}$$

Using the Simpler Rules

Relevant Rules: $f(x) = x^n, f'(x) = n * x^{n-1}$

$$f(x) = x^{-1}$$

$$f'(x) = -1 * x^{-2}$$

Using the Simpler Rules

Relevant Rules: $f(x) = x^n, f'(x) = n * x^{n-1}$

$$f(x) = x^{-1}$$

$$f'(x) = -1 * x^{-2}$$

$$f'(x) = -x^{-2}$$

Practice Problems

Find where functions are continuous and find derivatives

► $f(x) = 5$

► $f(x) = 3x - 7$

► $f(x) = 3x^2$

► $f(x) = \frac{x^2}{x}$

► $f(s) = s^{-2}$

► $f(y) = y(y + 7)(y - 3)$

► $f(z) = \frac{z^2 - 5z - 6}{z + 1}$

Practice Problems

Find where functions are continuous and find derivatives

- ▶ $f(x) = 5$, $f'(x) = 0$
- ▶ $f(x) = 3x - 7$, $f'(x) = 3$
- ▶ $f(x) = 3x^2$, $f'(x) = 6x$
- ▶ $f(x) = \frac{x^2}{x}$, $f'(x) = 1$
- ▶ $f(s) = s^{-2}$, $f'(s) = -2s^{-3}$ (not continuous at $s = 0$)
- ▶ $f(y) = y(y + 7)(y - 3)$, $f'(y) = 3y^2 + 8y - 21$
- ▶ $f(z) = \frac{z^2 - 5z - 6}{z + 1}$, $f'(z) = 1$ (not continuous at $z = -1$)

Higher Order Derivatives

Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

► $f(x) = x^5 + 3x^3 + 2x + 8$

Higher Order Derivatives

Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

- ▶ $f(x) = x^5 + 3x^3 + 2x + 8$

- ▶ $f'(x) = 5x^4 + 9x^2 + 2$

Higher Order Derivatives

Second derivatives (n^{th} derivatives): take a derivative a second (n^{th}) time

Rate of change of rate of change (velocity vs acceleration)

- ▶ $f(x) = x^5 + 3x^3 + 2x + 8$

- ▶ $f'(x) = 5x^4 + 9x^2 + 2$

- ▶ $f''(x) = 20x^3 + 18x$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z$$

$$f(x, y) = x + 4y$$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z$$

$$f(x, y) = x + 4y$$

Partial Derivatives

Similar to 'regular' derivative; treat additional variable(s) as constants. Written as ∂_x or $\frac{\partial f}{\partial x}(x, \dots)$

Ex: $y = 3xz$, $\partial_x = 3z$

Find ∂_x

$$f(x, z) = 7xz + 4x^2 + z \quad \partial_x = 7z + 8x$$

$$f(x, y) = x + 4y \quad \partial_x = 1$$

(Additional) Rules for derivatives

So far, we've just had simple functions but what if we are multiplying, dividing, or have an otherwise-more-advanced function?

- ▶ Product Rule: $f(x) * g(x)$
- ▶ Quotient Rule: $\frac{f(x)}{g(x)}$
- ▶ Chain Rule: $f(g(x))$
- ▶ Other: eg, exponentials: e^x , $\ln(x)$

(Additional) Rules for derivatives

So far, we've just had simple functions but what if we are multiplying, dividing, or have an otherwise-more-advanced function?

- ▶ Product Rule: $f(x) * g(x)$, $x^3 * x^2$
- ▶ Quotient Rule: $\frac{f(x)}{g(x)}$, $\frac{x^4+3x}{x^2}$
- ▶ Chain Rule: $f(g(x))$, $(x^2 + 1)^3$ (composition!)
- ▶ Other: eg, exponentials: e^x , $\ln(x)$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

► **Example:** $(3x+4)(x+2)$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

► **Example:** $(3x+4)(x+2)$

► To take the derivative using the product rule, we do the following: $f'(x) * g(x) + g'(x)f(x)$.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

- ▶ **Example: $(3x+4)(x+2)$**
- ▶ To take the derivative using the product rule, we do the following: $f'(x) * g(x) + g'(x)f(x)$.
- ▶ For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

► **Example: $(3x+4)(x+2)$**

► To take the derivative using the product rule, we do the following: $f'(x) * g(x) + g'(x)f(x)$.

► For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$.

► Then, we substitute these in:

$$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$$

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

► **Example:** $(3x+4)(x+2)$

► To take the derivative using the product rule, we do the following: $f'(x) * g(x) + g'(x)f(x)$.

► For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$.

► Then, we substitute these in:

$$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$$

► Simplify to get: $3x + 6 + 3x + 4 = 6x + 10$.

Product Rule

If we have two things multiplied together and need the derivative, we have two options: multiply everything and then take the derivative OR use the product rule.

- ▶ **Example: $(3x+4)(x+2)$**
- ▶ To take the derivative using the product rule, we do the following: $f'(x) * g(x) + g'(x)f(x)$.
- ▶ For us, our two functions are $f(x) = 3x + 4$ and $g(x) = x + 2$. The derivatives are $f'(x) = 3$ and $g'(x) = 1$.
- ▶ Then, we substitute these in:
$$f'(x) * g(x) + g'(x)f(x) = 3(x + 2) + 1(3x + 4)$$
- ▶ Simplify to get: $3x + 6 + 3x + 4 = 6x + 10$.
- ▶ To take the derivative using our previous approach, we first multiply: $3x^2 + 4x + 6x + 8 = 3x^2 + 10x + 8$. Then, just take the derivative of each term: $f'(x) = 6x + 10$.

Product Rule: A Motivating Example

Suppose that instead you had $(3x^2 + 3x + 4)(x^3 + 2x^2 + x + 2)$.
Now the product rule is looking a little nicer!

$$f(x) = (3x^2 + 3x + 4) \text{ and } g(x) = (x^3 + 2x^2 + x + 2)$$

$$f'(x) = 6x + 3 \text{ and } g'(x) = 3x^2 + 4x + 1.$$

We can substitute this into the formula:

$$f'(x) * g(x) + g'(x)f(x)$$

$$\rightarrow (6x + 3)(x^3 + 2x^2 + x + 2) + (3x^2 + 4x + 1)(3x^2 + 3x + 4)$$

.

Quotient Rule

Example: $\frac{3x^2}{x+2}$.

Formula is $\frac{f'(x) \cdot g(x) - g'(x) f(x)}{(g(x))^2}$

So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$,
therefore $f'(x) = 6x$ and $g'(x) = 1$.

Plug in to get:

$$\frac{6x(x+2) - 1(3x^2)}{(x+2)^2} = \frac{6x^2 + 12x - 3x^2}{(x+2)^2} = \frac{3x^2 + 12x}{(x+2)^2}$$

Practice: $\frac{x-4}{x+5} \cdot \frac{3x^3}{x+2}$

Quotient Rule

Example: $\frac{3x^2}{x+2}$.

Formula is $\frac{f'(x) \cdot g(x) - g'(x) f(x)}{(g(x))^2}$

So, we identify the following: $f(x) = 3x^2$ and $g(x) = x + 2$,
therefore $f'(x) = 6x$ and $g'(x) = 1$.

Plug in to get:

$$\frac{6x(x+2) - 1(3x^2)}{(x+2)^2} = \frac{6x^2 + 12x - 3x^2}{(x+2)^2} = \frac{3x^2 + 12x}{(x+2)^2}$$

Practice: $\frac{x-4}{x+5} \cdot \frac{3x^3}{x+2}$

Ans: $\frac{9}{(x+5)^2} \cdot \frac{6x^3+18x^2}{(x+2)^2}$

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$.
We can use the chain rule to evaluate this.

What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$.

So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

- The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$.

Chain Rule

Sometimes, you have a function to a power: $f(g(x)) = (x + 3)^3$.
We can use the chain rule to evaluate this.

What we do is we take the derivative of the function and multiply it by the derivative of the inside: $f'(g(x)) * g'(x)$.

So, for our example: $f(x) = x^3$ and $g(x) = (x + 3)$.

- ▶ The derivative of each is $f'(x) = 3x^2$ and $g'(x) = 1$.
- ▶ We substitute in to get: $3(x + 3)^2 * 1$.

More Chain Rule

► **Try:** $f(x) = (2x^2 + 8x)^4$ $f(x) = (9x - x^2)^6$

More Chain Rule

► **Try:** $f(x) = (2x^2 + 8x)^4$ $f(x) = (9x - x^2)^6$

► **Ans** $4(4x + 8)(2x^2 + 8x)^3$
 $6(9 - 2x)(9x - x^2)^5 = (54 - 12x)(9x - 2x^2)^5$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \text{ (a favorite of mine)}$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \text{ (used rarely, if ever)}$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \text{ (rarely used)}$$

You can make these more complicated by including a function of x .
How would we take the derivative in that case?

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \text{ (a favorite of mine)}$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x(\ln(a)) \text{ (used rarely, if ever)}$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \text{ (rarely used)}$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

EX: $\ln(3x)$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \text{ (a favorite of mine)}$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x(\ln(a)) \text{ (used rarely, if ever)}$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \text{ (rarely used)}$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

$$\textbf{EX: } \ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \text{ (a favorite of mine)}$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \text{ (used rarely, if ever)}$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \text{ (rarely used)}$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

EX: $\ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$

TRY: $\ln(x^2), e^{2x}$

Exponentials: e and \ln

You can take the derivative of continuous functions – including those with a log and/or e in them. The rules are a little hard, but once you learn them, it's not too bad:

$$f(x) = e^x \quad f'(x) = e^x \text{ (a favorite of mine)}$$

$$f(x) = e^{g(x)} \quad f'(x) = e^{g(x)} * g'(x)$$

$$f(x) = a^x \quad f'(x) = a^x (\ln(a)) \text{ (used rarely, if ever)}$$

$$f(x) = \ln(x) \quad f'(x) = \frac{1}{x}$$

$$f(x) = \ln(g(x)) \quad f'(x) = \frac{1}{x} * g'(x)$$

$$f(x) = \log_a(x) \quad f'(x) = \frac{1}{x \ln(a)} \text{ (rarely used)}$$

You can make these more complicated by including a function of x . How would we take the derivative in that case? Chain rule!

EX: $\ln(3x) \quad f'(x) = \frac{1}{3x} * 3 = \frac{1}{x}.$

TRY: $\ln(x^2), e^{2x}$ **ANS:** $\frac{2}{x}$ and $2e^{2x}$

Derivatives in Review

There are a few more handy rules and techniques that are important, perhaps even on your homework:

| Common Derivatives | | |
|--|---|--|
| $\frac{d}{dx}(x) = 1$ | $\frac{d}{dx}(\csc x) = -\csc x \cot x$ | $\frac{d}{dx}(a^x) = a^x \ln(a)$ |
| $\frac{d}{dx}(\sin x) = \cos x$ | $\frac{d}{dx}(\cot x) = -\csc^2 x$ | $\frac{d}{dx}(e^x) = e^x$ |
| $\frac{d}{dx}(\cos x) = -\sin x$ | $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$ | $\frac{d}{dx}(\ln(x)) = \frac{1}{x}, x > 0$ |
| $\frac{d}{dx}(\tan x) = \sec^2 x$ | $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$ | $\frac{d}{dx}(\ln x) = \frac{1}{x}, x \neq 0$ |
| $\frac{d}{dx}(\sec x) = \sec x \tan x$ | $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$ | $\frac{d}{dx}(\log_a(x)) = \frac{1}{x \ln a}, x > 0$ |

Figure 1: [Source on Stack Exchange](#)

And if you really want to explore more, check out all [these techniques](#).