# Day 1: Notation and Logic

Jean Clipperton

Math Camp 2021

#### Miscellaneous Info

- Going to jump around a little today
- ▶ PS 490 R workshop to help smooth the way with this course
- Will post today's slides by end of day

# Now, to the good stuff!

#### Overview of Math Camp

- ► TeX or Rmd are great for homework–Michelle will walk you through soon
- http://detexify.kirelabs.org/classify.html
- Canvas for symbol cheat sheet
- ▶ OH: we'll build in time at the end of class sessions in addition to during sessions. Please take advantage of this!
- ► Email: jean.clipperton@northwestern.edu, MichelleBuenoVasquez2024@u.northwestern.edu
- Will LOOSELY follow recommended book may jump around a bit
- Homeworks not graded but you will submit them

# Math (P)refresher: Review Topics

- Broad Overview
- ▶ Sets
- ► Algebra review
- Inequalities
- Combinatorics
- ► Summation
- Derivatives
- ► Integrals

#### **Broad Overview**

From the Top: Variables and Constants

Theories

Theories are how we frame statements (hypotheses, propositions) about the world, using concepts.

Concepts

Concepts are the ideas, aspects to be measured

Example

Actors and Laws

Theory: More *political actors* involved in a process hampers *legislative productivity* because it's harder to reach an agreement

Need to define how you count an actor and measure productivity

From the Top: Variables and Constants

#### **Broad Overview**

From the Top: Variables and Constants

Theories

Theories are how we frame statements (hypotheses, propositions) about the world, using concepts.

Concepts

Concepts are the ideas, aspects to be measured

Example

Actors and Laws

Theory: More *political actors* involved in a process hampers *legislative productivity* because it's harder to reach an agreement

Need to define how you count an actor and measure productivity

From the Top: Variables and Constants

Sets are collections of elements. Can relate variables to sets.

Sets are collections of elements. Can relate variables to sets.

#### Example

Suppose we are measuring voting —what is the variable and what are our possible values?

Variable: number of times voted on a series of bills.

Variable: actors in support of a bill.

Sets are collections of elements. Can relate variables to sets.

#### Example

Suppose we are measuring voting —what is the variable and what are our possible values?

Variable: number of times voted on a series of bills.

**Set** {3, 4, 0 }.

Variable: actors in support of a bill.

Sets are collections of elements. Can relate variables to sets.

#### Example

Suppose we are measuring voting —what is the variable and what are our possible values?

Variable: number of times voted on a series of bills.

**Set** {3, 4, 0 }.

Variable: actors in support of a bill.

**Set** { A, O, P }.

We can think about these sets' possible elements as follows

▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$ 

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line
- ▶ I: Imaginary numbers  $(i = \sqrt{-1})$

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line
- ▶ I: Imaginary numbers  $(i = \sqrt{-1})$
- C: Complex numbers (a + bi)

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line
- ▶ I: Imaginary numbers  $(i = \sqrt{-1})$
- C: Complex numbers (a + bi)

We can think about these sets' possible elements as follows

- ▶ N: Natural numbers  $\{(0), 1, 2, 3, ...\}$
- ► Z: Integers (negative and positive including zero)  $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- ► R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line
- ▶ I: Imaginary numbers  $(i = \sqrt{-1})$
- C: Complex numbers (a + bi)

#### Notes:

Subscript:  $Z_+$  only the positive or  $Q_-$  negative elements of the set Superscript:  $N^2$  dimensions of the space

#### Measurement

#### Measurement

- ► Nominal (categorical): no mathematical relationship between the variables
- Ordinal: categorical variable with set relationship (can compare items to one another)
- ▶ Interval: distance between numerical values has meaning (e.g. 0.1, 2-2 is two greater than 0)
- Ratio (interval-ratio): distance between numerical values has meaning AND zero is meaningful

#### Measurement: Examples

#### Measurement: Examples

- ► Nominal (categorical): no mathematical relationship between the variables **Eye color**
- Ordinal: categorical variable with set relationship (can compare items to one another) Age: above/under 18
- Interval: distance between numerical values has meaning (e.g. 0,1, 2 − 2 is two greater than 0) Approval from 1 to 5
- Ratio (interval-ratio): distance between numerical values has meaning AND zero is meaningful Number of years of grad school

# Variables and Sets: Putting Things Together

- Solution set: set of all solutions to an equation
- ► Sample space: set that contains all the values a variable can take
- Subsets: groupings that fall within other sets

# Variables and Sets: Putting Things Together

Set operations

Can combine sets by looking at the *difference*, *complement*, *intersection*, *union*, and *partition* of sets.

# Variables and Sets: Putting Things Together

#### Set operations

Can combine sets by looking at the *difference*, *complement*, *intersection*, *union*, and *partition* of sets.

#### Example

Consider the set of all men and all women. The intersection is what is in both sets ('and') while the union ('or') is what is the collection of the two. We'll go into this vocabulary more tomorrow.

#### Notation Refresher

- ▶ There exists, ∃ for all, ∀
- ▶ Union ∪, Intersection ∩
- ► Excluding \(\psi\), ¬
- ▶ Empty set ∅
- ▶ Element ∈
- ▶ Equivalent ≡
- ▶ Such that (s.t) or |, e.g.  $\{x|x > 7\}$
- Subset ⊂, ⊆ (these function roughly like the less than/less than equal to, but for sets)

#### It's all Greek to me!

You'll want to have a basic familiarity with Greek letters as they'll come up from time to time.

#### It's all Greek to me!

You'll want to have a basic familiarity with Greek letters as they'll come up from time to time.

Some commonly used letters include  $\delta$  (integrals),  $\Delta$  (difference/change),  $\beta$  (coefficients),  $\mu$  (mean),  $\sigma$  (standard deviation),  $\lambda$  (eigenvalues (linear algebra)),  $\epsilon$  (error)

Check in – all good?

#### Logic

What do I need to know?

We will use terms regarding proofs and you'll want to have a familiarity with these terms, but we won't be doing logic worksheets and/or testing you on these terms.