
Day 3 AM:

Jean Clipperton

Math Camp 2021



Sequences and Series

Sequences

A sequence is an ordered list of numbers. They can be infinite or finite, but all are *countable*.

Lingo

We refer to the elements by their position in the sequence – the third element would be x_3 . We can talk about the entire sequence as being generated by some equation or formula and represent it accordingly. So, if we take each element to the third, our sequence would be $\{1, 2^3, 3^3, ...\}$ and we could reference the sequence as $\{i^3\}_{i=1}^{\infty}$

Sequences and Series

Series

A series is the *sum* of a sequence. (Book reasoning unhelpful here – we care because you'll be adding probabilities in class).

Summation

We may have a large or otherwise complicated series of numbers to add. For example, suppose we wanted to add the numbers from 1-10. We could write the list out, $(1,2,3,\ldots,10)$ or we could use the summation operator:

$$\sum_{n=1}^{10} n$$

Sets

SETS! & Matrices!

Sets: Numbers

Recall from yesterday:

- ▶ N: Natural numbers {(0), 1, 2, 3, ...}
- ► Z: Integers (negative and positive including zero) $\{..., -3, -2, -1, 0, 1, 2, 3, ...\}$
- Q: Rational numbers (q for quotient, rational numbers, e.g. expressed as a fraction)
- R: Real numbers (positive, negative, zero, integers, fractions/rational); any point on the number line
- ▶ I: Imaginary numbers $(i = \sqrt{-1})$
- C: Complex numbers (a + bi)

Notes: Subscript: Z_+ only the positive or Q_- negative elements of the set Superscript: N^2 dimensions of the space

▶ Roster notation (most common): $\{x_1, x_2, ...\}$

- ▶ Roster notation (most common): $\{x_1, x_2, ...\}$
- ▶ Set builder notation (recipe): $\{x|x \in N, x < 4\}$

- ▶ Roster notation (most common): $\{x_1, x_2, ...\}$
- ▶ Set builder notation (recipe): $\{x | x \in N, x < 4\}$

Try:
$$\{x \in Z, x = x^2\}$$

- ▶ Roster notation (most common): $\{x_1, x_2, ...\}$
- ▶ Set builder notation (recipe): $\{x | x \in N, x < 4\}$

Try: $\{x \in Z, x = x^2\}$ "x in the set of (positive and negative) integers such that x is equal to its own square"

- ▶ Roster notation (most common): $\{x_1, x_2, ...\}$
- ▶ Set builder notation (recipe): $\{x | x \in N, x < 4\}$

Try: $\{x \in Z, x = x^2\}$ "x in the set of (positive and negative)

integers such that x is equal to its own square" The only number that satisfies this is $\{1\}$

There are types of sets:

► Finite (finite number of elements) or infinite (no limit)

There are types of sets:

- ► Finite (finite number of elements) or infinite (no limit)
- ▶ Countable or uncountable (elements can be counted or not (e.g $\{0,1\}$ is countable)

We care about sets and how elements are contained within them, and how the sets are shaped. We'll use this information in probability. It's also helpful when thinking about the possible responses and individuals who may fall in your dataset.

Open

Open sets—essentially the boundary is a little fuzzy. This is the set version of open brackets (). The more technical definition has to do with an 'epsilon ball' where, you can always nudge a little closer to the boundary of the set without actually reaching it.

We care about sets and how elements are contained within them, and how the sets are shaped. We'll use this information in probability. It's also helpful when thinking about the possible responses and individuals who may fall in your dataset.

Open

Open sets—essentially the boundary is a little fuzzy. This is the set version of open brackets (). The more technical definition has to do with an 'epsilon ball' where, you can always nudge a little closer to the boundary of the set without actually reaching it.

Closed

Closed sets have a clear and firm boundary. This is the set version of closed brackets []. Here, you can stand right on the edge of the set.

We care about sets and how elements are contained within them, and how the sets are shaped. We'll use this information in probability. It's also helpful when thinking about the possible responses and individuals who may fall in your dataset.

Open

Open sets—essentially the boundary is a little fuzzy. This is the set version of open brackets (). The more technical definition has to do with an 'epsilon ball' where, you can always nudge a little closer to the boundary of the set without actually reaching it.

Closed

Closed sets have a clear and firm boundary. This is the set version of closed brackets []. Here, you can stand right on the edge of the set.

Open and closed sets matter because this affects how we think about the contents of sets – what we call the 'elements'.

► (Proper) Subset: elements of some sets are contained within a second set, e.g. even integers and integers. (proper subset: \subseteq , subset: \subseteq)

- ▶ (Proper) Subset: elements of some sets are contained within a second set, e.g. even integers and integers. (proper subset: ⊂, subset: ⊆)
- ► Cardinality: number of elements
 - Singleton: one element

- ▶ (Proper) Subset: elements of some sets are contained within a second set, e.g. even integers and integers. (proper subset: ⊂, subset: ⊆)
- ► Cardinality: number of elements
 - ► Singleton: one element
- Empty set: no elements.
 - ▶ $A = \emptyset$ is VERY different from $A = \{\emptyset\}$

- ► (Proper) Subset: elements of some sets are contained within a second set, e.g. even integers and integers. (proper subset: ⊆, subset: ⊆)
- ► Cardinality: number of elements
 - ► Singleton: one element
- Empty set: no elements.
 - $A = \emptyset$ is VERY different from $A = \{\emptyset\}$
- Ordered and unordered sets: ordered set has significance in order (preference ranking) while in unordered, no significance in ranking (e.g. membership roster)

- ► (Proper) Subset: elements of some sets are contained within a second set, e.g. even integers and integers. (proper subset: ⊂, subset: ⊆)
- ► Cardinality: number of elements
 - ► Singleton: one element
- ► Empty set: no elements.
 - ▶ $A = \emptyset$ is VERY different from $A = \{\emptyset\}$
- Ordered and unordered sets: ordered set has significance in order (preference ranking) while in unordered, no significance in ranking (e.g. membership roster)
- ► Universal set (the universe: all elements)

Sets: Union and Intersection

Suppose
$$A = \{4, hat\}$$
 and $B = \{hat, 7\}$

Union (\cup)

Union is the combination of elements in either set (OR):

$$A \cup B = \{4, hat, 7\}$$

Sets: Union and Intersection

Suppose
$$A = \{4, hat\}$$
 and $B = \{hat, 7\}$

Union (∪)

Union is the combination of elements in either set (OR):

$$A \cup B = \{4, hat, 7\}$$

Intersection (\cap)

Intersection is the collection of elements present in both sets (AND): $A \cap B = \{hat\}$ (Def: $A \cap B = X : X \in A \text{ and } X \in B$) Sometimes the intersection is empty.




Figure 1: "Source: XKCD"

Disjoint

Disjoint sets are those where the intersection is empty.

Disjoint

Disjoint sets are those where the intersection is empty.

Partition

Sometimes we may wish to partition a set - to do this, we want to ensure that we *cover* the space (all the elements in the set are assigned to a separate subset), but we also want to make sure that we don't double assign. A proper partition is one where the collection of sets are disjoint and their union is the entire set.

Disjoint

Disjoint sets are those where the intersection is empty.

Partition

Sometimes we may wish to partition a set – to do this, we want to ensure that we *cover* the space (all the elements in the set are assigned to a separate subset), but we also want to make sure that we don't double assign. A proper partition is one where the collection of sets are disjoint and their union is the entire set.

Example: How to partition countries?

Disjoint

Disjoint sets are those where the intersection is empty.

Partition

Sometimes we may wish to partition a set – to do this, we want to ensure that we *cover* the space (all the elements in the set are assigned to a separate subset), but we also want to make sure that we don't double assign. A proper partition is one where the collection of sets are disjoint and their union is the entire set.


Example: How to partition countries? Could do by continents, population size, etc – but pay attention that, say Turkey, isn't assigned to two regions. Or that you don't forget Malta!

Difference Difference between A and B, $A \setminus B$ ("A difference B") is the set containing all the elements of A that are not also in B. $x \in A$ but $x \notin B$


Difference Difference between A and B, $A \setminus B$ ("A difference B") is the set containing all the elements of A that are not also in B. $x \in A$ but $x \notin B$

Complement Complement A' or A^c contains the elements that are not contained in A. $x \in A^c$ if $x \notin A$

Find NU GS\PS/SOC Grad Students



NU GS\PS/SOC Grad Students: AKA, where are the NU grad students who aren't in PS/SOC



Mutually exclusive sets: intersection is empty (e.g. even and odd numbers)

- Mutually exclusive sets: intersection is empty (e.g. even and odd numbers)
- ▶ Partition: collection whose subsets form the universal set (e.g. citizens who are over/under 18). Numerous ways to partition (try)

- Mutually exclusive sets: intersection is empty (e.g. even and odd numbers)
- ▶ Partition: collection whose subsets form the universal set (e.g. citizens who are over/under 18). Numerous ways to partition (try)
- ► Cartesian product: set containing all possible ordered pairs (a,b) where a is from set A and b is from set B for any two sets.

- Mutually exclusive sets: intersection is empty (e.g. even and odd numbers)
- ▶ Partition: collection whose subsets form the universal set (e.g. citizens who are over/under 18). Numerous ways to partition (try)
- ► Cartesian product: set containing all possible ordered pairs (a,b) where a is from set A and b is from set B for any two sets.
 - ▶ AxB where $A = \{apple, banana, kiwi\}$ and $B = \{2, 4\}$. $AxB = \{apple, 2\}$, $\{apple, 4\}$, $\{banana, 2\}$, $\{banana, 4\}$, $\{kiwi, 2\}$, $\{kiwi, 4\}$

Sets: Review

- ▶ Unions: OR, ∪
- ▶ Intersections: AND, ∩
- ▶ Ordered/Unordered
- ▶ Complements (not inside) c, written A^c , for example.
- ▶ Subsets and proper subsets (contained within): \subset , \subseteq
- Cardinality (number of elements)

- How many hands are possible?
- Does order matter?
- How would it affect the sample space? Would we have more or fewer possible hands?
- Now, you can swap one card. How many ways could you do it? Does it matter what cards you have?

- ► How many hands are possible? this is the *universe* and we would calculate it using $\binom{10}{4}$
- Does order matter?
- How would it affect the sample space? Would we have more or fewer possible hands?
- Now, you can swap one card. How many ways could you do it? Does it matter what cards you have?

- ► How many hands are possible? this is the *universe* and we would calculate it using $\binom{10}{4}$
- ▶ Does order matter? No!
- How would it affect the sample space? Would we have more or fewer possible hands?
- ▶ Now, you can swap one card. How many ways could you do it? Does it matter what cards you have?

- ► How many hands are possible? this is the *universe* and we would calculate it using $\binom{10}{4}$
- Does order matter? No!
- How would it affect the sample space? Would we have more or fewer possible hands? Permutation vs Combination: we would have many more possible hands
- ▶ Now, you can swap one card. How many ways could you do it? Does it matter what cards you have?

- ► How many hands are possible? this is the *universe* and we would calculate it using $\binom{10}{4}$
- ▶ Does order matter? No!
- How would it affect the sample space? Would we have more or fewer possible hands? Permutation vs Combination: we would have many more possible hands
- Now, you can swap one card. How many ways could you do it? Does it matter what cards you have? You could trade in any one of your 4 cards. In this scenario it does not matter.

Suppose you have a deck of ten cards, numbered 1-10, and are dealt four cards. Answer the following questions, labelling them with the appropriate terms we've discussed so far: $\$

► Finally, what if I told you that all 4 of my cards were even numbered. Would you be surprised? How would you know whether to be surprised?

Suppose you have a deck of ten cards, numbered 1-10, and are dealt four cards. Answer the following questions, labelling them with the appropriate terms we've discussed so far: $\$

► Finally, what if I told you that all 4 of my cards were even numbered. Would you be surprised? How would you know whether to be surprised? We will discuss this, and many other interesting things!, in class! soon!