

Robot Physicist

Team = [Manoj Kumar, Michele Ceru, Phil Yeres]

Advisor = Kyle Cranmer

Project mentors

Professor Kyle Cranmer, NYU Physics

Brenden Lake, CDS (unofficial mentor)

The problem:

Particle physics experiments
are expensive to perform



The problem:

Particle physics experiments are expensive to perform

Physicists optimize their experiments by running simulated experiments



The problem:


Particle physics experiments are expensive to perform

Physicists optimize their experiments by running simulated experiments


Our goal: design a proof of concept “robot physicist” that efficiently finds optimal experiment configurations



The black box




The Robot Physicist



for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$

The Robot Physicist




for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\boldsymbol{\theta} = [\theta_1, \theta_2, \dots, \theta_m]$:


$$\theta_1, \Phi_j \longrightarrow [\hat{x}_{1,\theta_1} \quad \hat{x}_{2,\theta_1} \quad \cdots \quad \hat{x}_{n,\theta_1}]$$

The Robot Physicist




for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:




The Robot Physicist




for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:




The Robot Physicist




for a single configuration of the experimental settings, Φ_j ...

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:




The Robot Physicist




for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:





The Robot Physicist




for a single configuration of the experimental settings, $\Phi_j \dots$

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:






The Robot Physicist




for a single configuration of the experimental settings, Φ_j ...

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:







The Robot Physicist




for a single configuration of the experimental settings, Φ_j ...

we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:







The Robot Physicist



for a single configuration of the experimental settings, $\Phi_j \dots$


we use a list of possible values of the unobservable parameter, $\theta = [\theta_1, \theta_2, \dots, \theta_m]$:




Project phase 2: physics data

Goal: choose the energy settings for an electron-positron collider in order to best measure the Weinberg angle parameter

Theory




Experiment



Project phase 2: physics data

- Run a simulated experiment, generating data with an unknown distribution



- Estimate $P(\theta | x, \Phi)$
- Predict the experimental settings that measure the Weinberg angle with the least uncertainty

Thanks!