

# Computability and Complexity

## COSC 4200

### Deterministic Finite Automata II

# Binary Numbers

Each binary string  $x \in \{0, 1\}^*$  corresponds to a natural number  $\text{bin}(x) \in \mathbb{N}$ .

$x$	$\text{bin}(x)$		$x$	$\text{bin}(x)$
$\epsilon$	0			
0	0		1000	8
1	1		1001	9
10	2		1010	10
11	3		1011	11
100	4		1100	12
101	5		1101	13
110	6		1110	14
111	7		1111	15

# Inductive Definition of $\text{bin}(x)$

For any  $x \in \{0, 1\}^*$ , let  $\text{bin}(x)$  be the number  $x$  encodes in binary. This is defined inductively as follows.

- The base case is  $\text{bin}(\epsilon) = 0$ .
- For any  $x$ , assuming  $\text{bin}(x)$  has already been defined,

$$\text{bin}(x0) = 2\text{bin}(x),$$

$$\text{bin}(x1) = 2\text{bin}(x) + 1.$$

# Inductive Definition of $\text{bin}(x)$

For any  $x \in \{0, 1\}^*$ , let  $\text{bin}(x)$  be the number  $x$  encodes in binary. This is defined inductively as follows.

- The base case is  $\text{bin}(\epsilon) = 0$ .
- For any  $x$ , assuming  $\text{bin}(x)$  has already been defined,

$$\text{bin}(x0) = 2\text{bin}(x),$$

$$\text{bin}(x1) = 2\text{bin}(x) + 1.$$

For example,  $\text{bin}(100) = 4$  and

$$\text{bin}(1000) = 2\text{bin}(100) = 2 \cdot 4 = 8,$$

$$\text{bin}(1001) = 2\text{bin}(100) + 1 = 2 \cdot 4 + 1 = 9.$$

# Multiples of 3 in Binary

Show that

$$T = \{x \in \{0, 1\}^* \mid \text{bin}(x) \text{ is a multiple of } 3\}$$

is regular.

$x$	$\text{bin}(x)$		$x$	$\text{bin}(x)$
$\epsilon$	0			
0	0		1000	8
1	1		1001	9
10	2		1010	10
11	3		1011	11
100	4		1100	12
101	5		1101	13
110	6		1110	14
111	7		1111	15


# Multiples of 3 in Binary

Show that


$$T = \{x \in \{0,1\}^* \mid \text{bin}(x) \text{ is a multiple of } 3\}$$

is regular.


$x$	$\text{bin}(x)$		$x$	$\text{bin}(x)$
$\epsilon$	0			
0	0		1000	8
1	1		1001	9
10	2		1010	10
11	3		1011	11
100	4		1100	12
101	5		1101	13
110	6		1110	14
111	7		1111	15



$\epsilon$ , 0, 11, 110, 1001, 1100, 1111, ... are accepted








$$Q = \{0, 1, 2\} \quad \Sigma = \{0, 1\}$$

$$\delta : Q \times \Sigma \rightarrow Q$$

$$\delta(q, b) = 2q + b \pmod{3}$$



$$Q = \{0, 1, 2\} \quad \Sigma = \{0, 1\}$$

$$\delta : Q \times \Sigma \rightarrow Q$$

$$\delta(q, b) = 2q + b \pmod{3}$$

**Claim.** For all  $x \in \{0, 1\}^*$ ,  $\delta^*(0, x) = \text{bin}(x) \pmod{3}$ .

# Induction on Natural Numbers

**Goal:** Prove  $(\forall n \geq 0) P(n)$ .

**Inductive proof:**

- **Base case:** Prove  $P(0)$ .
- **Inductive hypothesis:** Assume that  $P(k)$  holds for some  $k \geq 0$ .  
**Inductive step:** Use the inductive hypothesis to prove that  $P(k + 1)$  also holds.

In the proof, we show that  $P(0)$  holds and

$$(\forall k \geq 0) P(k) \Rightarrow P(k + 1).$$

# Induction on Binary Strings

**Goal:** Prove  $(\forall x \in \{0, 1\}^*) P(x)$ .

**Inductive proof:**

- **Base case:** Prove  $P(\epsilon)$ .
- **Inductive hypothesis:** Assume that  $P(x)$  holds for some  $x \in \{0, 1\}^*$ .  
**Inductive step:** Use the inductive hypothesis to prove that  $P(x0)$  and  $P(x1)$  also hold.

In the proof, we show that  $P(\epsilon)$  holds and

$$(\forall x \in \{0, 1\}^*) P(x) \Rightarrow P(x0) \wedge P(x1).$$

**Claim.** For all  $x \in \{0, 1\}^*$ ,  $\delta^*(0, x) = \text{bin}(x) \pmod{3}$ .

**Proof.** (by induction on strings)

**Base case:**  $x = \epsilon$  :  $\text{bin}(\epsilon) = 0$ .

$$\delta^*(0, \epsilon) = 0 = \text{bin}(\epsilon).$$

**Inductive hypothesis:** Suppose the claim holds for some  $x \in \{0, 1\}^*$ .

$$\delta^*(0, x) = \text{bin}(x) \pmod{3}.$$

**Inductive hypothesis:** Suppose the claim holds for some  $x \in \{0, 1\}^*$ .

$$\delta^*(0, x) = \text{bin}(x) \pmod{3}.$$

**Inductive step:** We must prove that the claim holds for the extensions  $x0$  and  $x1$  of  $x$  as well:

$$\delta^*(0, x0) = \text{bin}(x0) \pmod{3}$$

$$\delta^*(0, x1) = \text{bin}(x1) \pmod{3}$$

**Inductive hypothesis:** Suppose the claim holds for some  $x \in \{0, 1\}^*$ .

$$\delta^*(0, x) = \text{bin}(x) \pmod{3}.$$

**Inductive step:** We must prove that the claim holds for the extensions  $x0$  and  $x1$  of  $x$  as well:

$$\delta^*(0, x0) = \text{bin}(x0) \pmod{3}$$

$$\delta^*(0, x1) = \text{bin}(x1) \pmod{3}$$

For convenience, we'll work with both statements at the same time:

$$\delta^*(0, xb) = \text{bin}(xb) \pmod{3} \text{ for } b \in \{0, 1\}$$



$$\delta^*(0,xb) = \delta(\delta^*(0,x),b)$$

definition of  $\delta^*$

$$\begin{aligned}
 \delta^*(0,xb) &= \delta(\delta^*(0,x),b) \\
 &= 2\delta^*(0,x) + b \pmod{3}
 \end{aligned}$$

definition of  $\delta^*$   
definition of  $\delta$

$$\begin{aligned}
\delta^*(0, xb) &= \delta(\delta^*(0, x), b) && \text{definition of } \delta^* \\
&= 2\delta^*(0, x) + b \pmod{3} && \text{definition of } \delta \\
&= 2(\text{bin}(x) \pmod{3}) + b \pmod{3} && \text{inductive hypothesis}
\end{aligned}$$

$$\begin{aligned}
\delta^*(0, xb) &= \delta(\delta^*(0, x), b) && \text{definition of } \delta^* \\
&= 2\delta^*(0, x) + b \pmod{3} && \text{definition of } \delta \\
&= 2(\text{bin}(x) \pmod{3}) + b \pmod{3} && \text{inductive hypothesis} \\
&= 2\text{bin}(x) + b \pmod{3}
\end{aligned}$$

$$\begin{aligned}
\delta^*(0, xb) &= \delta(\delta^*(0, x), b) && \text{definition of } \delta^* \\
&= 2\delta^*(0, x) + b \pmod{3} && \text{definition of } \delta \\
&= 2(\text{bin}(x) \pmod{3}) + b \pmod{3} && \text{inductive hypothesis} \\
&= 2\text{bin}(x) + b \pmod{3} \\
&= \text{bin}(xb) \pmod{3} && \text{definition of bin}
\end{aligned}$$

$$\begin{aligned}
\delta^*(0, xb) &= \delta(\delta^*(0, x), b) && \text{definition of } \delta^* \\
&= 2\delta^*(0, x) + b \pmod{3} && \text{definition of } \delta \\
&= 2(\text{bin}(x) \pmod{3}) + b \pmod{3} && \text{inductive hypothesis} \\
&= 2\text{bin}(x) + b \pmod{3} \\
&= \text{bin}(xb) \pmod{3} && \text{definition of bin}
\end{aligned}$$

Therefore  $\delta^*(0, xb) = \text{bin}(xb) \pmod{3}$  for both  $b \in \{0, 1\}$ .

$$\begin{aligned}
\delta^*(0, xb) &= \delta(\delta^*(0, x), b) && \text{definition of } \delta^* \\
&= 2\delta^*(0, x) + b \pmod{3} && \text{definition of } \delta \\
&= 2(\text{bin}(x) \pmod{3}) + b \pmod{3} && \text{inductive hypothesis} \\
&= 2\text{bin}(x) + b \pmod{3} \\
&= \text{bin}(xb) \pmod{3} && \text{definition of bin}
\end{aligned}$$

Therefore  $\delta^*(0, xb) = \text{bin}(xb) \pmod{3}$  for both  $b \in \{0, 1\}$ .

It follows by mathematical induction that the claim holds for all  $x \in \{0, 1\}^*$ . □

# Regular Operations

We will see that the regular languages are *closed under the regular operations* of union, concatenation, and star. This means that if  $A, B \in \text{REG}$ , then

$$A \cup B \in \text{REG},$$

$$A \cdot B \in \text{REG},$$

$$A^* \in \text{REG}.$$

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}$$

$$A \cdot B = \{xy \mid x \in A \text{ and } y \in B\}$$

$$A^* = \{x_1 x_2 \cdots x_k \mid x_1, x_2, \dots, x_k \in A \text{ and } k \geq 0\}$$



## Theorem

*If  $A, B \in \text{REG}$ , then  $A \cup B \in \text{REG}$ .*

**Proof.** Let  $A, B \in \text{REG}$ . Then there exist two DFAs  $M_A = (Q_A, \Sigma, \delta_A, q_A, F_A)$  and  $M_B = (Q_B, \Sigma, \delta_B, q_B, F_B)$  with  $L(M_A) = A$  and  $L(M_B) = B$ . We will use  $M_A$  and  $M_B$  to construct a new DFA  $M$  with  $L(M) = A \cup B$ .

## Theorem

If  $A, B \in \text{REG}$ , then  $A \cup B \in \text{REG}$ .

**Proof.** Let  $A, B \in \text{REG}$ . Then there exist two DFAs  $M_A = (Q_A, \Sigma, \delta_A, q_A, F_A)$  and  $M_B = (Q_B, \Sigma, \delta_B, q_B, F_B)$  with  $L(M_A) = A$  and  $L(M_B) = B$ . We will use  $M_A$  and  $M_B$  to construct a new DFA  $M$  with  $L(M) = A \cup B$ .

*Product Construction.* We let  $M = (Q, \Sigma, \delta, q_0, F)$  where

- $Q = Q_A \times Q_B = \{(p, q) \mid p \in Q_A, q \in Q_B\}$

## Theorem

If  $A, B \in \text{REG}$ , then  $A \cup B \in \text{REG}$ .

**Proof.** Let  $A, B \in \text{REG}$ . Then there exist two DFAs  $M_A = (Q_A, \Sigma, \delta_A, q_A, F_A)$  and  $M_B = (Q_B, \Sigma, \delta_B, q_B, F_B)$  with  $L(M_A) = A$  and  $L(M_B) = B$ . We will use  $M_A$  and  $M_B$  to construct a new DFA  $M$  with  $L(M) = A \cup B$ .

*Product Construction.* We let  $M = (Q, \Sigma, \delta, q_0, F)$  where

- $Q = Q_A \times Q_B = \{(p, q) \mid p \in Q_A, q \in Q_B\}$
- $\delta : Q \times \Sigma \rightarrow Q$  is defined by

$$\delta((p, q), a) = (\delta_A(p, a), \delta_B(q, a))$$

for all  $(p, q) \in Q$  and  $a \in \Sigma$ .

## Theorem

If  $A, B \in \text{REG}$ , then  $A \cup B \in \text{REG}$ .

**Proof.** Let  $A, B \in \text{REG}$ . Then there exist two DFAs  $M_A = (Q_A, \Sigma, \delta_A, q_A, F_A)$  and  $M_B = (Q_B, \Sigma, \delta_B, q_B, F_B)$  with  $L(M_A) = A$  and  $L(M_B) = B$ . We will use  $M_A$  and  $M_B$  to construct a new DFA  $M$  with  $L(M) = A \cup B$ .

*Product Construction.* We let  $M = (Q, \Sigma, \delta, q_0, F)$  where

- $Q = Q_A \times Q_B = \{(p, q) \mid p \in Q_A, q \in Q_B\}$
- $\delta : Q \times \Sigma \rightarrow Q$  is defined by

$$\delta((p, q), a) = (\delta_A(p, a), \delta_B(q, a))$$

for all  $(p, q) \in Q$  and  $a \in \Sigma$ .

- $q_0 = (q_A, q_B)$ .

## Theorem

If  $A, B \in \text{REG}$ , then  $A \cup B \in \text{REG}$ .

**Proof.** Let  $A, B \in \text{REG}$ . Then there exist two DFAs  $M_A = (Q_A, \Sigma, \delta_A, q_A, F_A)$  and  $M_B = (Q_B, \Sigma, \delta_B, q_B, F_B)$  with  $L(M_A) = A$  and  $L(M_B) = B$ . We will use  $M_A$  and  $M_B$  to construct a new DFA  $M$  with  $L(M) = A \cup B$ .

*Product Construction.* We let  $M = (Q, \Sigma, \delta, q_0, F)$  where

- $Q = Q_A \times Q_B = \{(p, q) \mid p \in Q_A, q \in Q_B\}$
- $\delta : Q \times \Sigma \rightarrow Q$  is defined by


$$\delta((p, q), a) = (\delta_A(p, a), \delta_B(q, a))$$

for all  $(p, q) \in Q$  and  $a \in \Sigma$ .


- $q_0 = (q_A, q_B)$ .
- $F = \{(p, q) \in Q \mid p \in F_A \text{ or } q \in F_B\}$

# Example - Product Construction

$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$




$$B = \{x11 \mid x \in \{0,1\}^*\}$$




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$A \cup B$ :




$$B = \{x11 \mid x \in \{0,1\}^*\}$$




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$A \cup B$ :




$$B = \{x11 \mid x \in \{0,1\}^*\}$$






# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :






# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$




$A \cup B$ :




# Example - Product Construction


$$A = \{x \in \{0,1\}^* \mid |x| \text{ is even}\}$$



$$B = \{x11 \mid x \in \{0,1\}^*\}$$



$A \cup B$ :



**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

Now suppose the claim holds for some string  $x$ . For the inductive step, we must show the claim also holds for  $x0$  and  $x1$ . For all  $b \in \{0, 1\}$ ,

**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

Now suppose the claim holds for some string  $x$ . For the inductive step, we must show the claim also holds for  $x0$  and  $x1$ . For all  $b \in \{0, 1\}$ ,

$$\delta^*(q_0, xb) = \delta(\delta^*(q_0, x), b)$$

**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

Now suppose the claim holds for some string  $x$ . For the inductive step, we must show the claim also holds for  $x0$  and  $x1$ . For all  $b \in \{0, 1\}$ ,

$$\begin{aligned}\delta^*(q_0, xb) &= \delta(\delta^*(q_0, x), b) \\ &= \delta((\delta_A^*(q_A, x), \delta_B^*(q_B, x)), b)\end{aligned}$$

**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

Now suppose the claim holds for some string  $x$ . For the inductive step, we must show the claim also holds for  $x0$  and  $x1$ . For all  $b \in \{0, 1\}$ ,

$$\begin{aligned}\delta^*(q_0, xb) &= \delta(\delta^*(q_0, x), b) \\ &= \delta((\delta_A^*(q_A, x), \delta_B^*(q_B, x)), b) \\ &= (\delta_A(\delta_A^*(q_A, x), b), \delta_B(\delta_B^*(q_B, x), b))\end{aligned}$$



**Claim.** For all  $x \in \Sigma^*$ ,  $\delta^*(q_0, x) = (\delta_A^*(q_A, x), \delta_B^*(q_B, x))$ .

*Proof.* The proof is by induction on strings.

For the base case, we have

$$\delta^*(q_0, \epsilon) = q_0 = (q_A, q_B) = (\delta^*(q_A, \epsilon), \delta^*(q_B, \epsilon)).$$

Now suppose the claim holds for some string  $x$ . For the inductive step, we must show the claim also holds for  $x0$  and  $x1$ . For all  $b \in \{0, 1\}$ ,

$$\begin{aligned}\delta^*(q_0, xb) &= \delta(\delta^*(q_0, x), b) \\ &= \delta((\delta_A^*(q_A, x), \delta_B^*(q_B, x)), b) \\ &= (\delta_A(\delta_A^*(q_A, x), b), \delta_B(\delta_B^*(q_B, x), b)) \\ &= (\delta_A^*(q_A, xb), \delta_B^*(q_B, xb)). \quad \square\end{aligned}$$

For any  $x \in \Sigma^*$ , we have

$$x \in L(M) \iff \delta^*(q_0, x) \in F$$

For any  $x \in \Sigma^*$ , we have

$$\begin{aligned} x \in L(M) &\Leftrightarrow \delta^*(q_0, x) \in F \\ &\Leftrightarrow (\delta_A^*(q_A, x), \delta_B^*(q_B, x)) \in F \end{aligned}$$

For any  $x \in \Sigma^*$ , we have

$$\begin{aligned}x \in L(M) &\Leftrightarrow \delta^*(q_0, x) \in F \\&\Leftrightarrow (\delta_A^*(q_A, x), \delta_B^*(q_B, x)) \in F \\&\Leftrightarrow \delta_A^*(q_A, x) \in F_A \text{ or } \delta_B^*(q_B, x) \in F_B\end{aligned}$$

For any  $x \in \Sigma^*$ , we have

$$\begin{aligned} x \in L(M) &\Leftrightarrow \delta^*(q_0, x) \in F \\ &\Leftrightarrow (\delta_A^*(q_A, x), \delta_B^*(q_B, x)) \in F \\ &\Leftrightarrow \delta_A^*(q_A, x) \in F_A \text{ or } \delta_B^*(q_B, x) \in F_B \\ &\Leftrightarrow x \in L(M_A) \text{ or } x \in L(M_B) \end{aligned}$$

For any  $x \in \Sigma^*$ , we have

$$\begin{aligned}x \in L(M) &\Leftrightarrow \delta^*(q_0, x) \in F \\&\Leftrightarrow (\delta_A^*(q_A, x), \delta_B^*(q_B, x)) \in F \\&\Leftrightarrow \delta_A^*(q_A, x) \in F_A \text{ or } \delta_B^*(q_B, x) \in F_B \\&\Leftrightarrow x \in L(M_A) \text{ or } x \in L(M_B) \\&\Leftrightarrow x \in L(M_A) \cup L(M_B) = A \cup B.\end{aligned}$$

Therefore  $L(M) = A \cup B$ , so  $A \cup B$  is regular. □

# Closure Under Complementation

The *complement* of a language  $A$  is

$$A^c = \{x \in \Sigma^* \mid x \notin A\}.$$

## Theorem

*If  $A$  is regular, then  $A^c$  is regular.*

# Closure Under Complementation

The *complement* of a language  $A$  is

$$A^c = \{x \in \Sigma^* \mid x \notin A\}.$$

## Theorem

*If  $A$  is regular, then  $A^c$  is regular.*

**Proof.** Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA for  $A$ . Define a new DFA


$$M' = (Q, \Sigma, \delta, q_0, Q - F)$$

that flips the accepting and nonaccepting states of  $M$ . Then  $L(M') = L(M)^c = A^c$ . □




## Example - Closure Under Complementation

$C = \{w \in \{0,1\}^* \mid w \text{ contains } 010 \text{ as a substring}\}.$



$C^c = \{w \in \{0,1\}^* \mid w \text{ does not contain } 010 \text{ as a substring}\}.$



# Closure Under Intersection

## Theorem


*If  $A$  and  $B$  are regular languages, then  $A \cap B$  is regular.*

# Closure Under Intersection

## Theorem


*If  $A$  and  $B$  are regular languages, then  $A \cap B$  is regular.*

**Proof.** This follows from the previous two theorems. Observe that  $A \cap B = (A^c \cup B^c)^c$ .



The yellow region is  $A^c \cup B^c$ .

The complement of the yellow region is  $A \cap B$ .




The yellow region is  $A^c \cup B^c$ .

The complement of the yellow region is  $A \cap B$ .

Formally, we have

$$(A^c \cup B^c)^c =$$




The yellow region is  $A^c \cup B^c$ .

The complement of the yellow region is  $A \cap B$ .

Formally, we have

$$(A^c \cup B^c)^c = (A^c)^c \cap (B^c)^c \quad \text{De Morgan's Law}$$



The yellow region is  $A^c \cup B^c$ .


The complement of the yellow region is  $A \cap B$ .

Formally, we have

$$\begin{aligned}(A^c \cup B^c)^c &= (A^c)^c \cap (B^c)^c && \text{De Morgan's Law} \\ &= A \cap B && \text{Double Complement Law.}\end{aligned}$$

Now we apply closure properties:

- Since  $A$  and  $B$  are regular,  $A^c$  and  $B^c$  is also regular.



The yellow region is  $A^c \cup B^c$ .


The complement of the yellow region is  $A \cap B$ .

Formally, we have

$$\begin{aligned}(A^c \cup B^c)^c &= (A^c)^c \cap (B^c)^c && \text{De Morgan's Law} \\ &= A \cap B && \text{Double Complement Law.}\end{aligned}$$

Now we apply closure properties:

- Since  $A$  and  $B$  are regular,  $A^c$  and  $B^c$  is also regular.
- Therefore  $A^c \cup B^c$  is regular.



The yellow region is  $A^c \cup B^c$ .

The complement of the yellow region is  $A \cap B$ .

Formally, we have

$$\begin{aligned}(A^c \cup B^c)^c &= (A^c)^c \cap (B^c)^c && \text{De Morgan's Law} \\ &= A \cap B && \text{Double Complement Law.}\end{aligned}$$

Now we apply closure properties:

- Since  $A$  and  $B$  are regular,  $A^c$  and  $B^c$  is also regular.
- Therefore  $A^c \cup B^c$  is regular.
- Finally,  $(A^c \cup B^c)^c$  is regular.  $\square$



## Theorem

*If  $A$  and  $B$  are regular languages, then  $A \cap B$  is regular.*

We could also prove this by modifying the accepting states in the product construction.

## Theorem

*If  $A$  and  $B$  are regular languages, then  $A \cap B$  is regular.*

We could also prove this by modifying the accepting states in the product construction.

Change the “or” in  $F$  to an “and”:

$$\begin{aligned} F &= \{(p, q) \in Q \mid p \in F_A \text{ or } q \in F_B\} \\ &= F_A \times F_B \end{aligned}$$